

Daily physical activity and COPD

Citation for published version (APA):

Vaes, A. W. (2015). *Daily physical activity and COPD*. Maastricht University.

Document status and date:

Published: 01/01/2015

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Valorization addendum



Relevance

COPD is a highly prevalent disease, which negatively affects patients daily functioning. It is known that COPD patients experience problems with performance of activities of daily life (ADLs) and that the prevalence of physical inactivity levels is remarkably higher compared to healthy controls. These lower levels of physical activity have a tremendous impact on many health related outcomes and progression of the disease. Since physical activity is an important modifiable risk factor, it can therefore be an important disease management target in COPD.

Although physical activity has been studied frequently in COPD, less is known about ADLs. Moreover, strategies for improving physical activity are rarely studied. The current thesis investigated the impact of COPD on the performance of ADLs, strategies to improve functional exercise performance, symptom burden and physical activity level and the impact of longitudinal changes in physical activity level. This thesis provided new evidence in this field of expertise:

First, our findings showed that patients with COPD experience more problems during the performance of domestic ADLs, as indicated by a higher task-related oxygen uptake and ventilation and a higher degree of dyspnea and fatigue. These differences were already present in the earliest stages of the disease. Furthermore, patients in different BMI categories (underweight, normal, overweight or obese BMI) experience similar functional limitations during the performance of domestic ADLs.

Second, our data indicate that walking aids can improve functional exercise performance not only indoors, but also during outdoor use. Though, clinicians should base the selection of a walking aid preferably on an exercise field test instead of an indoor 6MWT, since test can be used to acquire more patient-meaningful information on walking ability and walking pattern and patients reported to use their walking aid mostly during outdoor activities. Moreover, patients' needs and daily use should also be taken into consideration.

Third, we demonstrated that a longitudinal decline in physical activity is significantly more common in subjects with COPD compared to healthy subjects, and that determinants of these longitudinal changes in physical activity are different. Moreover, a longitudinal decline to a low physical activity level was associated with an increased all-cause mortality risk. More important, findings suggest that once patients with COPD become physically low active, the high mortality risk cannot be reversed by increasing activity levels, showing the importance of encouraging an active lifestyle in patients with COPD.

Fourth, we showed the promising effects of activity monitor-based counseling on improvement of physical activity and health-related outcomes in patients with DMII. It can be assumed that activity monitor-based interventions in patients with COPD and CHF will have similar beneficial effects, however, more research is needed.