

# Effect of Intradialytic Aerobic Exercise among Hemodialysis Patients in the United Arab Emirates on Clinical Outcomes, Barriers to Physical Activity, and Quality of Life

Citation for published version (APA):

Salhab, N. (2019). *Effect of Intradialytic Aerobic Exercise among Hemodialysis Patients in the United Arab Emirates on Clinical Outcomes, Barriers to Physical Activity, and Quality of Life*. [Doctoral Thesis, Maastricht University]. ProefschriftMaken Maastricht. <https://doi.org/10.26481/dis.20191002ns>

## Document status and date:

Published: 01/01/2019

## DOI:

[10.26481/dis.20191002ns](https://doi.org/10.26481/dis.20191002ns)

## Document Version:

Publisher's PDF, also known as Version of record

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

Download date: 31 Mar. 2023

---

## VALORIZATION

## VALORIZATION

Chronic kidney disease (CKD) is a global health concern (Eckardt et al., 2013) with a prevalence of 13.4% worldwide (Hill et al., 2016). In 2017, it was estimated that 3.2 million patients globally were undergoing dialysis; this number was at 3 million in 2016, thus presenting a 6% annual growth rate (Fresenius Medical Care, 2017). In addition, CKD patients have increased mortality rate compared to the healthy population (Johansen et al., 2000). In 2015, the United Arab Emirates (UAE) had 1750 hemodialysis (HD) patients with an annual mortality rate of 8.5% (AlSahow et al., 2016). Furthermore, in 2018, home hemodialysis became an option for elderly, bed ridden Emirati patients (The national, 2018). All of this projects an increase in the economic burden of end stage renal disease.

Aside from the economical burden, dialysis patients present a social burden when it comes to their decreased quality of life (QOL) and their sedentary lifestyle. HD patients have decreased QOL as compared to healthy adults (Avramovic & Stefanovic, 2012). Furthermore, QOL tends to deteriorate with the advancement in CKD stages (Aggarwal et al., 2016). The situation in the UAE is not different and chronically ill patients have lower QOL scores when compared to healthy individuals (Ayoub & Hijjaz, 2013). Looking at HD patient's activity level, it was reported to be less than that of healthy sedentary controls (Johansen et al., 2000). As QOL and sedentarism could predict mortality rate (O'hare et al., 2003; Stack et al., 2005; Tentori et al., 2010; Avramovic & Stefanovic, 2012), there is an urge to find an adjunctive therapy to ease the social and economical pressure of HD patients. This thesis focused on IDE as a complementary therapy for HD patients that could impact on this social and economical burden.

Today, public health awareness campaigns are highlighting the importance of exercise in a healthy lifestyle, whether you are a healthy individual or a chronically ill patient. The global attention of stakeholders and health care professionals is focused on how to improve patients' health outcomes, increase the quality of life and decrease the health care costs.

Our project supported this global perspective by being a pioneer in exploring the effect of IDE on various clinical and social outcomes in HD patients in the UAE. After the completion of our study, we could say that intradialytic aerobic exercise (IDE) could be a safe option to enhance treatment from a different angle; this could be

possible with a minimal primary investment that covers the equipment's cost and the salary of a coordinator to assist the patients with IDE in collaboration with the nursing staff. Moreover, the initiation of IDE will require a primary assessment of the patient by his nephrologist, and the education for the staff about the benefits, and proper protocol of IDE.

Introducing IDE in HD population is a challenge, especially that many patients think of it as adding a burden to their situation. Thus, rises the need of more research focusing on the cost effectiveness of IDE, and on how it could alleviate the health and economic burden on the patient, the society, and the health care system. Cost effectiveness could be evaluated based on the hospitalization rate, the intake of phosphate binders, and maybe the frequency and duration of the weekly dialysis sessions. With the advancement of new technologies such as fitness trackers, step counter watches, and renal food applications, the motivation, assessment and accurate follow-up are now easier. In addition to the readily available, user friendly, low cost bio-impedance analyzers that offers an accurate body composition analysis. It is worth mentioning that a renal food application culturally specific for the UAE "KELA.ae" (developed by Dr. Mirey Karavetian and funded by Zayed University) has been launched in 2017. This application would not only benefit UAE dialysis patients, but also researchers in that field. It could minimize the cost of research that requires patient education on the renal diet. Such applications would also complement an IDE program, as nutrition and exercise goes together.

Besides, this dissertation highlighted the high prevalence of malnutrition among HD patients in the UAE. Therefore, future research should focus on IDE and malnutrition to position IDE as a preventive measure for the commonly reported deterioration of the nutritional status of HD patients over time (Karavetian et al., 2016).

In the UAE, 40% of the hemodialysis patients are Emirati citizens (AlSahow et al., 2016), which means that the government covers their health insurances. Hence, an initiative from the government side to support IDE programs might lessen the cost burden on the healthcare system. Also, we hope to see in the future more organizational efforts, whether from hospitals or private dialysis centers, on this matter. Organizations could take the initiative and implement IDE. For instance, a pediatric dialysis unit in Argentina is offering an amazing physical activity program for their patients. A small video about their work could be seen on <https://www.youtube.com/watch?v=4mPcKXL5B70&t=89s>. "Plagiarism" in this sense would not be a crime!