

Diagnosis and minimally invasive treatment of chronic discogenic low back pain

Citation for published version (APA):

Kallewaard, J. W. (2019). *Diagnosis and minimally invasive treatment of chronic discogenic low back pain*. [Doctoral Thesis, Maastricht University]. Maastricht University. <https://doi.org/10.26481/dis.20191106jk>

Document status and date:

Published: 06/11/2019

DOI:

[10.26481/dis.20191106jk](https://doi.org/10.26481/dis.20191106jk)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Valorization

Discogenic Low Back Pain: "burden for patient and society"

Valorization is a broad concept encompassing knowledge transfer from the research sector to other sectors for personal, social and economic value.¹

Low back pain is a vast problem worldwide causing a huge burden for the patient and for society, that is still growing. It is the main cause of years lived with disability in the western world.²

Chronic discogenic low back pain (CD-LBP) is defined as pain originating from the intervertebral disc, lasting for more than 3 months. The complaints of patients with CD-LBP result in medical consumption, absenteeism from work and disability. The recently published results from the Global Burden Group showed that chronic low back pain is a worldwide problem that is still growing.³ The growth is attributed to aging and the growing world population. The incidence of CD-LBP is estimated at 40% of chronic low back pain.⁴

The quality of life of patients with low back pain is reduced. In a study from our group evaluating the burden and costs of CD-LBP in patients selected for participation in a randomized controlled trial on a new treatment option, almost half of the patients with CD-LBP, (46%), reported severe pain (>7 of NRS 0-10), and 54% suffered moderate pain.⁵ This is in comparison with the findings of a European prevalence study in chronic pain patients⁶, that showed 34% have severe pain and 66% have moderate pain.

In 2007 a Chinese research group published a spectacular decrease in pain and improvement of quality of life, after an intradiscal methylene blue injection in a highly selected group of patients with CD-LBP.⁷ The publication reported in the methylene blue group, a mean reduction in VAS of 52.5%, a 35.8% mean reduction in the Oswestry disability scores and a 91.6% satisfaction rate compared with 0.70%, 1.68%, and 14.3%, respectively, in the placebo treatment group. These astonishing good results gave rise to skepticism, and if the results of that Chinese study would be true, this intervention would revolutionize the treatment of low back pain, rendering spinal surgery for back pain essentially obsolete.⁸ As with any treatment, the results of this study need to be replicated.

What did we find in our studies?

Diagnosing discogenic pain remains a difficult and controversial discussion. Since physical examination and radiology will only give an indication of discogenic pain, pressure-controlled discography remains a very useful test in the workup of diagnosing discogenic low back pain, in spite of the possible disadvantage of accelerating disc generation.⁹ In

the current thesis, we refuted the hypothesis that outcomes of discography are flawed by the transfer of pressure to adjacent discs.

The promising results of intradiscal methylene blue injection could not be reproduced in our multicenter RCT, which copied exactly the protocol of the Peng et al.¹⁰ No significant differences in outcome were found between the methylene blue and the placebo control group. Thus, the use of intradiscal methylene blue injections could not be supported.¹¹ At the moment only intradiscal biacuplasty is recommended for the minimally invasive treatment of CD-LBP. Biacuplasty is a cooled radiofrequency technique aiming to destroy painful nociceptors in the posterior part of the disc.¹²⁻¹⁴

When all therapies such as conservative therapy, medication, minimally invasive treatments fail and surgery is not an option, neurostimulation could be considered. The sustained effect over 12-months of high-frequency 10kHz stimulation of the spinal cord in chronic low back pain was documented in the literature in patients with suspected CD-LBP.¹⁵ Our prospective study with DRG L2 stimulation in a highly selected group of patients with CD-LBP showed a promising result after 12 months.¹⁶ Further studies are needed to define if there is a place for neurostimulation in CD-LBP in non-operated patients and to find what place it should have in the treatment algorithm of CD-LBP.

References

- 1 Finne H, Day A, Piccaluga A, Spithoven A, Walter P, Wellen D. A Composite Indicator for Knowledge Transfer. *Report from the European Commission's Expert Group on Knowledge Transfer Indicators*. Trondheim2011.
- 2 Itz CJ, Ramaekers BL, van Kleef M, Dirksen CD. Medical specialists care and hospital costs for low back pain in the Netherlands. *Eur J Pain*. 2017;21:705-715.
- 3 G. B. D. Disease Injury Incidence Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392:1789-1858.
- 4 Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G, Bogduk N. The prevalence and clinical features of internal disc disruption in patients with chronic low back pain. *Spine*. 1995;20:1878-1883.
- 5 Geurts JW, Willems PC, Kallewaard JW, van Kleef M, Dirksen C. The Impact of Chronic Discogenic Low Back Pain: Costs and Patient's Burden. *Pain Res Manag*. 2018;2018:4696180.
- 6 Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D. Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *Eur J Pain*. 2006;10:287-333.
- 7 Peng B, Zhang Y, Hou S, Wu W, Fu X. Intradiscal methylene blue injection for the treatment of chronic discogenic low back pain. *Eur Spine J*. 2007;16:33-38.
- 8 Bogduk N. A cure for back pain? *Pain*. 2010;149:7-8.
- 9 Carragee EJ, Don AS, Hurwitz EL, Cuellar JM, Carrino JA, Herzog R. 2009 ISSLS Prize Winner: Does discography cause accelerated progression of degeneration changes in the lumbar disc: a ten-year matched cohort study. *Spine (Phila Pa 1976)*. 2009;34:2338-2345.
- 10 Peng B, Pang X, Wu Y, Zhao C, Song X. A randomized placebo-controlled trial of intradiscal methylene blue injection for the treatment of chronic discogenic low back pain. *Pain*. 2010;149:124-129.
- 11 Kallewaard JW, Wintraecken VM, Geurts JW, Willems PC, van Santbrink H, Terwiel CTM, van Kleef M, van Kuijk SMJ. A multicenter randomized controlled trial on the efficacy of intradiscal methylene blue injection for chronic discogenic low back pain: the IMBI study. *Pain*. 2019;160:945-953.
- 12 Kapural L, Vrooman B, Sarwar S, Krizanac-Bengez L, Rauck R, Gilmore C, North J, Mekhail N. Radiofrequency intradiscal biacuplasty for treatment of discogenic lower back pain: a 12-month follow-up. *Pain Med*. 2015;16:425-431.
- 13 Desai MJ, Kapural L, Petersohn JD, Vallejo R, Menzies R, Creamer M, Gofeld M. A Prospective, Randomized, Multicenter, Open-label Clinical Trial Comparing Intradiscal Biacuplasty to Conventional Medical Management for Discogenic Lumbar Back Pain. *Spine (Phila Pa 1976)*. 2016;41:1065-1074.
- 14 Kapural L, Ng A, Dalton J, Mascha E, Kapural M, de la Garza M, Mekhail N. Intervertebral disc biacuplasty for the treatment of lumbar discogenic pain: results of a six-month follow-up. *Pain Med*. 2008;9:60-67.

- 15 Al-Kaisy A, Palmisani S, Smith T, Harris S, Pang D. The use of 10-kilohertz spinal cord stimulation in a cohort of patients with chronic neuropathic limb pain refractory to medical management. *Neuromodulation*. 2015;18:18-23; discussion 23.
- 16 Kallewaard JW, Nijhuis H, Huygen F, Wille F, Zuidema X, van de Minkelis J, Raza A. Prospective Cohort Analysis of DRG Stimulation for Failed Back Surgery Syndrome Pain Following Lumbar Discectomy. *Pain Pract*. 2019;19:204-210.