

Falls among older people

Citation for published version (APA):

McCrum, C. (2018). Falls among older people: are intervention duration and specificity the keys to making a difference? *BMJ (e)*, 361, [k2213]. <https://doi.org/10.1136/bmj.k2213>

Document status and date:

Published: 25/05/2018

DOI:

[10.1136/bmj.k2213](https://doi.org/10.1136/bmj.k2213)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



LETTERS

PREVENTING FALLS IN OLDER PEOPLE

Falls among older people—are intervention duration and specificity the keys to making a difference?

Christopher McCrum *human movement scientist and PhD candidate*

Maastricht University Medical Centre, 6200 MD Maastricht, Netherlands

As Oliver says,¹ strength and balance training reduces falls in older people, with greater benefits when the weekly training exceeds three hours.² Exposing enough people at risk of falls to interventions for enough time to yield population level improvements is one of the main challenges, so the required exposure time for interventions is an essential consideration.

Another intervention that may be more time efficient is reactive (or perturbation based) balance training.³ Although few randomised controlled trials have been conducted, evidence indicates that just a single session or a few training sessions, including repeated slips or trips, lead to long term benefits, reflected in reduced incidence of falls and improved stability control.^{4,5} The driving theory is that this training hones the skills required to avoid falls after balance disturbances, such as rapid reactive stepping, counter rotations of the upper body and limbs, and grasping reactions.^{3,6} We do not expect athletes to win Olympic medals without practising the skills of their sports, so why should we expect older adults to fall less often without practising the skills required for avoiding falls?

Implementing reactive balance training is more challenging than traditional exercise and should be conducted with care, but recent work has provided examples of how this can be achieved.⁶ Further research is required, but with the potential for similar

or greater reductions in falls in a fraction of the time compared with current practice, reactive balance training provides opportunities to tackle some of the current challenges facing our healthcare systems regarding falls.

Competing interests: None declared.

Full response at: <https://www.bmj.com/content/361/bmj.k1655/rr-0>.

- 1 Oliver D. David Oliver: Falls in older people—can we really make a difference? *BMJ* 2018;361:k1655. doi:10.1136/bmj.k1655 29666147
- 2 Sherrington C, Michaleff ZA, Fairhall N, et al. Exercise to prevent falls in older adults: an updated systematic review and meta-analysis. *Br J Sports Med* 2017;51:1750-8. doi:10.1136/bjsports-2016-096547 27707740
- 3 Mansfield A, Wong JS, Bryce J, Knorr S, Patterson KK. Does perturbation-based balance training prevent falls? Systematic review and meta-analysis of preliminary randomized controlled trials. *Phys Ther* 2015;95:700-9. doi:10.2522/ptj.20140090 25524873
- 4 Pai YC, Bhatt T, Yang F, Wang E. Perturbation training can reduce community-dwelling older adults' annual fall risk: a randomized controlled trial. *J Gerontol A Biol Sci Med Sci* 2014;69:1586-94. doi:10.1093/gerona/glu087 24966227
- 5 Epro G, Mierau A, McCrum C, Leyendecker M, Bruggemann GP, Karamanidis K. Retention of gait stability improvements over 1.5 years in older adults: effects of perturbation exposure and triceps surae neuromuscular exercise. *J Neurophysiol* 2018; (forthcoming). doi:10.1152/jn.00513.2017 29537914
- 6 Gerards MHG, McCrum C, Mansfield A, Meijer K. Perturbation-based balance training for falls reduction among older adults: Current evidence and implications for clinical practice. *Geriatr Gerontol Int* 2017;17:2294-303. doi:10.1111/ggi.13082 28621015

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://group.bmj.com/group/rights-licensing/permissions>