

Resolving the role of genetic defects and mtDNA copy number in mitochondrial disease and development

Citation for published version (APA):

Kamps, R. (2020). *Resolving the role of genetic defects and mtDNA copy number in mitochondrial disease and development*. [Doctoral Thesis, Maastricht University]. Maastricht University. <https://doi.org/10.26481/dis.20201120rk>

Document status and date:

Published: 01/01/2020

DOI:

[10.26481/dis.20201120rk](https://doi.org/10.26481/dis.20201120rk)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 01 Apr. 2023

Valorisation

Mitochondrial diseases are the most common genetic metabolic diseases, affecting approximately 1 in 5,000 individuals [1]. Mitochondrial diseases are a clinically and genetically heterogeneous group of disorders, which generally manifest in tissues or organs with a high-energy requirement [2]. Due to this complexity and the lack of proper genotype-phenotype correlations, at the start of the work performed in this thesis, a genetic diagnosis was obtained in only 25% of patients by sequential analysis of the mtDNA and single candidate genes. The application of next-generation sequencing as a research tool to first sequence gene panels and later all genes (the exome) has increased this figure considerably, which has facilitated a rapid introduction in diagnostics and clinical care [3]. The valorisation paragraph will deal with the introduction of NGS in routine genetic diagnostics.

Next-generation sequencing: from research to diagnostics

A genetic diagnosis is important for counseling patient and relatives, for preventing the transmission, for providing a more accurate prognosis and in some cases for offering therapeutic interventions. During the past years, NGS technology has developed as the preferred tool for identifying the genetic defects in patients with clinically and genetically heterogeneous diseases (**Chapter 2**). NGS has contributed in identifying the genetic cause in either the mtDNA (20%) or nDNA (WES; 49%), leading to an overall diagnostic yield of 69% [4]. Given the impressive increase not only in the percentage of patients, in which a genetic diagnosis is established (from 25% to 69%), but also in the speed (from years to months) and in costs (expensive Sanger Sequencing), a rapid introduction of NGS in routine genetic diagnosis of mitochondrial disease was warranted and has been established in the last couple of years. NGS costs rapidly declined and the capacity of platforms as the Illumina NovaSeq 6000 increased tremendously. For example 200 exomes can be sequenced in a single sequencing run. Crucial in this respect was also the development of novel improved bioinformatics tools to predict variant pathogenicity and the availability of superior databases with information on exomes (Exac) and genetic variants (ClinVar). The adage will be NGS first and guided by the genes and variants found, specific clinical and laboratory investigations can be carried out to unambiguously link the genetic defect(s) to the clinical manifestations. The laboratory findings may also be important, in case

functional validation is required. Our results show a few key points for proper introduction into the clinic:

1. NGS of both the mtDNA and the whole exome

It is essential investigate both the mtDNA and the exome to identify the genetic cause in mitochondrial disease [4-6]. Although the mtDNA can be analysed in the exome DNA, the coverage is insufficient to detect the necessary low level mutation loads in blood DNA, which might indicate a higher mutation load and follow-up in affected tissues. NGS of the mtDNA allows detection of point mutations and rearrangements alike with immediate characterization of the breakpoints. Only the mtDNA copy number has to be established by a separate test.

2. Novel mitochondrial disease genes

Novel genes are still being identified as a possible cause of mitochondrial disease. These are either genes, in which no genetic defect has been found so far, or genes, in which defects have been found in apparently non-mitochondrial disease. This implies that gene panel-based approaches fall short, as the mitochondrial panel is still not complete. Complete analysis of all genes in the exome (open exome) [4] is preferable to start with to identify all relevant genetic defects in the shortest timeframe and at the lowest cost.

3. Overlap between mitochondrial and other neuromuscular disorders

Our results demonstrate that many neuromuscular diseases exist, in which a mitochondrial defect can be the cause in the minority of cases. In case of gene panels, this would imply sequential analysis of multiple, and probably partly overlapping gene panels. Again, it is preferable to start with an open exome [4], as this would allow an unbiased identification of the underlying genetic causes.

4. De novo mutations

As we detected de novo mutations in the clinical exome, it would be preferable to include immediately the parents for whole exome sequencing [7] and analyze the data, using an autosomal recessive and *de novo* dominant disease model.

5. Multigenic disease

Especially in clinically heterogeneous disease and when consanguinity is involved, the option of having multiple genes causing the disease in the

patient, should be investigated. This can be done by investigating all gene defects, which could be potentially contribute to the clinical phenotype. One should not address all symptoms to the first gene identified. As this has been done in the past, it is clear that genotype-phenotype correlations from literature [8], when based on few patients and single gene analysis, should be considered with care.

Our proposed diagnostic strategy for most mitochondrial diseases will be NGS of the mtDNA, followed by whole exome sequencing of patients and parents, followed by analysis of all genes, using an autosomal recessive or *de novo* dominant genetic model. In case of new genes or unclassified variants, functional follow-up studies should be offered to demonstrate pathogenicity. In case an OXPHOS deficiency is observed in patient-derived fibroblasts, this can be done straightforward by complementation assays. If this is not the case, animal studies might provide the answer, which will often go beyond the capabilities of routine diagnostic labs. An *in vivo* model as zebrafish could fill this gap, but standardization, validation and high-throughput processing is required to keep-up with the rapid generation of unclassified variants and novel genes by WES and WGS these days.

As the combined approach will identify at max 69% of the genetic causes, still 31% of the patients will remain without a genetic diagnosis. As these are carefully selected patients from our cohort, it is likely that there will be technical issues, prohibiting the identification of the underlying defects. Applying WGS could solve the issue, but routine use is still limited by the financial costs and the complexity of data interpretation. Rapid and life-saving genetic diagnosis in less than 24 hours using rapid WGS (rWGS) is possible and already offered for early intervention cases within paediatric Intensive Care Units (ICUs) as discussed in **Chapter 7**. Introducing rWGS showed a diagnostic yield of 42 to 57% in serious ill infants in ICUs, and changes in medical management in 30 to 72%, and finally altered outcomes in 24 to 34% of the cases [9]. Sequencing the transcriptome by RNA-Seq seems at this moment a more realistic addition to WES to solve the remaining cases. RNA-Seq allowed analysis of the transcriptome at an unprecedented depth and reported the identification of causative, non-exonic regulatory variants in an additional 10% diagnostic yield of mitochondrial patients using RNA-Seq [10]. In another study of rare muscular disorders this was 35% yield, which clearly shows

the added value of this RNA-Seq approach [11]. It is beyond any doubt, that WGS and RNA-Seq will bring the paradigm shift started by WES to the next level: for a genetic diagnosis, geneticists and expert clinicians will no longer need to narrow down the differential clinical diagnosis to request sequencing of a specific gene or sets of genes, but all genetic information will be available at once from start. The results of this genetic analysis will directly guide the diagnostic patient evaluation, moving towards personalized patient care and treatment.

Gene-based treatment options

The identification of the gene defect in the patients has unfortunately not led to the anticipated major breakthrough in novel treatments. Only in few cases, a gene specific-treatment is possible and early start with these treatments prevents symptoms from occurring and can be lifesaving, as for example administration of thiamine to patients with fatal Leigh Syndrome due to *SLC19A3* mutations [5]. In case of *SLC25A32* mutations riboflavin is also the preferred treatment [12]. Our study revealed that *SLC25A32* patients improved when treated with riboflavin as a vitamin B2 precursor of FAD which will compensate the lack of FAD functioning as a crucial co-factor in the mitochondrial transport chain [13]. In these cases, the genetic defect directly revealed a successful treatment option. In other cases, clinical improvement by specific treatments at a moment when the genetic defect was not known, revealed when the genetic defect was found, which patients might benefit from a comparable treatment. We demonstrated that again riboflavin could be beneficial for patients in case of *ACAD9* defects [14]. Since increasing the levels of the riboflavin restored Complex I activity by compensating a disturbed *ACAD9* and FAD binding or by stimulating Complex I assembly factors. However, not all *ACAD9* patients are treatable with riboflavin [15]. A recent study on *ACAD9* disease-causing variants in 70 patients showed improvement of at least 20/31 (65%) *ACAD9* deficient patients to riboflavin and 11/31 (35%) patients had no effect, although 20 patients were not treated and additional data about the treatment were missing for approximately 19 patients in this cohort. However, this justifies administration of riboflavin to every *ACAD9* patient. Given the high frequency of *ACAD9* deficiency, we also propose that early riboflavin administration for phenotypically consistent patients whilst their genetic investigations are still underway [16]. Another, successful treatment of infantile-onset *ACAD9*-related cardiomyopathy was achieved by a combination of sodium

pyruvate, beta-blocker, and coenzyme Q₁₀ [17]. Finally, also a high fat diet resulted in clinical improvement in a patient with a OXPHOS Complex I deficiency due to a mutation in the *TMEM126B* gene [18].

Conclusively, still this is a minority, and gene and pathway based treatment strategies might fall short. More general, gene-correction/editing (CRISPR/Cas9) or cell-based strategies are more promising as a general treatment strategy, complemented by classical compound-screening strategies of the pharmaceutical industry.

References

1. Gorman, G.S., et al., *Prevalence of nuclear and mitochondrial DNA mutations related to adult mitochondrial disease*. *Ann Neurol*, 2015. **77**(5): p. 753-9.
2. Gorman, G.S., et al., *Mitochondrial diseases*. *Nat Rev Dis Primers*, 2016. **2**: p. 16080.
3. Neveling, K., et al., *A post-hoc comparison of the utility of sanger sequencing and exome sequencing for the diagnosis of heterogeneous diseases*. *Hum Mutat*, 2013. **34**(12): p. 1721-6.
4. Theunissen, T.E.J., et al., *Whole Exome Sequencing Is the Preferred Strategy to Identify the Genetic Defect in Patients With a Probable or Possible Mitochondrial Cause*. *Front Genet*, 2018. **9**: p. 400.
5. Gerards, M., et al., *Exome sequencing reveals a novel Moroccan founder mutation in SLC19A3 as a new cause of early-childhood fatal Leigh syndrome*. *Brain*, 2013. **136**(Pt 3): p. 882-90.
6. Kamps, R., et al., *Genetic defects in mtDNA-encoded protein translation cause paediatric, mitochondrial cardiomyopathy with early-onset brain disease*. *Eur J Hum Genet*, 2018. **26**(4): p. 537-551.
7. Sallevelt, S., et al., *Preimplantation genetic diagnosis for mitochondrial DNA mutations: analysis of one blastomere suffices*. *J Med Genet*, 2017. **54**(10): p. 693-697.
8. Theunissen, T.E.J., et al., *Rapid Resolution of Blended or Composite Multigenic Disease in Infants by Whole-Exome Sequencing*. *J Pediatr*, 2017. **182**: p. 371-374 e2.
9. Clark, M.M., et al., *Diagnosis of genetic diseases in seriously ill children by rapid whole-genome sequencing and automated phenotyping and interpretation*. *Sci Transl Med*, 2019. **11**(489).
10. Cummings, B.B., et al., *Improving genetic diagnosis in Mendelian disease with transcriptome sequencing*. *Sci Transl Med*, 2017. **9**(386).
11. Kremer, L.S., S.B. Wortmann, and H. Prokisch, *"Transcriptomics": molecular diagnosis of inborn errors of metabolism via RNA-sequencing*. *J Inher Metab Dis*, 2018. **41**(3): p. 525-532.
12. Schiff, M., et al., *SLC25A32 Mutations and Riboflavin-Responsive Exercise Intolerance*. *New England Journal of Medicine*, 2016. **374**(8): p. 795-797.
13. Hellebrekers, D., et al., *Novel SLC25A32 mutation in a patient with a severe neuromuscular phenotype*. *Eur J Hum Genet*, 2017. **25**(7): p. 886-888.
14. Gerards, M., et al., *Riboflavin-responsive oxidative phosphorylation complex I deficiency caused by defective ACAD9: new function for an old gene*. *Brain*, 2011. **134**(Pt 1): p. 210-9.
15. Nouws, J., et al., *ACAD9, a complex I assembly factor with a moonlighting function in fatty acid oxidation deficiencies*. *Hum Mol Genet*, 2014. **23**(5): p. 1311-9.

16. Repp, B.M., et al., *Clinical, biochemical and genetic spectrum of 70 patients with ACAD9 deficiency: is riboflavin supplementation effective?* Orphanet J Rare Dis, 2018. **13**(1): p. 120.
17. Kadoya, T., et al., *Successful treatment of infantile-onset ACAD9-related cardiomyopathy with a combination of sodium pyruvate, beta-blocker, and coenzyme Q10.* J Pediatr Endocrinol Metab, 2019. **32**(10): p. 1181-1185.
18. Theunissen, T.E.J., et al., *Selection and Characterization of Palmitic Acid Responsive Patients with an OXPHOS Complex I Defect.* Frontiers in Molecular Neuroscience, 2017. **10**.