

# Unravelling atherosclerotic heterogeneity by single cell RNA sequencing

## Citation for published version (APA):

McCracken, I. R., Taylor, R. S., Henderson, N. C., Sluimer, J. C., & Baker, A. H. (2018). Unravelling atherosclerotic heterogeneity by single cell RNA sequencing. *Current Opinion in Lipidology*, 29(6), 488-489. <https://doi.org/10.1097/MOL.0000000000000559>

## Document status and date:

Published: 01/12/2018

## DOI:

[10.1097/MOL.0000000000000559](https://doi.org/10.1097/MOL.0000000000000559)

## Document Version:

Publisher's PDF, also known as Version of record

## Document license:

Taverne

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.



# Unravelling atherosclerotic heterogeneity by single cell RNA sequencing

Ian R. McCracken<sup>a</sup>, Richard S. Taylor<sup>b</sup>, Neil C. Henderson<sup>b</sup>  
Judith C. Sluimer<sup>a,c</sup>, and Andrew H. Baker<sup>a,c</sup>

The atherosclerotic plaque is a complex environment containing immune and stromal cells, many switching phenotype or even transitioning to alternative cell types [1–4]. Previous attempts to characterize heterogeneity in atherosclerosis have been limited by the requirement for predefined cell surface markers, thus negating new population discovery [5<sup>¶</sup>]. Consequently, cellular heterogeneity is poorly characterized and likely oversimplified [6–8].

This challenge in the atherosclerosis field has recently been addressed by the utility of single cell RNA sequencing (scRNA-seq), allowing transcriptome analysis at a single cell resolution [9<sup>¶¶</sup>, 10<sup>¶¶</sup>]. In contrast to previous priori-dependant approaches, scRNA-seq allows for the unbiased, comprehensive characterization of cellular identity within a heterogeneous population [5<sup>¶</sup>]. Importantly, droplet-based microfluidic platforms permit the sequencing of thousands of cells in parallel [11].

Two recent studies have utilized a droplet-based scRNA-seq platform to map leukocyte heterogeneity in control and fat-fed low-density lipoprotein receptor deficient (*Ldlr*<sup>-/-</sup>) [9<sup>¶¶</sup>] or apolipoprotein E (*Apoe*<sup>-/-</sup>) mice [10<sup>¶¶</sup>], respectively. Unsupervised clustering revealed more than 10 clusters in each model, with increased heterogeneity with advancing disease. Both publications show that the atherosclerotic plaques were predominantly composed of macrophages and T cells. Macrophages were found to account for a lower proportion of aortic leukocytes in *Apoe*<sup>-/-</sup> mice compared to that observed in *Ldlr*<sup>-/-</sup> mice, although the proportion of macrophages increased with disease progression in both models [9<sup>¶¶</sup>, 10<sup>¶¶</sup>].

Importantly, scRNA-seq identified a new macrophage cluster (identified by enriched expression of TREM2) and changed prevailing views on macrophage biology, as the identified inflammatory and resident macrophage populations demonstrated only a weak resemblance to polarized M1 and M2 macrophage phenotypes, again highlighting the limitations of the current classification system [10<sup>¶¶</sup>].

Understandably, the authors initially employed this new technology using homogenous mouse models, but it will be clearly important to utilize this technology in human vessels with underlying atherosclerosis where heterogeneity will be greater and more difficult to clearly quantify. Owing to changes in cell phenotype occurring asynchronously, the transcriptional signatures of transitioning cells can be used to reconstruct the associated trajectory [12]. This technology, termed as pseudotime analysis, has previously been utilized to identify potential regulators of myeloid and lymphocyte differentiation [13,14]. Consequently, application of pseudotime analysis may be crucial to identifying putative regulators of important dynamic processes such as the transdifferentiation of smooth muscle cells to macrophage-like cells [1,2]. Additionally, using an in-silico receptor-ligand screen such as that used by Camp et al. [15] could provide insight into the level of crosstalk between cell types, thus providing further insight into the pathogenesis of atherosclerosis. Together, single cell technology and advanced computational analyses will provide exciting opportunities to exploit cellular heterogeneity in the cardiovascular system, which will aid in understanding the healthy and disease vessel wall cellular and molecular dynamics, as well as refining future therapeutic and diagnostic strategies. However, complete characterization of atherosclerosis is likely to require the advent of new technologies capable of integrating the transcriptional characterization of single cells with genomic, epigenetic, and spatial proteomic data.

<sup>a</sup>Centre for Cardiovascular Science, <sup>b</sup>Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK and <sup>c</sup>Department of Pathology, CARIM School for Cardiovascular Diseases, Maastricht University Medical Center, Maastricht, The Netherlands

Correspondence to Andrew H. Baker, Centre for Cardiovascular Science, University of Edinburgh, 47 Little France Crescent, Edinburgh, EH16 4TJ, UK. Tel: +0131 24 26728; e-mail: Andy.Baker@ed.ac.uk.

**Curr Opin Lipidol** 2018, 29:488–489

DOI:10.1097/MOL.0000000000000559

## Acknowledgements

None.

## Financial support and sponsorship

I.R.M. is supported by the MRC Precision Medicine Doctoral Training Programme. J.C.S. received a VIDI fellowship of the Netherlands Organization of Scientific research (91718364). A.H.B. received funding from BHF CVR grant (RM/17/3/33381), ERC Advanced Grant VASMIR, and the BHF Chair of Translational Cardiovascular Sciences.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Rong JX, Shapiro M, Trogan E, Fisher EA. Transdifferentiation of mouse aortic smooth muscle cells to a macrophage-like state after cholesterol loading. *Proc Natl Acad Sci U S A* 2003; 100:13531–13536.
  2. Feil S, Fehrenbacher B, Lukowski R, *et al.* Transdifferentiation of vascular smooth muscle cells to macrophage-like cells during atherogenesis. *Circ Res* 2014; 115:662–667.
  3. Gistera A, Hansson GK. The immunology of atherosclerosis. *Nat Rev Nephrol* 2017; 13:368–380.
  4. Li Y, Lui KO, Zhou B. Reassessing endothelial-to-mesenchymal transition in cardiovascular diseases. *Nat Rev Cardiol* 2018; 15:445–456.
  5. Papalexli E, Satija R. Single-cell RNA sequencing to explore immune cell heterogeneity. *Nat Rev Immunol* 2018; 18:35–45.
- This article reviews the current and future applications of scRNA-seq to characterise immune cell heterogeneity.
6. Nahrendorf M, Swirski FK. Abandoning M1/M2 for a Network Model of Macrophage Function. *Circ Res* 2016; 119:414–417.
  7. Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Rep* 2014; 6:13.
  8. Tabas I, Bornfeldt KE. Macrophage phenotype and function in different stages of atherosclerosis. *Circ Res* 2016; 118:653–667.
  9. Cochain C, Vafadarnejad E, Arampatzis P, *et al.* Single-cell RNA-seq reveals ■ the transcriptional landscape and heterogeneity of aortic macrophages in murine atherosclerosis. *Circ Res* 2018; 122:1661–1674.
- This study uses scRNA-seq to determine CD45+ leukocyte heterogeneity in *Ldlr*<sup>-/-</sup> mice with a focus on characterizing macrophage populations.
10. Winkels H, Ehinger E, Vassallo M, *et al.* Atlas of the immune cell repertoire in ■ mouse atherosclerosis defined by single-cell RNA-sequencing and mass cytometry. *Circ Res* 2018; 122:1675–1688e.
- This study uses scRNA-seq alongside mass cytometry to characterise leukocyte heterogeneity in *Apoe*<sup>-/-</sup> and *Ldlr*<sup>-/-</sup> mice.
11. Haque A, Engel J, Teichmann SA, *et al.* A practical guide to single-cell RNA-sequencing for biomedical research and clinical applications. *Genome Med* 2017; 9:75.
  12. Trapnell C, Cacchiarelli D, Grimsby J, *et al.* The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol* 2014; 32:381–386.
  13. Drissen R, Buza-Vidas N, Woll P, *et al.* Distinct myeloid progenitor-differentiation pathways identified through single-cell RNA sequencing. *Nat Immunol* 2016; 17:666–676.
  14. Tsang JC, Yu Y, Burke S, *et al.* Single-cell transcriptomic reconstruction reveals cell cycle and multilineage differentiation defects in *Bcl11a*-deficient hematopoietic stem cells. *Genome Biol* 2015; 16:178.
  15. Camp JG, Sekine K, Gerber T, *et al.* Multilineage communication regulates human liver bud development from pluripotency. *Nature* 2017; 546:533–538.