

Cardiovascular magnetic resonance : a key to imaging cardiac function

Citation for published version (APA):

Schalla, S. M. (2015). *Cardiovascular magnetic resonance : a key to imaging cardiac function*. Datawyse / Universitaire Pers Maastricht. <https://doi.org/10.26481/dis.20150706ss>

Document status and date:

Published: 01/01/2015

DOI:

[10.26481/dis.20150706ss](https://doi.org/10.26481/dis.20150706ss)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Stellingen

behorende bij het proefschrift

Cardiovascular magnetic resonance

A key to imaging cardiac function

1. Magnetic resonance (MR) fluoroscopy allows cardiac catheterization and closure of atrial septal defects in an experimental model. Small intracardiac shunts are reliably detected with noninvasive MR imaging.
2. Real-time MR imaging saves time at the cost of reduced image quality. It enables the rapid assessment of hemodynamics and ischemia induced regional wall motion abnormalities. If image quality is sufficient we might replace standard scans with real-time scans. But who is accepting “sufficient” image quality? And if we saved time we would immediately add a new imaging sequence to the scan protocol anyway.
3. Infarct size is accurately measured with MR imaging. Hypertrophied hearts are more sensitive to ischemic injury than normal hearts.
4. MR imaging is the method of choice to detect and characterize myocardial infarcts. Using echocardiography, small infarcts will be missed.
5. Patients with cardiac syndrome X are suffering from microvascular dysfunction. In these patients, the subendocardial perfusion reserve assessed with MR imaging is significantly lower than the subepicardial perfusion reserve.
6. Focal myocardial fibrosis in patients with dilated cardiomyopathy is detected and quantified by MR imaging. Focal fibrosis is related to inflammation rather than increased interstitial fibrosis.
7. The systolic function of the right ventricle, the forgotten chamber, can accurately be assessed with MR imaging and is more impaired in dilated cardiomyopathy than in ischemic heart disease.
8. MR scanners are huge machines and cardiac MR imaging requires more time than other imaging modalities. Thus, we clearly have two reasons why the results of MR scans must be of superior quality.
9. Artsen willen steeds alles in beeld brengen om iets uit te sluiten (“uitsluitertis”). Vervolgens controleren ze regelmatig of ook alles goed blijft (“controlitis”).
10. Cardiologen en radiologen zijn in staat om samen te werken.
11. Auch für das Promovieren gilt: Doppelt hält besser!!