

Urocortin 3 Gene Transfer Increases Function of the Failing Murine Heart

Citation for published version (APA):

Giamouridis, D., Gao, M. H., Lai, N. C., Tan, Z., Kim, Y. C., Guo, T., Miyanohara, A., Blankesteyn, M. W., Biessen, E. A. L., & Hammond, H. K. (2019). Urocortin 3 Gene Transfer Increases Function of the Failing Murine Heart. *Human Gene Therapy*, 30(1), 10-20. <https://doi.org/10.1089/hum.2018.103>

Document status and date:

Published: 01/01/2019

DOI:

[10.1089/hum.2018.103](https://doi.org/10.1089/hum.2018.103)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Urocortin 3 Gene Transfer Increases Function of the Failing Murine Heart

Dimosthenis Giamouridis,¹⁻³ Mei Hua Gao,^{1,2} N. Chin Lai,^{1,2} Zhen Tan,^{1,2} Young Chul Kim,^{1,2} Tracy Guo,^{1,2} Atsushi Miyanojara,² Matthijs W. Blankestijn,³ Erik A. L. Biessen,^{3,4} and H. Kirk Hammond^{1,2,*}

¹Veterans Affairs San Diego Healthcare System, San Diego, California; ²Department of Medicine, University of California San Diego, San Diego, California; ³Department of Pharmacology and Toxicology, Cardiovascular Research Institute Maastricht University, Maastricht, The Netherlands; and ⁴Institute for Molecular Cardiovascular Research, RWTH Aachen, Aachen, Germany.

Peptide infusions of peptides the corticotropin releasing factor family, including urocortin 2, stresscopin, and urocortin 3 (UCn3), have favorable acute effects in clinical heart failure (HF), but their short half-lives make them unsuitable for chronic therapy. This study asked whether *UCn3* gene transfer, which provides sustained elevation of plasma UCn3 levels, increases the function of the failing heart. HF was induced by transmural left ventricular (LV) cryoinjury in mice. LV function was assessed 3 weeks later by echocardiography. Those with ejection fractions (EF) <40% received intravenous saline or intravenous adeno-associated virus type-8 encoding murine *UCn3* (AAV8.m*UCn3*; 1.9×10^{13} genome copies/kg). Five weeks after randomization, repeat echocardiography, assessment of LV function (+dP/dt, -dP/dt), and quantification of Ca²⁺ transients and sarcomere shortening in isolated cardiac myocytes were conducted, and assessment of LV Ca²⁺ handling and stress proteins was performed. Three weeks after myocardial infarction, prior to treatment, EFs were reduced (mean 31%, from 63% in sham-operated animals). Mice randomized to receive *UCn3* gene transfer showed increased plasma UCn3 (from 0.1 ± 0.01 ng/mL in the saline group to 5.6 ± 1.1 ng/mL; $n = 12$ each group; $p < 0.0001$). Compared to mice that received saline, *UCn3* gene transfer was associated with higher values for EF ($p = 0.0006$); LV +dP/dt ($p < 0.0001$), and LV -dP/dt ($p < 0.0001$). Cardiac myocytes from mice that received *UCn3* gene transfer showed higher peak Ca²⁺ transients ($p = 0.0005$), lower time constant of cytosolic Ca²⁺ decline (tau, $p < 0.0001$), and higher rates of sarcomere shortening (+dL/dt, $p = 0.03$) and lengthening (-dL/dt, $p = 0.04$). LV samples from mice that received *UCn3* gene transfer contained higher levels of SERCA2a ($p = 0.0004$ vs. HF) and increased amounts of phosphorylated troponin I ($p = 0.04$ vs. HF). *UCn3* gene transfer is associated with improved Ca²⁺ handling and LV function in mice with HF and reduced EF.

Keywords: gene therapy, heart failure with reduced EF, SERCA2a, AAV8

INTRODUCTION

CARDIOVASCULAR DISEASE is the most common cause of death in the United States. Heart failure (HF) affects 6 million people in the United States and has a dismal outlook, despite optimal therapy.¹ Because of such a poor prognosis, new approaches, including gene transfer, are warranted. Urocortin 3 (UCn3) is a 38-amino acid peptide in the corticotropin-releasing factor (CRF) family that binds with high affinity to corticotropin-releasing

hormone receptor-2 (CRHR2). Intravenous infusion of urocortin 2 (UCn2) peptide, which has 42% sequence homology with UCn3,² has beneficial cardiovascular effects,³ but UCn3 has not been studied as extensively for its cardioprotective activity. In preclinical studies, UCn3 peptide infusion provided cardiac protection against ischemia-reperfusion injury.⁴ Stresscopin, which has 96% sequence homology with UCn3, is efficacious when briefly infused in clinical HF.⁵ A recent study in

*Correspondence: Dr. H. Kirk Hammond, Department of Medicine, UCSD, VA San Diego Healthcare System, 3350 La Jolla Village Drive, San Diego, CA 92161. E-mail: khammond@ucsd.edu

healthy individuals and HF patients showed that UCn2 and UCn3 peptide infusions have beneficial effects, including reduced systemic vascular resistance, and increased cardiac output.⁶

The major impediment to the translation of these approaches to use in clinical HF is the short half-life (10 min) of these peptides.⁷ Sustained hemodynamic effect of these short-acting peptides would require continuous intravenous (i.v.) infusion, which hampers their clinical usefulness in chronic therapy for HF. A solution for such a problem would be using gene transfer to provide sustained plasma levels of the peptide. In animal studies, such an approach has been used to achieve sustained increases in plasma peptide concentration and increases in cardiac function.^{8,9} Pre-clinical studies have shown that such an approach, using i.v. delivery of an adeno-associated virus type 8 (AAV8) encoding UCn2, provided sustained increases in plasma UCn2 and is also beneficial for the normal and failing heart.^{8,9}

Data were recently published comparing the long-term cardiovascular effects of chronic exposure to sustained high plasma levels of UCn2 versus UCn3 using gene transfer in normal animals.¹⁰ Similar beneficial effects were found on cardiac function. However, UCn2 gene transfer increased glucose disposal and resulted in a significant decline in fasting glucose—an effect not shared by UCn3 gene transfer. These data indicate that UCn2 gene transfer may be an optimal selection for patients with diabetes and HF, while UCn3 may be better suited for the majority of patients with HF who are not diabetic. However, no previous studies have established that chronic elevation of plasma UCn3 is effective in the treatment of HF. The present study tested the hypothesis that UCn3 gene transfer would improve function of the failing heart.

MATERIALS AND METHODS

AAV8.UCn3 vector

HEK293T cells were transfected with the pRep2/Cap8 and pAd-Helper plasmid¹¹ to produce a helper-virus free AAV8 vector encoding murine urocortin-3 (UCn3) driven by a chicken β -actin (CBA) promoter (AAV8.CBA.mUCn3; Fig. 1). Plasmid pRep2/Cap8 was obtained from the University of Pennsylvania Vector Core. Virus vectors were then purified and concentrated, as previously described.^{9,12} Virus titers were determined by quantitative real-time polymerase chain reaction (RT-PCR), with virus genome DNA prepared from purified virus.

Animal use

The Animal Use and Care Committee of the VA San Diego Healthcare System approved the studies. Eighty-eight C57BL/6J mice (51 males) aged 8–12 weeks and weighing 24.5 ± 0.4 g were obtained (Jackson Laboratories, Bar Harbor, ME). Of these, 20 (9 males) underwent thoracotomy and heart manipulation but no cryoinjury; 68 underwent cryoinjury to induce HF.

HF model

Cryoinjury was used to induce anterior wall myocardial infarction (MI) and subsequent left ventricular (LV) chamber dilation and reduced function, as previously described.¹³ Animals were intubated and mechanically ventilated with oxygen and 1.5% isoflurane. To expose the heart, a thoracotomy was performed at the fourth left intercostal space, and the pericardium was opened. A cryoprobe 3.5 mm in diameter (Brymill, Ellington, CT) was applied to the LV anterior free wall for 10 s, followed by rinsing (25°C saline) to avoid traumatic detachment from the LV. This process was then repeated. Three weeks after cryoinjury, echocardiography was performed to assess heart function. Those with LV ejection fraction (EF) <40% were included in the study and received i.v. saline or i.v. AAV8.UCn3 (1.9×10^{13} genome copies [gc]/kg).

UCn3 gene transfer

Under anesthesia (1.5% isoflurane via nose-cone), the jugular vein was exposed, and a syringe with a 31-gauge needle was inserted to deliver AAV8.UCn3 (1.9×10^{13} gc/kg in 100 μ L) or a similar volume of saline.

Echocardiography

Echocardiography was performed, as previously described.¹⁴ Three weeks after MI, mice were anesthetized initially with 5% and then maintained at 1–1.5% isoflurane, and using a Vevo 3100 ultrasound system (FUJIFILM Visualsonics, Toronto, Canada), echocardiography was performed to document reduced LV function (EF <40%) and to record LV chamber dimensions. Echocardiographic assessment of heart function was then repeated 5 weeks after randomization of mice to receive i.v. delivery of AAV8.UCn3 or saline.

LV systolic and diastolic function

Sodium pentobarbital (80 mg/kg administered intraperitoneally [i.p.]) was used to anesthetize the animals, and a 1.4F micromanometer catheter (SPR 839; Millar Instruments, Houston, TX) was




Figure 1. *Urocortin 3 (UCn3)* transgene and study design. **(A)** AAV8.CBA.mUCn3 vector map. ITR, inverted terminal repeat; CMV.en, cytomegalovirus enhancer; CBA, chicken β -actin promoter; *UCn3*, murine *urocortin 3*; β GpA, β globin polyadenylation signal. **(B)** Study design. A representative cross-section of the left ventricle (LV) following cryoinjury, which resulted in transmural infarction of $27 \pm 2\%$ of the LV ($n=5$). **(C)** UCn3 plasma levels in mice with heart failure (HF) 5 weeks after intravenous (i.v.) saline (Sal) or i.v. AAV8.mUCn3 (*UCn3*). Mean \pm standard error (SE) are shown. Numbers in bars denote group size. p -Value is from a Student's t -test (unpaired, two-tailed). Color images available online at www.liebertpub.com/hum

advanced via the right carotid artery across the aortic valve and into the LV cavity. LV pressure was recorded, stored digitally, and then processed (IOX V2.9.5; Emka Technologies, Christchurch, VA), as previously reported.¹⁴ Subsequently, blood and tissue samples were obtained. The first derivatives of LV pressure development (LV +dP/dt) and decline (LV -dP/dt) were used to assess LV systolic and diastolic function, respectively, in a manner that was relatively load independent and a more accurate estimate of contractility than EF.

RT-PCR, immunoblotting

LV and liver samples were collected and stored at -80°C for quantitative RT-PCR and Western blotting. Total RNA was isolated and reverse transcribed into cDNA that was used in quantitative PCR, as previously described.⁸ The gene-specific primers used in PCR are listed in Table 1. Immunoblotting was performed, as described previously.⁸ The following antibodies were used: Phospho-PKA-RIIa, phospho-PKA-RIIb, CamKII, phospho-CamKII, and cardiac ankyrin repeat protein (CARP; Santa Cruz Biotechnology, Santa Cruz, CA); phospho-PKA catalytic subunit, total PKA, p-TnI, total TnI (Cell Signaling Technology, Danvers, MA); PLB (Thermo Fisher

Scientific, Waltham, MA); Ser 16 and Thr 17-phospho-PLB (Badrilla Ltd., Leeds, United Kingdom); SERCA2a (Enzo Life Sciences, Farmingdale, NY); and phospho-RYR2 and MYLK3 (Abcam, San Francisco, CA).

Table 1. Primers used in qRT-PCR

Gene	Forward	Reverse
α -MHC	5'-AAAGGCTGAGAGGAAGTACC	5'-ACCAGCCTTCTCCTCTGC
β -MHC	5'-GCTGAAAGCAGAAGAGATTATC	5'-TGGAGTTCTTCTCTTGGAG
α -SK-actin	5'-GTGTCACCCACAACGTGC	5'-AGGCCACATAGCACAGC
ANF	5'-CCTCGTCTTGGCCTTTTGG	5'-CATCTTCTACCGGCATCTTC
BNP	5'-GAAGTCCTAGCCAGTCTCC	5'-CAGCTTGAGATATGTGTACC
CARP	5'-CTGAACCTGTGGATGTGCC	5'-GGCTCCAGCCTCCATTAAT
FHL-1	5'-TGCAACAAGTGCCTACTCG	5'-CAATGTTTGGCGAAGCTGGTC
Coll1 α 1	5'-GCCAAGAAGACATCCCTGAAG	5'-GGGTCCCTCGACTCTAC
Coll3 α 1	5'-GCACAGCAGTCCAACGTAGA	5'-TCTCCAAATGGGATCTCTGG
MMP8	5'-GACTCTGGTGATTTCTTGCTAAG	5'-CACCATGGTCTCTTGAGACG
MMP9	5'-CGTCGTGATCCCCACTTACT	5'-GAACACACAGGGTTTGCCTTC
TIMP1	5'-GACAGCTTTCTGCAACTCGG	5'-CTTGTGGACATATCCACAGAGG
IL-1b	5'-CCCATCCTCTGTGACTCATG	5'-AAGGCCACAGGATTTTGTGC
IL-6	5'-CGGAGAGGAGACTTCCAG	5'-TTCTGCAAGTGCATCATCGTC
IL-10	5'-GCTCTTACTGACTGGCATGAG	5'-CGCAGCTCTAGGAGCATGTG
UCn3	5'-ACGCACCTCCAGATCAAAG	5'-AATTCTTGGCCTTGTGATG

qRT-PCR, quantitative real-time polymerase chain reaction; α -MHC, alpha-myosin heavy chain; β -MHC, beta-myosin heavy chain; α -SK-actin, alpha-skeletal actin; ANF, atrial natriuretic peptide; BNP, brain natriuretic peptide; CARP, cardiac ankyrin repeat protein; FHL-1, four and a half LIM domains protein 1; Coll, collagen; MMP, matrix metalloproteinase; TIMP1, tissue inhibitor of metalloproteinase 1; IL, interleukin; UCn3, urocortin 3.

Cardiac myocyte isolation and Ca²⁺ transients

Cardiac myocytes were isolated, as previously described.¹⁴ Mice were heparinized (5,000 IU/kg i.p.), and then anesthetized with sodium pentobarbital (150 mg/kg i.p.). Hearts were extracted and quickly cannulated via the aorta for perfusion. Using a peristaltic pump, a calcium-free medium containing Joklik-modified minimal essential medium (Gibco-BRL; Thermo Fisher Scientific) was perfused for 3 min (3 mL/min), followed by perfusion of the same buffer with the addition of 20 μ M of CaCl₂ and collagenase type II (Worthington Biochemical Corp., Lakewood, NJ) for another 12–13 min. Hearts were then placed into a beaker with digestion solution, atria were removed, and ventricles were sliced into pieces and mixed gently with a wide-nozzle pipette (3–4 min) to dissociate the myocytes. Isolated cardiac myocytes were plated onto laminin pre-coated glass coverslips and loaded with Indo-1/AM (with 0.02% pluronic F-127) for 20 min. Coverslips then were placed in a field stimulation perfusion chamber, rinsed to remove excess stain, and mounted on a Nikon TMD inverted microscope (Nikon, Melville, NY) equipped with a 40 \times objective interfaced to a photomultiplier-based fluorescence measuring system (PTI, Edison, NJ). During the measurements, cardiac myocytes were continuously perfused with KRH solution (125 mM of NaCl, 5 mM of KCl, 1.2 mM of NaH₂PO₄, 6 mM of glucose, 1.2 mM of MgCl₂, and 25 mM of HEPES, pH 7.37–7.38) containing 2 mM of CaCl₂, and they were field stimulated at 0.3 Hz. Using an excitation wavelength of 365 nm via a monochromator, fluorescent emission was directed to two photomultiplier tubes through 20 nm band-pass filters centered at 405 and 485 nm, respectively. Indicator of the [Ca²⁺]_i is the ratio F₄₀₅/F₄₈₅. Multiple cardiac myocytes from each heart of three to five mice per group were used to obtain the Ca²⁺ transient data.

Quantification of fibrosis, LV apoptosis, and infarct size

Fibrosis. Samples of liver and viable (uninfarcted) transmural LV samples were formalin fixed and paraffin embedded. Five-micron sections were mounted and counterstained with hematoxylin and eosin and Masson's trichrome.

Infarct size. A separate group of animals ($n=5$) underwent cryoinjury. Eight days later, hearts were arrested in diastole (0.6% KCl) and the right ventricle removed, and LV (including septum) was pressed between two microscopic slides (1.2 mm apart) and scanned using an EPSON V850 pro scanner (Epson, Nagano, Japan). Using ImageJ

v1.49 software, the area of transmural infarct was measured and recorded as percent of total LV.

Apoptosis. Paraffin-embedded tissue sections were rehydrated and then microwaved for 5 min in 10 mM of sodium citrate buffer (pH 6.0) for epitope retrieval. After quenching of endogenous peroxidase activity and blocking with normal goat serum, the tissue sections were incubated (18 h at 4°C) with anti-cleaved caspase-3 antibody (1:200; Cell Signaling Technology) to assess apoptosis. After washing, immunostaining was performed using an ImmPRESS HRP kit (Vector Laboratories, Burlingame, CA) according to the manufacturer's instructions.

Statistical analysis

Data acquisition and analysis were performed without knowledge of group identity. Group sizes were determined by power calculations, and male and female mice were used. Since the hypothesis was that *UCn3* gene transfer would improve function of the failing heart, the key statistical comparison was between mice with HF that received i.v. saline and those that received *UCn3* gene transfer, and Student's *t*-test was used (unpaired, two-tailed). In order to confirm the presence of LV dysfunction, a sham-operated age- and sex-matched control group was included and, when appropriate, additional between-group *t*-tests (control vs. HF) were performed, with Bonferroni correction. Data represent mean \pm standard error (SE). Analyses were performed using GraphPad Prism (GraphPad Software, Inc., San Diego, CA). The null hypothesis was rejected when $p < 0.05$.

RESULTS

Plasma UCn3 levels

Plasma levels of UCn3 were measured 5 weeks after gene transfer. Mice that received *UCn3* gene transfer showed increased plasma UCn3 concentrations (HF: 0.1 ± 0.01 ng/mL, $n=12$; HF + *UCn3*: 5.6 ± 1.1 ng/mL, $n=12$; $p < 0.0001$; Fig. 1). There was no sex difference in plasma UCn3 concentrations in HF saline-injected mice ($p=0.5$), but in mice that received *UCn3* gene transfer, male mice had higher plasma UCn3 levels than female mice (male: 8.5 ± 1 ng/mL, $n=6$; female: 2.7 ± 0.8 ng/mL; $n=6$; $p=0.001$).

MI-induced HF

Sixteen mice died in the first 3 weeks after MI, and 12 did not show EF <40%. Thus, of 68 mice that underwent infarction by cryoinjury, 40 met the

enrollment criteria 3 weeks later, giving a yield of 59%. This method is superior to coronary occlusion, where the yield can be <10%.¹⁵ No mice died in the 5-week interval after treatment.

Infarct size. Cryoinjury resulted in transmural infarction of $27 \pm 2\%$ of the LV ($n=5$). Figure 1B shows a representative cross-section of the LV following cryoinjury infarction.

Echocardiography

Three weeks after MI (before randomization), reductions in EF ($p < 0.0001$) and velocity of circumferential fiber shortening were seen (corrected for heart rate, VCFc; $p < 0.0001$; Table 2). In addition, end-systolic dimensions (ESD) and end-diastolic dimensions (EDD) were increased among HF mice compared to controls (Table 2), confirming that cryoinjury-induced MI resulted in LV chamber dilation and dysfunction. There were no pre-treatment group differences in HF mice in EF, VCFc, EDD, or ESD. However, 8 weeks after MI (5 weeks after saline or *UCn3* gene transfer,) beneficial between-group differences conferred by *UCn3* gene transfer on EF were seen. Indeed, the final EF was nine percentage units higher (a 32% relative increase) in mice that received *UCn3* gene transfer (HF: $28 \pm 1\%$, $n=13$; HF + *UCn3*: $37 \pm 1\%$, $n=13$; $p=0.0006$; Table 3). In addition, *UCn3* gene transfer was associated with higher VCFc ($p=0.0001$; Table 3).

LV function

Eight weeks after MI, mice with HF that received *UCn3* gene transfer showed an elevation in LV peak +dP/dt (HF: $3,941 \pm 159$ mmHg/s, $n=11$; HF + *UCn3*: $7,353 \pm 582$ mmHg/s, $n=12$; $p < 0.0001$; Fig. 2A), LV peak -dP/dt (HF: $-3,713 \pm 211$ mmHg/s, $n=11$; HF + *UCn3*: $-6,302 \pm 270$ mmHg/s, $n=12$; $p < 0.0001$; Fig. 2B) and developed LV pressure (HF: 76 ± 2 mmHg/s, $n=11$; HF + *UCn3*: 95 ± 4 mmHg/s, $n=12$; $p=0.0006$; Fig. 2C). Heart rates of the HF

saline-treated group were lower than the HF *UCn3*-treated group (HF: 350 ± 13 mmHg/s, $n=11$; HF + *UCn3*: 440 ± 20 mmHg/s, $n=12$; $p=0.001$; Fig. 2D). After *UCn3* gene transfer, there were no sex differences in LV peak +dP/dt (male: $7,288 \pm 586$ mmHg/s, $n=7$; female: $7,444 \pm 1,228$ mmHg/s, $n=5$; $p=0.9$) or in LV peak -dP/dt (male: $-6,457 \pm 303$ mmHg/s, $n=7$; female: $-6,086 \pm 519$ mmHg/s, $n=5$; $p=0.5$).

Cytosolic Ca²⁺ transients and sarcomere length

Cardiac myocytes were isolated 8 weeks after MI (5 weeks after randomization to saline or *UCn3* gene transfer) to assess Ca²⁺ transients. Cardiac myocytes from mice that received *UCn3* gene transfer showed higher peak cytosolic Ca²⁺ concentration (Fig. 3A and B; $p=0.0005$) and lower Tau (Fig. 3D; $p < 0.0001$). Time to peak Ca²⁺ concentration showed no group difference. Sarcomere shortening was also measured in these cardiac myocytes, and it was found that *UCn3* gene transfer was associated with a higher shortening rate (+dL/dt; HF: 7.8 ± 0.5 $\mu\text{m/s}$, $n=39$; HF + *UCn3*: 9.8 ± 0.7 $\mu\text{m/s}$, $n=20$; $p=0.03$; Fig. 3E) and a higher lengthening rate (-dL/dt; HF: 6.7 ± 0.4 $\mu\text{m/s}$, $n=39$; HF + *UCn3*: 8.3 ± 0.6 $\mu\text{m/s}$, $n=20$; $p=0.04$; Fig. 3F). These data are consistent with group differences in Ca²⁺ transients and pressure development and decline *in vivo* (Fig. 2A-C).

Necropsy

Body and liver weights showed no group differences (Table 4). HF mice compared to control mice showed higher weights of LV ($p=0.002$) and LV/body weight ratio ($p < 0.0002$), which were increased similarly after MI in HF and HF + *UCn3* groups (Table 4).

LV Ca²⁺ handling protein expression

After MI, LV content of sarco/endoplasmic reticulum Ca²⁺ ATPase (SERCA2a) protein was higher after *UCn3* gene transfer (HF: 0.41 ± 0.02 ,

Table 2. Echocardiography 3 weeks after MI (no treatment)

	Control (16)	HF (13)	HF + <i>UCn3</i> (13)	<i>p</i> , control vs. HF
HR (bpm)	541 \pm 6	535 \pm 5	544 \pm 6	0.5
EDD (mm)	3.7 \pm 0.1	4.5 \pm 0.1	4.6 \pm 0.1	<0.0001
ESD (mm)	2.4 \pm 0.1	3.8 \pm 0.1	4.0 \pm 0.1	<0.0001
IVSd (mm)	0.7 \pm 0.03	0.8 \pm 0.02	0.8 \pm 0.03	<0.0001
LVEF (%)	63 \pm 2	31 \pm 2	30 \pm 2	<0.0001
VCFc (circ/s)	22.2 \pm 0.9	9.6 \pm 0.6	9.8 \pm 0.6	<0.0001

p-Values from student's *t*-tests (unpaired, two-tailed) with Bonferroni correction for multiple comparisons; numbers in parentheses indicate group size. MI, myocardial infarction; HF, heart failure; HR, heart rate; bpm, beats per minute; EDD, LV end - diastolic diameter; ESD, LV end - systolic diameter; IVSd, interventricular septum thickness in diastole; LVEF, left ventricular ejection; VCFc, velocity of circumferential fiber shortening (corrected for heart rate); control, sham-operated animals.

Table 3. Echocardiography 5 weeks after UCn3 gene transfer

	Control (16)	HF (13)	HF + UCn3 (13)	p	
				Control vs. HF	HF vs. HF + UCn3
HR (bpm)	544±6	535±7	539±6	0.6	1
EDD (mm)	3.6±0.1	4.6±0.1	4.6±0.2	<0.0002	1
ESD (mm)	2.4±0.1	4.0±0.1	3.8±0.2	<0.0002	0.4
IVSd (mm)	0.7±0.02	0.9±0.05	0.8±0.06	<0.0002	0.8
LVEF (%)	62±3	28±1	37±1	<0.0002	0.0006
VCFc (circ/s)	21.8±1	9.3±0.5	12.3±0.5	<0.0002	0.001

p-Values from a Student's *t*-test (unpaired, two-tailed) with Bonferroni correction for multiple comparisons; numbers in parentheses indicate group size.

$n=4$; HF + UCn3: 0.52 ± 0.01 , $n=4$; $p=0.0004$; Fig. 4A). LV content of phosphorylated troponin I (p-TnI) was lower after MI ($p=0.01$) but was normalized after UCn3 gene transfer. It was found that p-TnI was 50% higher in animals with HF that received UCn3 (control: 0.31 ± 0.01 , $n=4$; HF: 0.20 ± 0.03 , $n=4$; HF + UCn3: 0.30 ± 0.02 , $n=4$; $p=0.04$; Fig. 4B). Selected additional proteins relevant to Ca^{2+} handling showed reduced phosphorylation (p-RYR2, p-16-PLB, and p17-PLB) after MI, but no HF versus HF + UCn3 group differences were seen (Table 5).

LV mRNA expression of markers of hypertrophy and fibrosis

As expected, there were group differences in LV mRNA expression of several markers of LV stress 8 weeks after MI. However, *brain natriuretic peptide* (BNP; $p=0.006$), *atrial natriuretic peptide* (ANF;

$p=0.05$), *α -skeletal actin* ($p=0.02$), and *CARP* ($p=0.01$) were lower in HF + UCn3 than in HF mice. The levels of LV *beta-myosin heavy chain* (β -MHC) mRNA were lower in HF + UCn3 versus HF mice ($p<0.01$; Table 6). Although several genes related to fibrosis showed increased LV mRNA expression in HF versus control mice, their expression was not altered by UCn3 gene transfer.

Fibrosis and LV apoptosis

Fibrosis. There were no group differences in fibrosis in the liver or in the uninfarcted region of LV, and there were no differences in inflammatory infiltrates (Fig. 5A).

LV apoptosis. Quantification of active caspase-3 staining showed more apoptosis after MI ($p<0.0001$), with similar amounts of apoptosis in HF and HF + UCn3 mice (control: $0.04\pm 0.01\%$,




Figure 2. *In vivo* assessment of LV function. Eight weeks after myocardial infarction (MI) 5 weeks after i.v. delivery of AAV8.UCn3 (1.9×10^{13} gc/kg; HF + UCn3) or saline (HF), mice underwent physiological studies to assess LV function. (A and B) Peak rate of LV pressure development (A, +dP/dt) and peak rate of LV pressure decline (B, -dP/dt). These data indicate UCn3 gene transfer increased both peak +dP/dt and peak -dP/dt in mice with HF, indicating increased systolic and diastolic LV function. (C) LV developed pressure (LVP) was increased by UCn3 gene transfer. (D) Heart rate (HR) in anesthetized animals was lower in the saline group. Individual data are shown (mean±SE are indicated); p-values are from a Student's *t*-test (unpaired, two-tailed). Color images available online at www.liebertpub.com/hum




Figure 3. Cytosolic Ca^{2+} transients. **(A)** Representative Indo-1 Ca^{2+} transient recordings from cardiac myocytes from one heart in each group. **(B)** Mean peak Ca^{2+} transients from multiple cardiac myocytes from each group, showing increased peak Ca^{2+} transients in cardiac myocytes isolated from mice with HF versus HF mice treated with *UCn3* gene transfer (HF + *UCn3*). **(C)** Time-to-peak cytosolic Ca^{2+} concentration showed no group difference between mice with HF that received saline versus *UCn3* gene transfer. **(D)** Time constant of cytosolic Ca^{2+} decline (τ) showed shorter τ (more rapid decline) in cardiac myocytes isolated from mice following *UCn3* gene transfer. **(E)** Rate of sarcomere shortening in systole (+dL/dt) was higher in HF animals that received *UCn3* gene transfer. **(F)** Rate of sarcomere lengthening in diastole (-dL/dt) was higher in HF animals that received *UCn3* gene transfer. In **(B-F)**, summary data are from cardiac myocytes isolated from three to five mice per group. Mean \pm SE are shown; p -values are from a Student's t -test (unpaired, two-tailed). Color images available online at www.liebertpub.com/hum

$n = 14$; HF: $0.26 \pm 0.05\%$, $n = 8$; HF + *UCn3*: $0.31 \pm 0.06\%$, $n = 7$; Fig. 5B).

DISCUSSION

The major finding of this study is that *UCn3* gene transfer increased the function of the failing heart. Three weeks after MI, LV EF was reduced from a mean value of 63% (control) to 30–31%. Five weeks later, mice that received *UCn3* gene transfer showed increased EF, while those that received saline showed a further decline. The final EF was nine percentage units higher (32% relative increase) in mice that received *UCn3* gene transfer

($p = 0.0006$; Table 3). Furthermore, LV peak +dP/dt showed a 1.9-fold increase in HF + *UCn3* versus HF ($p < 0.0001$; Fig. 2A). Finally, LV diastolic function after MI was also enhanced by *UCn3* gene transfer. LV peak -dP/dt, a measure of diastolic function, showed a 1.7-fold increase in HF + *UCn3* versus HF ($p < 0.0001$; Fig. 2B).

What explains these improvements in function of the failing heart associated with *UCn3* gene transfer? If the evidence for improved function resided in EF alone, one could propose that the benefit might reflect a vasodilating effect of *UCn3*^{5,6} because a reduction in systemic pressure would be expected to enhance EF. However, the favorable

Table 4. Necropsy data 5 weeks post saline or AAV8.*UCn3* administration in mice with HF

	Con (15)	HF (16)	HF + <i>UCn3</i> (13)	p	
				Control vs. HF	HF vs. HF + <i>UCn3</i>
BW (g)	26.9 \pm 1.1	26.5 \pm 0.7	26.7 \pm 0.9	1	1
LV (mg)	84 \pm 4	110 \pm 6	102 \pm 5	0.002	0.8
LV/BW (mg/g)	3.1 \pm 0.1	5.0 \pm 0.2	4.8 \pm 0.3	<0.0002	1
Liver/BW (mg/g)	45 \pm 1	45 \pm 1	43 \pm 1	1	0.8

p -Values from a Student's t -test (unpaired, two-tailed) with Bonferroni correction for multiple comparisons; numbers in parentheses indicate group size. BW, body weight.




Figure 4. LV SERCA2a and phosphorylation of troponin I (TnI) in mice with HF 5 weeks after *UCn3* gene transfer. **(A)** LV SERCA2a content was higher in mice with HF that received *UCn3* gene transfer ($p=0.0004$ vs. HF). **(B)** Troponin I (TnI) phosphorylation at serine 23 and 24 was higher in HF mice that received *UCn3* gene transfer ($p=0.04$ vs. HF). Individual mouse data are shown (mean \pm SE are indicated); Con, sham-operated animals. p -Values are from a Student's t -test (unpaired, two-tailed) with Bonferroni correction for multiple comparisons. Color images available online at www.liebertpub.com/hum

effects of *UCn3* gene transfer also were seen in LV peak $+dP/dt$, an event that occurs prior to aortic valve opening, and therefore is not as affected by systemic pressure. Likewise, isolated cardiac myocytes from HF mice that received *UCn3* gene transfer showed a higher sarcomere shortening rate ($+dL/dt$) versus HF alone, confirming that the benefits reside within the cardiac myocyte *per se* and do not reflect vasodilatory enhancement of LV function.

The group difference in HR during measures of LV peak $+dP/dt$ (Fig. 2) may have contributed to the increase in LV peak $+dP/dt$ via a positive force–frequency relationship. However, LV peak $+dP/dt$ was increased 90% in mice with HF treated with *UCn3* gene transfer, and the force–frequency effect would account for a much smaller change.¹⁶ Finally, the increase in VCFc (at matched HR) and the increase in isolated cardiac myocyte shortening rates indicate changes in function not related to HR.

Table 5. LV protein levels (immunoblotting)

Protein	Con (4)	HF (4)	HF + UCn3 (4)	p	
				Control vs. HF	HF vs. HF + UCn3
p-PKA-RIIa	0.6 \pm 0.1	0.7 \pm 0.1	0.6 \pm 0.1	1	0.6
p-PKA-RIIb	0.3 \pm 0.15	0.3 \pm 0.03	0.2 \pm 0.03	1	0.8
Total-PKA	0.4 \pm 0.01	0.3 \pm 0.02	0.3 \pm 0.02	1	0.8
p-PKA/Total	1.8 \pm 0.3	2.0 \pm 0.10	1.9 \pm 0.1	1	0.8
CamKII	0.6 \pm 0.1	0.5 \pm 0.1	0.6 \pm 0.1	1	0.8
p-CamKII	0.5 \pm 0.1	0.4 \pm 0.02	0.6 \pm 0.1	0.8	0.2
p-CamKII/Total	0.9 \pm 0.15	0.9 \pm 0.1	1.1 \pm 0.2	1	1
p-RYR2/Total	2.2 \pm 0.4	0.7 \pm 0.1	1.2 \pm 0.1	0.01	0.06
p16-PLB	0.1 \pm 0.1	0.7 \pm 0.1	0.9 \pm 0.11	1	0.8
p16-PLB/Total	3.7 \pm 0.6	1.1 \pm 0.2	2.1 \pm 0.5	0.01	0.2
p17-PLB	6.3 \pm 0.7	2.4 \pm 0.8	2.2 \pm 0.4	0.02	1
p17-PLB/Total	28.6 \pm 3.5	3.5 \pm 1.1	5.4 \pm 1.5	0.001	0.6
CARP	10.3 \pm 1	14 \pm 1.8	14 \pm 1.1	0.2	1
MYLK3	0.6 \pm 0.1	1.0 \pm 0.3	0.7 \pm 0.1	0.4	0.8

Immunoblotting data presenting protein levels normalized for GAPDH or vinculin. Data are mean \pm SE. p -Values from a Student's t -test (unpaired, two-tailed) with Bonferroni correction for multiple comparisons; numbers in parentheses indicate group size.

p-TnI, S23/24 phosphorylated troponin I; PKA-RIIa, protein kinase A IIa; CamKII, Ca²⁺/calmodulin-dependent protein kinase II; RYR, ryanodine receptor; PLB, phospholamban; MYLK3, myosin light chain kinase 3.

Table 6. LV mRNA expression of markers of hypertrophy and fibrosis

Gene	HF (4), fold control	HF + UCn3 (4), fold control	p
α -MHC	0.5 \pm 0.1	0.6 \pm 0.1	0.40
β -MHC	25 \pm 5	6.6 \pm 0.5	0.01
α -skeletal actin	5.3 \pm 1	2.3 \pm 0.3	0.02
ANF	15 \pm 3.8	5.6 \pm 0.6	0.05
BNP	3.4 \pm 0.4	1.6 \pm 0.2	0.006
CARP	3.5 \pm 0.3	2.3 \pm 0.04	0.01
FHL-1	1.3 \pm 0.2	1.6 \pm 0.1	0.16
Col1	3.8 \pm 0.7	2.7 \pm 0.2	0.21
Col3	3.8 \pm 0.6	3.0 \pm 0.3	0.30
MMP8	0.4 \pm 0.1	0.5 \pm 0.2	0.89
MMP9	0.4 \pm 0.1	0.4 \pm 0.1	0.74
TIMP1	0.7 \pm 0.01	0.7 \pm 0.1	0.46
IL-1b	0.4 \pm 0.02	0.4 \pm 0.1	0.89
IL-6	3.9 \pm 0.4	4 \pm 1.3	0.93
IL-10	0.8 \pm 0.1	1.2 \pm 0.3	0.17

Data are mean \pm SE. *p*-Values are from a Student's *t*-test (unpaired, two-tailed); numbers in parentheses indicate group size.

Col1, collagen 1; col3, Collagen 3; MMP8, matrix metalloproteinase 8; MMP9, matrix metalloproteinase 9; IL-1b, interleukin 1b; IL-6, interleukin 6; IL-10, interleukin 10.

What then might explain intrinsic improvement in function of the LV and cardiac myocytes isolated from failing hearts treated with *UCn3* gene transfer? The rate of cardiac myocyte shortening

during systole (+dL/dt, *p*=0.03; Fig. 3E) and lengthening during diastole (−dL/dt, *p*=0.04; Fig. 3F) were increased in the HF + *UCn3* group versus the HF group. These findings show tight correlations with LV peak +dP/dt and peak −dP/dt, physiological measures of the intact heart. *UCn3* gene transfer increases Ca²⁺ handling, and subsequently increases LV systolic and diastolic function. It was found that cardiac myocytes from HF + *UCn3* mice showed increased peak Ca²⁺ transients (*p*=0.0005; Fig. 3B) and reduced time constant of Ca²⁺ decline (tau, *p*<0.0001; Fig. 3D) compared to those from HF mice. This may explain the increases in LV systolic and diastolic function.

To determine molecular underpinnings for increased LV Ca²⁺ handling, LV protein expression of key Ca²⁺ related proteins was assessed. LV SERCA2a returns cytosolic Ca²⁺ to the sarcoplasmic reticulum, and consequently, increased expression of this protein would increase cytosolic Ca²⁺ decline, as was observed (Fig. 3D). An increase in cytosolic Ca²⁺ decline, by increasing Ca²⁺ availability to myofilaments, could also increase systolic function. It was found that LV SERCA2a protein levels were similar in control




Figure 5. Histological analysis of LV and liver and LV apoptosis 5 weeks after *UCn3* gene transfer. **(A)** Hematoxylin and eosin (H&E) and Masson's trichrome staining were performed on liver and transmural sections of LV (original magnification 20 \times ; presented image magnification H&E and Masson's trichrome: 7 \times), no evidence of inflammatory infiltrates was seen. LV fibrosis of the viable (non-infarcted LV) was similar in all three groups 8 weeks after MI. There were no group differences in liver fibrosis. **(B)** LV apoptosis was assessed using active caspase-3 staining on transmural LV sections, which showed a group difference between control and HF groups. Standard, control for Masson's trichrome. Individual data are shown (mean \pm SE); *p*-values are from a Student's *t*-test (unpaired, two-tailed). Color images available online at www.liebertpub.com/hum

and HF mice but were higher after *UCn3* gene transfer ($p=0.0004$ vs. HF; Fig. 4A). Upregulation of SERCA2a would be anticipated to increase peak systolic Ca^{2+} transient amplitude and rate of Ca^{2+} decline (Fig 3A, B, and D), and subsequently increase systolic and diastolic function of isolated cardiac myocytes (Fig. 3E and F) and the intact heart (Fig. 2A and B). Cardiac myocytes would no longer be exposed to sustained high plasma levels of UCn3 once isolated. However, one would anticipate that increases in SERCA2a protein, which has a half-life of 2–3 days,¹⁷ would endure for the 2 h interval required for cell isolation and study.

Improved Ca^{2+} handling continues to be seen in the cardiac myocytes that were isolated from animals that received *UCn3* gene transfer. Although increased LV SERCA2a provides a potential mechanism for UCn3's effects on LV Ca^{2+} handling, it does not elucidate the molecular mechanism by which this occurs, which will require further studies.

Troponin I phosphorylation plays a role in cardiac contractility and relaxation and is impaired under ischemic conditions. In the current study, phosphorylation of TnI at Serine residues 23 and 24 was reduced in the HF group but restored to normal by *UCn3* gene transfer (HF vs. HF+*UCn3*, $p=0.04$; Fig. 4B). The implications of the restoration of LV p-TnI to normal levels vis-à-vis cardiac function and the mechanism by which UCn3 affected this change were not established in the present study. However, reduction in TnI phosphorylation at Ser 23/24 is seen in explanted hearts from patients with end-stage HF,¹⁸ and Ser 23/24 phosphorylation of TnI is associated with more rapid relaxation,¹⁹ and therefore may be of mechanistic importance for the enhanced diastolic function conferred by *UCn3* gene transfer in the present study.

There were increases in the HF group (vs control) in *ANF*, *BNP*, α -skeletal actin, and β -MHC, which are molecular markers of hypertrophy and stress.^{20–22} These markers were significantly less increased after *UCn3* gene transfer (Table 6). These group difference likely are a consequence of improved LV function in HF after *UCn3* gene transfer. LV *CARP* expression is increased in clinical HF and in many animal models of HF, and may exacerbate adverse cardiac remodeling and apoptosis.^{23,24} Increased LV *CARP* mRNA was seen in HF mice, as was somewhat less of an increase in the HF + *UCn3* group ($p=0.01$; Table 6). LV mRNA expression of *Col 1*, *Col3*, and *IL-6* were increased similarly in HF and HF + *UCn3* mice (Table 6).

HF is associated with protean abnormalities, and it is uncertain whether the benefits seen in

mice after UCn3 gene transfer would also be seen in clinical HF. The present study tested *UCn3* gene transfer in HF with reduced EF on endpoints obtained 5 weeks after initiation of therapy. Will *UCn3* gene transfer have enduring effects, and will such effects be seen in models of HF with preserved EF? Despite these limitations, it is promising that a simple i.v. injection of AAV8.*UCn3* has such benefits on function of the failing heart.

AAV8 was selected because the target was the liver, and the beneficial effects reside in increased plasma concentrations and do not depend on increasing LV expression *per se*. In head-to-head experiments of various vectors and promoters, AAV8.CBA was superior to others.⁸ After i.v. AAV8 vector delivery, the primary organ in which one detects transgene expression is the liver,^{8,10} which is the target of this paracrine-based gene transfer. Another potential advantage is less pre-existing anti-AAV8 antibodies in humans versus AAV9.²⁵

The National Institutes of Health has requested that preclinical studies be conducted in similar proportions of male and female animals when possible,²⁶ a requirement embraced by the Food and Drug Administration for Investigational New Drug applications. In the current study, the same amount of AAV8.*UCn3* (1.9×10^{13} gc/kg i.v.) was associated with threefold higher plasma UCn3 concentrations in male versus female mice. Even so, the mean plasma UCn3 concentration in females after gene transfer was >20-fold higher than that of mice that did not receive vector and was a sufficient amount to evoke equivalent LV +dP/dt and LV -dP/dt as was found in male mice. These data indicate that lower doses of AAV8.*UCn3* than used in the current study will suffice to obtain a maximal effect. The dose–response effect of AAV8.*UCn3* in each sex will be addressed in separate studies.

In conclusion, *UCn3* gene transfer increases systolic and diastolic function of the failing heart. These beneficial effects of *UCn3* gene transfer reverberate from Ca^{2+} transients and sarcomere shortening in isolated cardiac myocytes to measures of LV function *in vivo*. The mechanism for these benefits can be attributed to improved LV Ca^{2+} handling, which directly affects sarcomere shortening, and is possibly a consequence of increased expression of LV SERCA2a and perhaps restoration of TnI phosphorylation. These studies indicate that chronically increasing plasma levels of UCn3 may be beneficial in HF with reduced EF. Translating this to possible clinical application will require proof of safety and efficacy and in other models of HF, and studies in larger mammals.

ACKNOWLEDGMENTS

This work was supported by National Institutes of Health grant (P01 HL66941) and Merit grants from the Department of Veteran's Affairs (1101 BX001515 and 1101 BX003774).

AUTHOR DISCLOSURE

H.K.H. is a founder, board member, and unpaid consultant of Renova Therapeutics. Renova redundant as stated played no role in the studies. None of the other authors have any disclosures.

REFERENCES

- Lloyd-Jones D, Adams RJ, Brown TM, et al. Executive summary: heart disease and stroke statistics—2010 update: a report from the American Heart Association. *Circulation* 2010;121:948–954.
- Lewis K, Li C, Perrin MH, et al. Identification of urocortin III, an additional member of the corticotropin-releasing factor (CRF) family with high affinity for the CRF2 receptor. *Proc Natl Acad Sci U S A* 2001;98:7570–7575.
- Chan WY, Frampton CM, Crozier IG, et al. Urocortin-2 infusion in acute decompensated heart failure: findings from the UNICORN study. *JACC Heart Fail* 2013;1:433–441.
- Brar BK, Jonassen AK, Egorina EM, et al. Urocortin-II and urocortin-III are cardioprotective against ischemia reperfusion injury: an essential endogenous cardioprotective role for corticotropin releasing factor receptor type 2 in the murine heart. *Endocrinology* 2004;145:24–35.
- Gheorghiadu M, Greene SJ, Ponikowski P, et al. Haemodynamic effects, safety, and pharmacokinetics of human stresscopin in heart failure with reduced ejection fraction. *Eur J Heart Fail* 2013;15:679–689.
- Stirrat CG, Venkatasubramanian S, Pawade T, et al. Cardiovascular effects of urocortin 2 and urocortin 3 in patients with chronic heart failure. *Br J Clin Pharmacol* 2016;82:974–982.
- Patel K, Rademaker MT, Kirkpatrick CM, et al. Comparative pharmacokinetics and pharmacodynamics of urocortins 1, 2 and 3 in healthy sheep. *Br J Pharmacol* 2012;166:1916–1925.
- Gao MH, Lai NC, Miyahara A, et al. Intravenous adeno-associated virus serotype 8 encoding urocortin-2 provides sustained augmentation of left ventricular function in mice. *Hum Gene Ther* 2013;24:777–785.
- Lai NC, Gao MH, Giamouridis D, et al. Intravenous AAV8 encoding urocortin-2 increases function of the failing heart in mice. *Hum Gene Ther* 2015;26:347–356.
- Giamouridis D, Gao MH, Lai NC, et al. Effects of urocortin 2 versus urocortin 3 gene transfer on left ventricular function and glucose disposal. *JACC Basic Transl Sci* 2018;3:249–264.
- Xiao X, Li J, Samulski RJ. Production of high-titer recombinant adeno-associated virus vectors in the absence of helper adenovirus. *J Virol* 1998;72:2224–2232.
- Gao G, Qu G, Burnham MS, et al. Purification of recombinant adeno-associated virus vectors by column chromatography and its performance *in vivo*. *Hum Gene Ther* 2000;11:2079–2091.
- van den Bos EJ, Mees BM, de Waard MC, et al. A novel model of cryoinjury-induced myocardial infarction in the mouse: a comparison with coronary artery ligation. *Am J Physiol Heart Circ Physiol* 2005;289:1291–1230.
- Gao MH, Lai NC, Roth DM, et al. Adenylylcyclase increases responsiveness to catecholamine stimulation in transgenic mice. *Circulation* 1999;99:1618–1622.
- Bayat H, Swaney JS, Ander AN, et al. Progressive heart failure after myocardial infarction in mice. *Basic Res Cardiol* 2002;97:206–213.
- Palakodeti V, Oh S, Oh BH, et al. Force-frequency effect is a powerful determinant of myocardial contractility in the mouse. *Am J Physiol* 1997;273:H1283–H1290.
- Louch WE, Hougen K, Mørk HK, et al. Sodium accumulation promotes diastolic dysfunction in end-stage heart failure following Serca2 knockout. *J Physiol* 2010;588:465–478.
- Messer AE, Jacques AM, Marston SB. Troponin phosphorylation and regulatory function in human heart muscle: dephosphorylation of Ser23/24 on troponin I could account for the contractile defect in end-stage heart failure. *J Mol Cell Cardiol* 2007;42:247–259.
- Salhi HE, Hassel NC, Siddiqui JK, et al. Myofilament calcium sensitivity: mechanistic insight into TnI Ser-23/24 and Ser-150 phosphorylation integration. *Front Physiol* 2016;7:567.
- Battistoni A, Rubattu S, Volpe M. Circulating biomarkers with preventive, diagnostic and prognostic implications in cardiovascular diseases. *Int J Cardiol* 2012;157:160–168.
- Krenz M, Robbins J. Impact of beta-myosin heavy chain expression on cardiac function during stress. *J Am Coll Cardiol* 2004;44:2390–2397.
- Copeland O, Nowak KJ, Laing NG, et al. Investigation of changes in skeletal muscle alpha-actin expression in normal and pathological human and mouse hearts. *J Muscle Res Cell Motil* 2010;31:207–214.
- Chen C, Shen L, Cao S, et al. Cytosolic CARP promotes angiotensin II- or pressure overload-induced cardiomyocyte hypertrophy through calcineurin accumulation. *PLoS One* 2014;9:e104040.
- Shen L, Chen C, Wei X, et al. Overexpression of ankyrin repeat domain 1 enhances cardiomyocyte apoptosis by promoting p53 activation and mitochondrial dysfunction in rodents. *Clin Sci (Lond)* 2015;128:665–678.
- Boutin S, Monteilhet V, Veron P, et al. Prevalence of serum IgG and neutralizing factors against adeno-associated virus (AAV) types 1, 2, 5, 6, 8, and 9 in the healthy population: implications for gene therapy using AAV vectors. *Hum Gene Ther* 2010;21:704–712.
- Clayton JA, Collins FS. Policy: NIH to balance sex in cell and animal studies. *Nature* 2014;509:282–283.

Received for publication May 23, 2018;
accepted after revision July 11, 2018.

Published online: July 12, 2018.