

Metabolic profile before and after short-term overfeeding with a high-fat diet: a comparison between South Asian and white men.

Citation for published version (APA):

Wulan, S. N., Westerterp, K. R., & Plasqui, G. (2014). Metabolic profile before and after short-term overfeeding with a high-fat diet: a comparison between South Asian and white men. *British Journal of Nutrition*, 111(10), 1853-1861. <https://doi.org/10.1017/S0007114514000014>

Document status and date:

Published: 01/01/2014

DOI:

[10.1017/S0007114514000014](https://doi.org/10.1017/S0007114514000014)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Metabolic profile before and after short-term overfeeding with a high-fat diet: a comparison between South Asian and white men

Siti N. Wulan^{1,2*}, Klaas R. Westerterp¹ and Guy Plasqui¹

¹Department of Human Biology, Nutrition and Toxicology Research Institute (NUTRIM) – School for Nutrition Toxicology and Metabolism, Maastricht University Medical Center (MUMC+), Universiteit Single (UNS), 50, PO Box 616, 6200 MD Maastricht, The Netherlands

²Laboratory of Food Quality and Nutrition, Department of Food and Agricultural Product Technology, Faculty of Agricultural Technology, Brawijaya University, Malang, East Java, Indonesia

(Submitted 20 June 2013 – Final revision received 14 November 2013 – Accepted 13 December 2013 – First published online 10 February 2014)

Abstract

For the same BMI, South Asians have a higher body fat percentage and an adverse metabolic profile compared with whites. The objective of the present study was to determine the metabolic profiles of South Asian and white men matched for body fat percentage in response to short-term overfeeding with a high-fat diet. A total of ten healthy non-diabetic South Asian men matched for body fat percentage with ten white men were included in the study. A weight-maintenance diet (containing 30% fat, 55% carbohydrate and 15% protein) was provided for 3 d followed by 4 d of overfeeding (150% of energy requirement) with a high-fat diet (60% fat, 25% carbohydrate and 15% protein). Before and after the overfeeding period, plasma glucose, insulin, TAG, NEFA, total cholesterol and HDL-cholesterol concentrations were determined. Glucose clearance was calculated using a 2 h oral glucose tolerance test. The results revealed that in South Asian and white men, respectively, overfeeding with a high-fat diet decreased plasma TAG concentrations by 0.4 (SD 0.6) and 0.4 (SD 0.5) mmol/l ($P_{\text{diet}} = 0.008$; $P_{\text{ethnicity}} = 0.24$), increased HDL-cholesterol concentrations by 0.12 (SD 0.1) and 0.14 (SD 0.2) mmol/l ($P_{\text{diet}} = 0.001$; $P_{\text{ethnicity}} = 0.06$) and decreased glucose clearance by 48.8 (SD 53.5) and 37.2 (SD 34.2) ml/min per m² body surface ($P_{\text{diet}} = 0.004$; $P_{\text{ethnicity}} = 0.18$). There was a significant interaction between diet and ethnicity with regard to the changes in total and LDL-cholesterol concentrations ($P = 0.01$ and 0.007 , respectively), which trended towards a larger increase in South Asian subjects than in white subjects. Despite a similar body fat percentage, short-term overfeeding with a high-fat diet had more adverse effects on the lipid profile of South Asians than on that of whites.

Key words: Overfeeding; High-fat diet; South Asian men; White men

The prevalence of energy imbalance and related obesity is rapidly increasing worldwide, including in a number of Asian countries^(1–3). This is linked to an increase in the prevalence of non-communicable diseases such as diabetes, CVD and hypertension⁽¹⁾. It is estimated that 300 million people will have diabetes by the year 2025 and that among them more than 100 million people live in Asia⁽⁴⁾. However, diabetes in Asia has developed in a shorter time period (3–5-fold increase within 30 years), in a younger age group (45–64 years old) and in people with a lower BMI compared with that in Western countries⁽⁵⁾.

Compared with whites with the same BMI and of the same age and sex, Asians have been found to have a higher body fat percentage and a lower fat-free mass (FFM)/appendicular skeletal muscle mass, as has been reported recently by us⁽⁶⁾ and by others^(7–13). Among Asians, South Asians (people from

the Indian subcontinent) have the most pronounced difference in body fat compared with whites⁽¹⁴⁾. Metabolic consequences, as revealed by the prevalence of the metabolic syndrome, are also greater than those in other Asian countries^(4,5). Raji *et al.*⁽¹⁵⁾ reported that even at a normal BMI Asian Indians exhibit clustering of cardiovascular risk factors such as insulin resistance, dyslipidaemia and pro-coagulant state.

A strong gene–environment interaction, which is triggered by lifestyle changes due to modernisation, is suggested to be the cause of the rapid increase in the prevalence of diabetes in Asia^(3,16). Furthermore, migrant Asian groups have also been reported to have higher susceptibility to adverse environmental influences than co-inhabitants of different races⁽¹⁶⁾. Major dietary changes such as a large increase in the consumption of fats and added sugars in the

Abbreviations: FFM, fat-free mass; LPL, lipoprotein lipase; OGTT, oral glucose tolerance test; TEE, total energy expenditure.

* **Corresponding author:** S. N. Wulan, fax +31 433670976, email siti.wulan@maastrichtuniversity.nl

diet have accelerated in Asia in recent decades⁽¹⁷⁾. Distinct genetic background represented partly by differences in body composition between South Asians and whites may contribute to the differences in metabolic responses when they are exposed to a high-fat diet challenge. It has been shown that a high-fat diet results in liver fat accumulation in a short time period⁽¹⁸⁾, leading to metabolic complications⁽¹⁹⁾. In the present study, South Asians and whites were matched for body fat percentage, which is one of the confounding factors affecting metabolic profiles in individuals. The objective was to determine the effect of short-term overfeeding with a high-fat diet on the metabolic profile of South Asians and whites.

Subjects and methods

Subjects

A total of ten healthy non-diabetic South Asian men and ten white men were included in the study. They were matched for body fat percentage on an individual basis; as a consequence, the variation in body fat percentage was comparable between the groups. The number of subjects was determined based on a previous study that detected a significant reduction in insulin sensitivity after an intervention with a high-fat diet in ten white subjects⁽²⁰⁾. Subject characteristics are given in Table 1. Asian subjects had four grandparents from South Asia, while white subjects were non-Hispanic Europeans. Subjects were selected based on the following inclusion criteria: healthy; not having metabolic diseases (diabetes or CVD); not using medications; aged between 20 and 40 years with BMI ranging between 18 and 29 kg/m² for South Asians and 22 and 33 kg/m² for whites; having a stable body weight (as defined by weight change <5 kg) for the last 3 months before participating in the study; not being on a diet; not

being an athlete. Verbal and written information about the study was given to all subjects before obtaining their consent. Written informed consent was obtained from all subjects. The study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects were approved by the Medical Ethics Committee of Maastricht University Medical Centre, MEC No. 10-3-013, and registered in the public trial registry (www.ccmo.nl no. NL31217.068.10).

Experimental protocol

The present study was a dietary intervention study carried out in a free-living condition. Body composition was measured before the start of the dietary intervention to ensure that the body fat percentage of the two ethnic groups matched. Cardiorespiratory fitness (defined as VO₂max corrected for FFM) was measured before the start of the dietary intervention. Daily physical activity levels were measured for seven consecutive days with an accelerometer. Energy requirements of a weight-maintenance diet for 3 d were calculated based on FFM and daily physical activity levels of each subject (see below). The metabolic profiles were determined just before and after overfeeding with the high-fat diet for 4 d. All measurements were carried out at the Metabolic Research Unit Maastricht (MRUM), Maastricht University, Maastricht, The Netherlands.

Determination of body composition

Body composition was determined according to a three-compartment model based on body weight, body volume and total body water. Body weight and body volume were determined in the morning in a fasting state. Body volume was determined by hydrodensitometry with simultaneous measurement of residual lung capacity using the helium dilution technique. Total body water was determined using the ²H dilution method according to the Maastricht protocol⁽²¹⁾. Body composition was calculated from body density and total body water using the equation of Siri⁽²²⁾.

Assessment of cardiorespiratory fitness

Physical fitness was assessed with an incremental test on a bicycle ergometer using the protocol of Kuipers *et al.*⁽²³⁾. During the test, VO₂ and CO₂ production were measured continuously and heart rate was monitored (Polar heart rate monitor; Polar Electro Oy). After 5 min of warm-up at 100 W, the workload was increased by 50 W every 2.5 min. When the heart rate reached a value of 35 beats/min below the age-predicted maximal heart rate (220 beats/min – age) or the respiratory quotient (RQ = CO₂ production/VO₂) exceeded 1, the workload was increased by 25 W every 2.5 min until exhaustion. VO₂max was determined by averaging the last few points of maximum VO₂. Cardiorespiratory fitness was defined as VO₂max corrected for FFM.

Table 1. Subject characteristics
(Mean values and standard deviations)

Characteristics	South Asians (n 10)		Whites (n 10)		P*
	Mean	SD	Mean	SD	
Age (years)	27	2	24	2	0.001
Weight (kg)	68.9	7.4	88.3	18.1	0.006
Height (m)	1.72	0.06	1.80	0.08	0.02
BMI (kg/m ²)	23.3	3.0	27.0	4.2	0.04
Fat mass (%)	25.5	6.4	23.7	7.5	0.58
FFM (kg)	51.0	3.7	66.5	8.5	0.001
Waist circumference (cm)	85.9	7.7	96.0	12.0	0.04
Hip circumference (cm)	97.0	5.5	105.4	8.7	0.02
Waist:hip ratio	0.9	0.04	0.91	0.06	0.31
Waist:height ratio	0.50	0.05	0.53	0.06	0.23
VO ₂ max corrected for FFM (ml O ₂ /min)	2780	265	3249	152	0.15
Physical activity accelerometer (10 ³ counts/d)	1123	239	1304	442	0.27
Daily PAL	1.64	0.06	1.69	0.11	0.27

FFM, fat-free mass; PAL, physical activity level.

* Differences between the groups were assessed using an independent *t* test.

Measurement of daily physical activity levels

Daily physical activity levels were measured using a Direct Life triaxial accelerometer for movement registration (Tracmor-D; Philips NewWellness Solutions; <http://www.directlife.philips.com>). The device is a small (3.2 × 3.2 × 0.5 cm), lightweight (12.5 g) instrument. The accelerometer was attached to the lower back of the subjects by means of an elastic belt. It registered accelerations minute by minute, in the mediolateral (*x*-axis), longitudinal (*y*-axis) and anterioposterior (*z*-axis) positions of the trunk as described elsewhere⁽²⁴⁾.

The subjects were instructed to wear the accelerometer for seven consecutive days during waking hours, except during water activities. The subjects were advised to maintain their habitual physical activity levels during the dietary intervention period. Tracmor-D output is expressed as activity counts/min. To determine the average Tracmor-D counts/d, Tracmor-D activity counts/min were summed over the entire monitoring period and divided by the number of monitoring days.

Daily physical activity levels (PAL) were calculated based on the activity counts/d using the following formula⁽²⁴⁾:

$$\text{PAL} = 1.354 + 256 \times 10^{-9} \times \text{counts/d.}$$

Total energy expenditure (TEE) was calculated using the formula of Bonomi *et al.*⁽²⁴⁾, by including activity counts/d (from the accelerometer) and FFM:

$$\text{TEE accelerometer} = 0.04 + 0.17 \text{ FFM} + 1.67 \times 10^{-6} \times \text{counts/d.}$$

Determination of energy intake

The energy balance (weight-maintenance) diet to be consumed at home for 3 d before the baseline measurement was calculated on the basis of TEE of each subject as calculated above. The macronutrient composition of the energy balance diet before the baseline measurement was 30 % fat, 55 % carbohydrate and 15 % protein.

The high-fat diet was formulated to have 50 % excess energy above the TEE of subjects when on the balance diet⁽²⁵⁾. The macronutrient composition of the high-fat diet was 60 % fat, 25 % carbohydrate and 15 % protein^(26,27). The fatty acid composition of the diet was 40 % SFA and 60 % unsaturated fatty acids.

A written instruction was given to prepare the diet at home. During the balance diet period, subjects were provided with amounts of the diet that exceeded their TEE and were allowed to eat more or less than the amounts prescribed, according to their preference (*ad libitum*). Any additional intake from the prescribed foods was recorded. All unfinished foods were collected and returned to the university to calculate the actual energy intake. During the overfeeding period, the subjects were asked to completely consume all the foods prescribed, but on failing to do so, to record their intake and return the unfinished foods. The diet consisted of normal ready-to-eat foods combining typical Western and Asian diets. Foods were selected by reviewing the ingredient content to ensure that there was no (or only a minimal) effect of certain ingredients on fat oxidation (such as spices). During the overfeeding

period, the subjects were also provided with decaffeinated coffee and fruit tea, as caffeine has been reported to increase fat oxidation. Alcohol intake was limited during the dietary intervention period (only one serving/d when needed). During the dietary intervention period, the subjects were asked to wear the accelerometer and to maintain their habitual physical activity levels; this allowed us to monitor the energy balance of the subjects during the entire dietary intervention period.

Oral glucose tolerance test

A 2 h oral glucose tolerance test (OGTT) was carried out before and after overfeeding with the high-fat diet. After an overnight fast, the subjects underwent an OGTT test. A fasting blood sample was collected ($t = 0$), after which the subjects drank a glucose solution (82.5 g glucose monohydrate dissolved in 600 ml water) within 5 min⁽²⁸⁾. This method is a modification of the American Diabetes Association (ADA) guidelines⁽²⁹⁾, as it has been reported to improve overall tolerability and have only a minor impact on the intra-subject variation of plasma glucose and insulin responses with no effect on the diagnostically relevant 2 h endpoint irrespective of the body composition⁽²⁸⁾. Venous blood samples were collected every 15 min in the first hour ($t = 15, 30, 45$ and 60) and every 30 min in the second hour ($t = 90$ and 120). Plasma glucose concentrations were measured to determine glucose tolerance according to the ADA criteria⁽³⁰⁾.

Venous plasma glucose and insulin concentrations determined during the OGTT were used to assess pancreatic β -cell function and insulin sensitivity. These parameters were assessed using the updated homeostasis model assessment⁽³¹⁾ and the oral glucose insulin sensitivity index⁽³²⁾, respectively. The oral glucose insulin sensitivity index is a measure of insulin sensitivity derived from glucose clearance in the OGTT. It is expressed as glucose clearance (ml/min) per m² body surface by taking into account the subject's body weight and height (body surface), the dose of glucose solution, the concentration of plasma glucose at time 0, 90 and 120 min, and the concentration of insulin at time 0 and 90 min⁽²⁹⁾.

Blood analysis

Blood samples were collected in tubes containing 30 μ l of 0.2 M-EDTA. Plasma samples were immediately centrifuged at 3000 rpm for 10 min, frozen in liquid N₂ and stored at -80°C until analysis. Glucose (Roche) concentrations were determined enzymatically. Insulin concentrations were determined using a RIA (Linco Research). Fasting blood lipid profiles were measured as follows: NEFA using the Wako Nefa C test kit; TAG with correction for free glycerol (Sigma Diagnostics); total cholesterol (Roche) using the oxidase phenol 4-aminoantipyrene peroxidase method. HDL-cholesterol concentrations were measured using the precipitation method, while LDL-cholesterol concentrations were determined using the Friedewald equation⁽³³⁾.

Statistical analysis

Data are presented as means and standard deviations. Statistical comparison of the subjects' baseline characteristics, energy intake and macronutrient composition of the diet consumed between the ethnic groups was made using an independent-samples *t* test. Repeated-measures ANOVA were carried out to assess within-group and between-group differences in parameters before and after overfeeding with the high-fat diet as well as the interaction between ethnicity and diet. The SPSS program version 16 (SPSS) was used for statistical analysis, and the statistical significance was set at $P < 0.05$.

Results

Subject characteristics

South Asian subjects were of Indian (n 8) and Pakistani (n 2) origins. White subjects were of Dutch (n 3), German (n 2), French (n 1), British (n 1), Danish (n 1), Polish (n 1) and Icelander (n 1) origins. For South Asian subjects, measurements were carried out within 3 years (n 5) and within 1 year (n 5) of their stay in The Netherlands. Subject characteristics are given in Table 1. South Asian subjects were slightly older than white subjects. South Asian subjects had a significantly lower BMI than white subjects (23.3 (SD 3.0) *v.* 27.0 (SD 4.2) kg/m²; $P = 0.04$), but there was no difference in body fat percentage (25.5 (SD 6.4) and 23.7 (SD 7.5)% for South Asian and white subjects, respectively; $P = 0.58$). The FFM in South Asian subjects was lower than that in white subjects ($P = 0.001$). White subjects had significantly higher waist circumference and hip circumference than South Asian subjects, resulting in no difference being observed in the waist:hip ratio between the groups ($P = 0.31$). Cardiorespiratory fitness, defined as VO_2max corrected for FFM, did not differ between the ethnic groups ($P = 0.15$), and there was no difference in daily physical activity counts ($P = 0.27$). Additionally, in the South Asian group, of the ten subjects, a family history of diabetes, dyslipidaemia and CVD was found in four, five and four subjects, respectively. In the white group, of the ten subjects, a family

history of diabetes, dyslipidaemia and CVD was found in three, five and five subjects, respectively.

Energy intake and macronutrient composition during the dietary intervention period

Compliance with the dietary intervention is summarised in Table 2. White subjects had a significantly higher energy requirement due to a higher FFM and therefore had a significantly higher energy intake. During the overfeeding period, South Asian subjects consumed 143 (SD 12)% of their energy requirement, whereas white subjects consumed 152 (SD 16)% of their energy requirement ($P = 0.16$). There was no difference in the macronutrient composition of the diet consumed between South Asian and white subjects. The percentage of fat intake (as % energy intake) during the balance diet period was 29 (SD 1.0) and 28.8 (SD 0.5)% ($P = 0.33$) for South Asian and white subjects, respectively. The percentage of fat intake during the overfeeding period was 59.3 (SD 1.3) and 58.8 (SD 2.3)% ($P = 0.53$) for South Asian and white subjects, respectively. There was no difference in the fatty acid composition of the diet consumed between South Asian and white subjects ($P = 0.49$).

Plasma parameters

Plasma parameters are given in Table 3. Short-term overfeeding with the high-fat diet decreased fasting TAG concentrations ($P = 0.008$), with no difference being observed between the ethnic groups ($P = 0.24$), and increased HDL-cholesterol concentrations ($P = 0.001$), which tended to be higher in white subjects ($P = 0.06$). There was a significant interaction between diet and ethnicity with regard to the changes in total cholesterol ($P = 0.01$) and LDL-cholesterol ($P = 0.007$) concentrations, which trended towards a larger increase in South Asian subjects than in white subjects, whereas fasting NEFA concentrations remained unaffected.

There was no effect of overfeeding with the high-fat diet on fasting glucose and fasting insulin concentrations and hence

Table 2. Energy intake and macronutrient composition during the weight-maintenance and dietary intervention periods (Mean values and standard deviations)

	South Asians (n 10)		Whites (n 10)		P^*
	Mean	SD	Mean	SD	
Energy balance (weight-maintenance) diet period					
Energy intake (kJ/d)	10800	1000	12900	2200	0.01
Fat (% energy intake)	29.1	1.0	28.8	0.5	0.33
Carbohydrate (% energy intake)	55.6	1.3	55.6	0.6	0.99
Protein (% energy intake)	15.3	0.6	15.6	0.6	0.19
Overfeeding period					
Energy intake (kJ/d)	15400	1300	19500	2400	0.001
Percentage of overfeeding (% from energy balance)	143	12	152	16	0.16
Fat intake (% energy intake)	59.3	1.3	58.8	2.3	0.53
Carbohydrate (% energy intake)	26.2	1.5	26.4	2.3	0.85
Protein (% energy intake)	14.5	0.3	14.9	0.7	0.20
SFA (% fat intake)	40.9	1.1	40.5	1.2	0.49
Unsaturated fatty acids (% fat intake)	59.1	1.1	59.5	1.2	0.49

* Differences between the groups were assessed using an independent *t* test.

Table 3. Plasma parameters*
(Mean values and standard deviations)

Parameters	South Asians (n 10)				Whites (n 10)				P†		
	Before OHFD		After OHFD		Before OHFD		After OHFD				
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Diet	Ethnicity	Diet × ethnicity
NEFA ($\mu\text{mol/l}$)	391.8	103.9	461.2	149.1	420.0	168.9	388.6	95.3	0.60	0.65	0.17
TAG (mmol/l)	1.5	0.9	1.1	0.5	1.2	0.6	0.8	0.3	0.008	0.24	0.77
Total chol (mmol/l)	5.7	0.9	6.1	1.0	5.5	0.9	5.2	0.8	0.51	0.14	0.01
HDL-chol (mmol/l)	1.1	0.3	1.2	0.3	1.4	0.4	1.5	0.4	0.001	0.063	0.71
LDL-chol (mmol/l)	3.9	0.8	4.4	0.7	3.5	1.0	3.3	0.9	0.29	0.058	0.007
HDL-chol:total chol ratio	0.19	0.04	0.2	0.04	0.26	0.1	0.3	0.1	0.001	0.03	0.008
LDL-chol:total chol ratio	0.68	0.05	0.72	0.03	0.64	0.1	0.63	0.1	0.32	0.046	0.04
Glucose (mmol/l)	5.1	0.5	5.2	0.4	5.0	0.4	5.0	0.3	0.99	0.43	0.55
Insulin (pmol/l)	124.7	44.9	119.7	37.1	123.3	66.3	130.4	69.8	0.80	0.86	0.25
HOMA index	4.1	1.8	3.9	1.4	3.7	2.1	3.9	2.2	0.94	0.84	0.43
OGIS index (ml/min per m^2)‡	387.5	107.2	338.7	61.4	435.9	63.1	398.7	53.2	0.004	0.18	0.64

OHFD, overfeeding with a high-fat diet; chol, cholesterol; HOMA, homeostasis model assessment; OGIS, oral glucose insulin sensitivity.

* Plasma parameters were determined in a fasting condition, except for the OGIS index, which was derived during the 2 h oral glucose tolerance test (OGTT).

† Differences in the changes in parameters within the groups (P_{diet}) and between the groups ($P_{\text{ethnicity}}$) and the interaction between diet and ethnicity were assessed using repeated-measures ANOVA.

‡ For the OGIS index (glucose clearance, ml/min per m^2 body surface), data were based on seven South Asians and seven whites, matched for body fat percentage (25.9 (SD 7.1) and 24.4 (SD 8.4) %, $P=0.72$, for South Asians and whites respectively), who had completed the OGTT.

there were no differences in the homeostasis model assessment index before and after overfeeding with the high-fat diet ($P=0.94$) and no differences between the ethnic groups ($P=0.84$). Overfeeding with the high-fat diet decreased insulin sensitivity ($P=0.004$), expressed as the oral glucose insulin sensitivity index, which was calculated from glucose clearance in 2 h OGTT (ml/min) per m^2 body surface. There was no difference in the changes in glucose clearance between the ethnic groups ($P=0.18$). Fig. 1(a) and (b) shows that the glucose response during the 2 h OGTT was not affected by the diet ($P=0.16$) and did not differ between the ethnic groups ($P=0.6$), whereas the insulin response (Fig. 1(c) and (d)) increased significantly with overfeeding ($P=0.046$), but no difference was observed between the ethnic groups ($P=0.23$).

Discussion

In the present study, two ethnic groups were matched for body fat percentage, confirming the evidence that South Asian men have a lower BMI for the same body fat percentage when compared with white men. Despite a similar body fat percentage, South Asian men had a more adverse lipid profile, i.e. a larger increase in total cholesterol and LDL-cholesterol concentrations as a response to overfeeding with the high-fat diet, whereas glucose clearance (insulin sensitivity) decreased similarly in both ethnic groups.

Cross-sectional studies comparing South Asians and whites have consistently reported an unfavourable metabolic profile such as dyslipidaemia in South Asians when matched or adjusted for BMI, sex and age^(34–36), which is expected, given that the South Asians have a higher body fat percentage at the same BMI than whites. In the present study, subjects were matched for body fat percentage, and the baseline metabolic profile measured before the overfeeding period was comparable between the ethnic groups, except for the HDL-cholesterol:total cholesterol ratio, which tended to be lower

in South Asian subjects. Another study has reported comparable fasting TAG, total cholesterol, HDL-cholesterol and LDL-cholesterol concentrations but higher plasma NEFA concentrations in South Asian Indians compared with body fat percentage-matched white men⁽³⁷⁾.

In the present study, fasting plasma NEFA concentrations were unaffected by overfeeding with the high-fat diet, and there was no difference between the ethnic groups. In the fasting state, lipolysis is activated and fatty acids released by the action of hormone-sensitive lipase are directed primarily into the venous plasma⁽³⁸⁾. In the fed state, lipoprotein lipase (LPL) is activated and the fatty acids released from circulating TAG are directed into the tissue for esterification and storage in white adipose tissue⁽³⁸⁾. In the overfeeding condition, the postprandial (fed) state may be slightly extended as a result of extended meal ingestion; the release of NEFA from subcutaneous fat presumably declines due to the prolonged presence of high insulin concentrations. However, it is a common observation either during infusion of TAG emulsions or after ingestion of high-fat meals⁽³⁹⁾ that the plasma NEFA concentration is relatively maintained, despite an elevation in insulin concentrations⁽³⁸⁾. In addition, after the consumption of a high-fat meal, both the inward flux of fatty acids from the plasma into adipose tissue and the release of LPL-derived NEFA into the venous plasma are greater than those after the consumption of a lower-fat mixed meal⁽⁴⁰⁾. LPL appears to operate continuously, generating a pool of fatty acids, of which some are always released into the venous plasma, while a proportion are diverted into the tissue for esterification and storage in the postprandial state⁽³⁸⁾.

We observed a decrease in fasting TAG concentrations after short-term overfeeding with the high-fat diet, but the decrease did not differ between the ethnic groups. Diets that replace fat with carbohydrate isoenergetically have been consistently observed to worsen certain elements of the plasma lipid profile⁽⁴¹⁾, particularly TAG^(42,43). In a multiethnic




Fig. 1. Glucose (a) and insulin (c) responses during overfeeding with a high-fat diet (OHFD) in South Asians (before, ○; after, ●) and whites (before, □; after, ■) and the corresponding AUC of glucose (b) and insulin (d) during 2 h oral glucose tolerance test (OGTT) (□, before OHFD; ■, after OHFD). Values are means and standard deviations represented by vertical bars. Differences within and between the groups after the dietary intervention were assessed using repeated-measures ANOVA. * Mean value obtained after OHFD was significantly different from that obtained before OHFD for South Asian subjects. † Mean value obtained after OHFD was significantly different from that obtained before OHFD for white subjects. Data were available for fourteen subjects, seven subjects in each group matched for body fat (25.9 (sd 7.1)% and 24.4 (sd 8.4)% for South Asian and white subjects, respectively; $P=0.72$), who had completed the OGTT: AUC glucose – $P_{\text{diet}}=0.16$, $P_{\text{ethnicity}}=0.60$, and $P_{\text{diet} \times \text{ethnicity}}=0.53$; AUC insulin – $P_{\text{diet}}=0.046$, $P_{\text{ethnicity}}=0.23$, and $P_{\text{diet} \times \text{ethnicity}}=0.53$.

population study in Canada, Merchant *et al.*⁽⁴³⁾ reported that South Asian Indians consumed large amounts of carbohydrate followed by Europeans, Aboriginals and Chinese. In that study, higher carbohydrate intake was found to be associated with higher fasting TAG concentrations and those with the highest percentile of carbohydrate intake were found to have the highest fasting TAG concentrations. In the present study, reduction in fasting plasma TAG concentrations may be attributed to a lower proportion of carbohydrate in the high-fat diet (25% energy from carbohydrate in the overfeeding period equals 37.5% carbohydrate compared with 55% carbohydrate in the balance diet period), as also reported by others^(44–46). High-carbohydrate diets are hypothesised to increase TAG concentrations by inducing fatty acid production in the liver and inhibiting the action of LPL through increased apo CIII production, particularly in the presence of insulin resistance⁽⁴⁷⁾. Higher fat intakes may increase circulating chylomicron concentrations, but do not affect LPL and the clearance of circulating TAG and hence do not affect TAG concentrations⁽⁴⁸⁾.

A significant interaction between diet and ethnicity was found with regard to the changes in total cholesterol and LDL-cholesterol concentrations, which trended towards a larger increase in South Asian subjects than in white subjects.

In addition, overfeeding with the high-fat diet increased HDL-cholesterol concentrations, but the increase tended to be higher in white subjects than in South Asian subjects. Furthermore, the changes in HDL-cholesterol:total cholesterol ratio and LDL-cholesterol:total cholesterol ratio were different between the ethnic groups. Thus, overfeeding with the high-fat diet clearly resulted in a more unfavourable lipoprotein profile in South Asian subjects. The effect of a high-fat diet on lipoproteins varies considerably depending on the type of fat consumed, i.e. fatty acid composition of the diet⁽⁴⁹⁾. SFA increase the concentrations of both total cholesterol and LDL-cholesterol, while unsaturated fatty acids lower the concentrations of total cholesterol and LDL-cholesterol⁽⁴⁹⁾. Both SFA and unsaturated fatty acids increase HDL-cholesterol concentrations, although this increase is greater with SFA⁽⁴⁹⁾. A recent review by Hooper *et al.*⁽⁵⁰⁾ has reported that reduced and modified fat intakes decrease total cholesterol and LDL-cholesterol concentrations, with no effect on HDL-cholesterol concentrations. In the present study, during the overfeeding period, the subjects were given a diet containing 60% energy from fat, of which 40% of the fat consumed was from SFA and 60% was from unsaturated fatty acids. No differences in the adherence to the diet and in fat intake and fatty acid composition were observed between the ethnic groups.

Given that the subjects in the present study had a comparable range of body fat percentages and comparable metabolic profiles at baseline except for HDL-cholesterol, which tended to be present at lower concentrations in South Asian subjects, the unfavourable lipoprotein profile in South Asian subjects as a response to the high-fat diet needs to be investigated further. This may possibly be due to differences in the rate of LDL clearance and production⁽⁵¹⁾. Furthermore, data obtained from thirty-one study cohorts of twelve countries revealed that South Asian Indian men and women have the lowest HDL-cholesterol concentrations compared with Central and Northern Europeans, Japanese and Chinese across the glucose level categories⁽⁵²⁾.

Overfeeding^(53,54) and a high-fat diet⁽⁵⁵⁾ have been reported to be associated with disturbances in glucose metabolism and insulin sensitivity. As expected, short-term overfeeding with the high-fat diet increased the insulin response (AUC insulin) during the 2 h OGTT. We also calculated the glucose clearance (ml/min) per m² body surface, which allowed us to correct for the differences in body height between South Asian and white subjects (indirectly FFM). The result remains that short-term overfeeding with the high-fat diet decreased glucose clearance (ml/min) per m² body surface, with no difference being observed between the ethnic groups. In response to a meal, insulin suppresses lipolysis and the release of NEFA into the circulation and stimulates skeletal muscle glucose uptake⁽⁵⁶⁾. *In vivo*, elevation of NEFA concentrations induced by intralipid plus heparin infusion has consistently been shown to reduce glucose oxidation and glucose uptake in the skeletal muscle⁽⁵⁷⁾. Others have suggested the importance of CD36 in NEFA transport and ectopic TAG accumulation; incomplete oxidation generates intermediates that can impair insulin signalling⁽⁵⁸⁾.

A limitation of the present study is the low number of subjects, which may not reflect the general population of South Asians and whites. Additionally, the short-term dietary intervention may reflect an early response to overfeeding with a high-fat diet. Although an early response may precede the metabolic complications in the long term, it cannot be simply extrapolated to assess the response in a longer term. However, this is a well-controlled dietary intervention study and, to our knowledge, no such studies comparing two ethnic groups have been carried out.

In summary, we observed a more adverse lipid profile in South Asian men compared with white men as a response to short-term overfeeding with a high-fat diet, whereas insulin sensitivity decreased similarly. Although the ethnic groups differed with respect to age in the present study, a regression analysis to assess the predictors of baseline metabolic profiles by including age, fat mass percentage and ethnicity revealed that age did not have a significant effect in the groups studied, presumably because all of them were young and the age difference between the groups was small. The differences in the lipid profile may be associated with differences in body fat distribution towards a more central fat deposition⁽⁵⁹⁾ in South Asians. Goel *et al.*⁽⁶⁰⁾ reported that subcutaneous abdominal fat is associated with the metabolic syndrome independent of visceral fat in Asian Indians. These results may

partly explain the high prevalence of the metabolic syndrome in South Asians, which is triggered by lifestyle changes, i.e. increased consumption of high-energy-density foods.

Acknowledgements

The authors thank Jos Stegen, Loek Wouters and Paul Schoffelen for their technical assistance and help with the analyses. They also deeply appreciate and thank all subjects who participated in the study.

The present study was partly funded by a scholarship from the Directorate General of Higher Education (DGHE), the Ministry of Education and Culture of the Republic of Indonesia, to S. N. W. The Ministry of Education and Culture of the Republic of Indonesia had no role in the design and analysis of the study and in the writing of this article.

The authors' contributions are as follows: S. N. W. conducted the research, carried out the data analysis and wrote the manuscript under the supervision of K. R. W. and G. P., who designed the study, interpreted the data and reviewed the manuscript.

None of the authors has any conflicts of interest to declare.

References

1. Popkin BM (2006) Global nutrition dynamics: the world is shifting rapidly toward a diet linked with noncommunicable diseases. *Am J Clin Nutr* **84**, 289–298.
2. Subramanian SV, Perkins JM, Ozaltin E, *et al.* (2011) Weight of nations: a socioeconomic analysis of women in low- to middle-income countries. *Am J Clin Nutr* **93**, 413–421.
3. Misra A & Khurana L (2008) Obesity and the metabolic syndrome in developing countries. *J Clin Endocrinol Metab* **93**, S9–S30.
4. King H, Aubert RE & Herman WH (1998) Global burden of diabetes, 1995–2025: prevalence, numerical estimates, and projections. *Diabetes Care* **21**, 1414–1431.
5. Yoon KH, Lee JH, Kim JW, *et al.* (2006) Epidemic obesity and type 2 diabetes in Asia. *Lancet* **368**, 1681–1688.
6. Wulan SN, Westerterp KR & Plasqui G (2012) Dietary and 24-h fat oxidation in Asians and whites who differ in body composition. *Am J Clin Nutr* **95**, 1335–1341.
7. Deurenberg P & Deurenberg-Yap M (2001) Differences in body-composition assumptions across ethnic groups: practical consequences. *Curr Opin Clin Nutr Metab Care* **4**, 377–383.
8. Deurenberg P & Deurenberg-Yap M (2003) Validity of body composition methods across ethnic population groups. *Forum Nutr* **56**, 299–301.
9. Deurenberg-Yap M, Chew SK & Deurenberg P (2002) Elevated body fat percentage and cardiovascular risks at low body mass index levels among Singaporean Chinese, Malays and Indians. *Obes Rev* **3**, 209–215.
10. Chang CJ, Wu CH, Chang CS, *et al.* (2003) Low body mass index but high percent body fat in Taiwanese subjects: implications of obesity cutoffs. *Int J Obes Relat Metab Disord* **27**, 253–259.
11. Rush EC, Goedecke JH, Jennings C, *et al.* (2007) BMI, fat and muscle differences in urban women of five ethnicities from two countries. *Int J Obes (Lond)* **31**, 1232–1239.

12. Sampei MA, Novo NF, Juliano Y, *et al.* (2008) Anthropometry and body composition in ethnic Japanese and Caucasian adolescent boys. *Pediatr Int* **50**, 679–686.
13. Rush EC, Freitas I & Plank LD (2009) Body size, body composition and fat distribution: comparative analysis of European, Maori, Pacific Island and Asian Indian adults. *Br J Nutr* **102**, 632–641.
14. Deurenberg-Yap M, Schmidt G, van Staveren WA, *et al.* (2001) Body fat measurement among Singaporean Chinese, Malays and Indians: a comparative study using a four-compartment model and different two-compartment models. *Br J Nutr* **85**, 491–498.
15. Raji A, Seely EW, Arky RA, *et al.* (2001) Body fat distribution and insulin resistance in healthy Asian Indians and Caucasians. *J Clin Endocrinol Metab* **86**, 5366–5371.
16. Ramachandran A, Ma RC & Snehalatha C (2010) Diabetes in Asia. *Lancet* **375**, 408–418.
17. Popkin BM (2001) The nutrition transition and obesity in the developing world. *J Nutr* **131**, 871S–873S.
18. van der Meer RW, Hammer S, Lamb HJ, *et al.* (2008) Effects of short-term high-fat, high-energy diet on hepatic and myocardial triglyceride content in healthy men. *J Clin Endocrinol Metab* **93**, 2702–2708.
19. Brons C, Jacobsen S, Hiscock N, *et al.* (2012) Effects of high-fat overfeeding on mitochondrial function, glucose and fat metabolism, and adipokine levels in low-birth-weight subjects. *Am J Physiol Endocrinol Metab* **302**, E43–E51.
20. Westerbacka J, Lammi K, Hakkinen AM, *et al.* (2005) Dietary fat content modifies liver fat in overweight nondiabetic subjects. *J Clin Endocrinol Metab* **90**, 2804–2809.
21. Westterterp KR, Wouters L & van Marken Lichtenbelt WD (1995) The Maastricht protocol for the measurement of body composition and energy expenditure with labeled water. *Obes Res* **3**, Suppl. 1, 49–57.
22. Siri WE (1993) Body composition from fluid spaces and density: analysis of methods. 1961. *Nutrition* **9**, 480–492, discussion 480, 491.
23. Kuipers H, Keizer HA, Verstappen FT, *et al.* (1985) Influence of a prostaglandin-inhibiting drug on muscle soreness after eccentric work. *Int J Sports Med* **6**, 336–339.
24. Bonomi AG, Plasqui G, Goris AH, *et al.* (2010) Estimation of free-living energy expenditure using a novel activity monitor designed to minimize obtrusiveness. *Obesity (Silver Spring)* **18**, 1845–1851.
25. Joosen AM, Bakker AH, Zorenc AH, *et al.* (2006) PPAR-gamma activity in subcutaneous abdominal fat tissue and fat mass gain during short-term overfeeding. *Int J Obes (Lond)* **30**, 302–307.
26. Schrauwen-Hinderling VB, Kooi ME, Hesselink MK, *et al.* (2005) Intramyocellular lipid content and molecular adaptations in response to a 1-week high-fat diet. *Obes Res* **13**, 2088–2094.
27. Schrauwen P, van Marken Lichtenbelt WD, Saris WH, *et al.* (1997) Changes in fat oxidation in response to a high-fat diet. *Am J Clin Nutr* **66**, 276–282.
28. Sevenpiper JL, Jenkins DJ, Josse RG, *et al.* (2001) Dilution of the 75-g oral glucose tolerance test improves overall tolerability but not reproducibility in subjects with different body compositions. *Diabetes Res Clin Pract* **51**, 87–95.
29. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus (1997) Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* **20**, 1183–1197.
30. American Diabetes Association (2010) Standards of medical care in diabetes – 2010. *Diabetes Care* **33**, Suppl. 1, S11–S61.
31. Levy JC, Matthews DR & Hermans MP (1998) Correct homeostasis model assessment (HOMA) evaluation uses the computer program. *Diabetes Care* **21**, 2191–2192.
32. Mari A, Pacini G, Murphy E, *et al.* (2001) A model-based method for assessing insulin sensitivity from the oral glucose tolerance test. *Diabetes Care* **24**, 539–548.
33. Friedewald WT, Levy RI & Fredrickson DS (1972) Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* **18**, 499–502.
34. Forouhi NG, Jenkinson G, Thomas EL, *et al.* (1999) Relation of triglyceride stores in skeletal muscle cells to central obesity and insulin sensitivity in European and South Asian men. *Diabetologia* **42**, 932–935.
35. Anand SS, Tarnopolsky MA, Rashid S, *et al.* (2011) Adipocyte hypertrophy, fatty liver and metabolic risk factors in South Asians: the Molecular Study of Health and Risk in Ethnic Groups (mol-SHARE). *PLoS One* **6**, e22112.
36. Chandalia M, Lin P, Seenivasan T, *et al.* (2007) Insulin resistance and body fat distribution in South Asian men compared to Caucasian men. *PLoS One* **2**, e812.
37. Abate N, Chandalia M, Snell PG, *et al.* (2004) Adipose tissue metabolites and insulin resistance in nondiabetic Asian Indian men. *J Clin Endocrinol Metab* **89**, 2750–2755.
38. Frayn KN, Shadid S, Hamlani R, *et al.* (1994) Regulation of fatty acid movement in human adipose tissue in the postabsorptive-to-postprandial transition. *Am J Physiol* **266**, E308–E317.
39. Karpe F, Olivecrona T, Walldius G, *et al.* (1992) Lipoprotein lipase in plasma after an oral fat load: relation to free fatty acids. *J Lipid Res* **33**, 975–984.
40. Frayn KN, Coppack SW, Fielding BA, *et al.* (1995) Coordinated regulation of hormone-sensitive lipase and lipoprotein lipase in human adipose tissue *in vivo*: implications for the control of fat storage and fat mobilization. *Adv Enzyme Regul* **35**, 163–178.
41. Hellerstein MK (2002) Carbohydrate-induced hypertriglyceridemia: modifying factors and implications for cardiovascular risk. *Curr Opin Lipidol* **13**, 33–40.
42. Parks EJ & Hellerstein MK (2000) Carbohydrate-induced hypertriglyceridemia: historical perspective and review of biological mechanisms. *Am J Clin Nutr* **71**, 412–433.
43. Merchant AT, Anand SS, Kelemen LE, *et al.* (2007) Carbohydrate intake and HDL in a multiethnic population. *Am J Clin Nutr* **85**, 225–230.
44. Sharman MJ, Gomez AL, Kraemer WJ, *et al.* (2004) Very low-carbohydrate and low-fat diets affect fasting lipids and postprandial lipemia differently in overweight men. *J Nutr* **134**, 880–885.
45. Maki KC (2004) Dietary factors in the prevention of diabetes mellitus and coronary artery disease associated with the metabolic syndrome. *Am J Cardiol* **93**, 12C–17C.
46. Hu T, Mills KT, Yao L, *et al.* (2012) Effects of low-carbohydrate diets versus low-fat diets on metabolic risk factors: a meta-analysis of randomized controlled clinical trials. *Am J Epidemiol* **176**, Suppl. 7, S44–S54.
47. Grundy SM, Abate N & Chandalia M (2002) Diet composition and the metabolic syndrome: what is the optimal fat intake? *Am J Med* **113**, Suppl. 9B, 25S–29S.
48. Grundy SM (1998) Hypertriglyceridemia, atherogenic dyslipidemia, and the metabolic syndrome. *Am J Cardiol* **81**, 18B–25B.
49. Samaha FF (2005) Effect of very high-fat diets on body weight, lipoproteins, and glycemic status in the obese. *Curr Atheroscler Rep* **7**, 412–420.

50. Hooper L, Summerbell CD & Thompson R, *et al.* (2012) Reduced or modified dietary fat for preventing cardiovascular disease *The Cochrane Database of Systematic Reviews*, issue 5, CD002137.
51. Sniderman AD, De Graaf J, Couture P, *et al.* (2009) Regulation of plasma LDL: the apoB paradigm. *Clin Sci (Lond)* **118**, 333–339.
52. Zhang L, Qiao Q, Tuomilehto J, *et al.* (2010) Distinct ethnic differences in lipid profiles across glucose categories. *J Clin Endocrinol Metab* **95**, 1793–1801.
53. Samocha-Bonet D, Campbell LV, Viardot A, *et al.* (2010) A family history of type 2 diabetes increases risk factors associated with overfeeding. *Diabetologia* **53**, 1700–1708.
54. Samocha-Bonet D, Campbell LV, Mori TA, *et al.* (2012) Overfeeding reduces insulin sensitivity and increases oxidative stress, without altering markers of mitochondrial content and function in humans. *PLOS ONE* **7**, e36320.
55. Bachmann OP, Dahl DB, Brechtel K, *et al.* (2001) Effects of intravenous and dietary lipid challenge on intramyocellular lipid content and the relation with insulin sensitivity in humans. *Diabetes* **50**, 2579–2584.
56. Magkos F, Fabbrini E, Conte C, *et al.* (2012) Relationship between adipose tissue lipolytic activity and skeletal muscle insulin resistance in nondiabetic women. *J Clin Endocrinol Metab* **97**, E1219–E1223.
57. Lewis GF, Carpentier A, Adeli K, *et al.* (2002) Disordered fat storage and mobilization in the pathogenesis of insulin resistance and type 2 diabetes. *Endocr Rev* **23**, 201–229.
58. Fabbrini E, Magkos F, Mohammed BS, *et al.* (2009) Intrahepatic fat, not visceral fat, is linked with metabolic complications of obesity. *Proc Natl Acad Sci U S A* **106**, 15430–15435.
59. Misra A & Vikram NK (2003) Clinical and pathophysiological consequences of abdominal adiposity and abdominal adipose tissue depots. *Nutrition* **19**, 457–466.
60. Goel K, Misra A, Vikram NK, *et al.* (2010) Subcutaneous abdominal adipose tissue is associated with the metabolic syndrome in Asian Indians independent of intra-abdominal and total body fat. *Heart* **96**, 579–583.