

Comments on Kanis et al.

Citation for published version (APA):

Geusens, P., & van den Bergh, J. (2019). Comments on Kanis et al. Characteristics of recurrent fractures. *Osteoporosis International*, 30(2), 529-530. <https://doi.org/10.1007/s00198-018-4735-y>

Document status and date:

Published: 01/02/2019

DOI:

[10.1007/s00198-018-4735-y](https://doi.org/10.1007/s00198-018-4735-y)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



Comments on Kanis et al.: Characteristics of recurrent fractures

P. Geusens^{1,2} · J. van den Bergh^{3,4}

Received: 16 September 2018 / Accepted: 8 October 2018 / Published online: 18 December 2018
© International Osteoporosis Foundation and National Osteoporosis Foundation 2018

Dear Editor

We have read with great interest the paper “Characteristics of recurrent fractures” of Kanis et al. [1] and appreciate their conclusion that the high imminent fracture risk within 1 year provides the rationale for very early intervention immediately after a sentinel fracture, in line with previous recommendations [2–4]. These data, presumably of untreated subjects, are therefore of extreme relevance for the Fracture Liaison Service (FLS).

However, it was difficult to deduce the number of subsequent fractures per patient and the absolute 1-year subsequent fracture incidence.

1. From the abstract, we understand that over 10 years, subsequent fractures were sustained in 28% of 1498 individuals with a sentinel hip fracture (according to Table 5, this number refers to women only). From this, we could deduce that 419 women had 984 subsequent fractures, which suggests a mean of 2.3 subsequent fractures per woman.
2. From the text, we understand that subsequent fractures were sustained in 35% of 2074 individuals with a sentinel hip fracture (women and men combined). This indicates that 726 subjects had 1221 fractures, a mean of 1.7 subsequent fractures per subject after a hip fracture.
3. Furthermore, it is stated that in individuals with a subsequent fracture following a sentinel hip fracture ($n = 726$), 45% sustained the first subsequent fracture within 1 year of the sentinel fracture. From this, we could calculate that 326 patients had a subsequent fracture after a recent hip fracture, which indicates an absolute imminent 1-year subsequent fracture risk of 15.7% after a sentinel hip fracture. We could only calculate this 1-year absolute fracture incidence after a sentinel hip fracture.

It would therefore be of great interest for clinicians who run or want to start an FLS if the authors could provide the results for absolute 1- and 2-year subsequent fracture incidence in women and men according to each sentinel fracture.

Sincerely,
Piet Geusens
Joop van den Bergh

Compliance with ethical standards

Conflicts of interest None.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

✉ P. Geusens
piet.geusens@scarlet.be

- ¹ Department of Internal Medicine, Rheumatology, Maastricht University Medical Centre+ (MUMC+), PO Box 616, 6200 MD Maastricht, the Netherlands
- ² University Hasselt, Agoralaan gebouw A, B-3590 Diepenbeek, Belgium
- ³ NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre+ (MUMC+), PO Box 616, 6200, MD Maastricht, the Netherlands
- ⁴ Department of Internal Medicine, VieCuri Medical Centre, PO Box 1926, 5900 BX Venlo, the Netherlands

References

1. Kanis JA, Johansson H, Odén A, Harvey NC, Gudnason V, Sanders KM, Sigurdsson G, Siggeirsdottir K, Fitzpatrick LA, Borgström F, McCloskey EV (2018) Characteristics of recurrent fractures. *Osteoporos Int* 29(8):1747–1757
2. Akesson K, Marsh D, Mitchell PJ, McLellan AR, Stenmark J, Pierroz DD, Kyer C, Cooper C, Fracture Working Group IOF (2013) Capture the Fracture: a Best Practice Framework and global campaign to break the fragility fracture cycle. *Osteoporos Int* 24(8): 2135–2152
3. Eisman JA, Bogoch ER, Dell R, Harrington JT, McKinney RE Jr, McLellan A, Mitchell PJ, Silverman S, Singleton R, Siris E, ASBMR Task Force on Secondary Fracture Prevention (2012) Making the

- first fracture the last fracture: ASBMR task force report on secondary fracture prevention. *J Bone Miner Res* 27(10):2039–2046
4. Lems WF, Dreinhöfer KE, Bischoff-Ferrari H, Blauth M, Czerwinski E, da Silva J, Herrera A, Hoffmeyer P, Kvien T, Maalouf G, Marsh D, Puget J, Puhl W, Poor G, Rasch L, Roux C, Schüler S, Serio B, Tarantino U, van Geel T, Woolf A, Wyers C, Geusens P (2017 May) EULAR/EFORT recommendations for management of patients older than 50 years with a fragility fracture and prevention of subsequent fractures. *Ann Rheum Dis* 76(5):802–810