

From fear of movement-related pain and avoidance to chronic pain disability

Citation for published version (APA):

Meulders, A. (2019). From fear of movement-related pain and avoidance to chronic pain disability: a state-of-the-art review. *Current Opinion in Behavioral Sciences*, 26, 130-136.
<https://doi.org/10.1016/j.cobeha.2018.12.007>

Document status and date:

Published: 01/04/2019

DOI:

[10.1016/j.cobeha.2018.12.007](https://doi.org/10.1016/j.cobeha.2018.12.007)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



From fear of movement-related pain and avoidance to chronic pain disability: a state-of-the-art review[☆]

Ann Meulders^{1,2}

This review addresses recent developments in fear of movement-related pain and avoidance research, adopting a contemporary learning approach focusing on cognitive, emotional, and behavioral factors, and emphasizing the role of the motivational context including pain-avoidance goals but also other valued life goals. We particularly demonstrated the role of classical conditioning in the acquisition, generalization, and extinction of fear of movement-related pain, and how safety-seeking behavior during exposure-based treatment may hamper genuine extinction causing relapse. We further highlighted the importance of newly developed methods to study operant conditioning of behavioral avoidance itself, and its relationship with individual vulnerability factors (e.g. catastrophizing) and resiliencies (e.g. positive affect) to further our understanding of mechanisms underlying the development and persistence of chronic pain disability.

Addresses

¹Research Group Health Psychology, KU Leuven, Leuven, Belgium

²Experimental Health Psychology, Department of Clinical Psychological Science, Maastricht University, The Netherlands

Corresponding author: Meulders, Ann (ann.meulders@kuleuven.be)

Current Opinion in Behavioral Sciences 2019, 26:130–136

This review comes from a themed issue on **Pain and aversive motivation**

Edited by **Ben Seymour** and **Joshua Johansen**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 31st January 2019

<https://doi.org/10.1016/j.cobeha.2018.12.007>

2352-1546/© 2018 Elsevier Ltd. All rights reserved.

Introduction

Pain is a biologically salient signal and adaptive response to bodily threat that prompts protective action. The protective response repertoire initially includes increased arousal, selective attention, escape, and painful facial expressions, as well as long-term recuperative avoidance and safety-seeking behaviors. Pain is thus a *motivational state* initiating early defensive responses followed by recuperative behaviors aimed to promote recovery. Given its intrinsically

alarming function, pain is a potent motivator for learning—it stimulates the detection of stimuli predicting the occurrence of pain and bodily threat in order to avoid or minimize their impact. Pavlovian or classical conditioning is the prime mechanism underlying the prediction of potentially harmful stimuli in the environment [1,2]. When pairing an initially neutral stimulus (conditioned stimulus; CS) with an aversive outcome (unconditioned stimulus; US), this stimulus in itself may start to elicit defensive responses (conditioned response; CR).

Contemporary fear-avoidance models of chronic musculoskeletal pain advance pain-related fear and avoidance as key contributors to the transition from acute to chronic pain [3^{**},4]. Catastrophic misinterpretations of pain as being harmful may give rise to pain-related fear, which in turn can initiate avoidance behavior intended to avert bodily threat. When protective action serves to reduce genuine bodily threat, it is highly adaptive, but when pain persists beyond healing time and turns chronic, it may paradoxically increase suffering and disability (Figure 1). In this paper, we will review the role of learning mechanisms in the development, spreading, and persistence of fear of movement-related pain and avoidance, and how they contribute to pain-related disability.


The role of learning in pain-related fear and avoidance

The acquisition of pain-related fear

Because of its biological significance, pain can be considered a US, which by itself triggers protective responses such as psychophysiological arousal (e.g. elevated startle responses, skin conductance, and muscle tone) and escape behavior. Diverse stimuli can precede pain and thus serve as CSs including exteroceptive (tactile, visual, auditory), interoceptive (visceral, olfactory), and proprioceptive stimuli (movement, posture). For patients with musculoskeletal pain, proprioceptive information is particularly relevant to predict pain episodes. For example, a neutral movement (e.g. bending to pick something up) preceding a pain episode (e.g. shooting back pain) may come to signal bodily harm and elicit *fear of movement-related pain* and spur avoidance behavior (Figure 2). So far, we have described situations in which discrete cues predict pain-onset, typically inducing *cued pain-related fear*. Some patients, however, report spontaneous pain


[☆] Ann Meulders is a postdoctoral researcher of the Research Foundation Flanders (FWO-Vlaanderen), Belgium (grant ID 12E3717N) and is also supported by a Vidi grant from the Netherlands Organisation for Scientific Research (NWO), The Netherlands (grant ID 452-17-002).

Figure 1



Fear-avoidance model of pain.

Figure 2



Fear of movement-related pain acquisition and generalization.

Note – The red radiating dot represents a shooting back pain (i.e. pain-US). During acquisition bending forward (CS+) is the movement that is followed by the pain-US, whereas leaning backward (CS-) is the movement that is not followed by the pain-US. The generalization stimuli (GSs) are novel intermediate movements between the CS+ and the CS-.

flare-ups without cues predicting the (increases in) pain [5]. In situations of apparent unpredictable pain, long-lasting contextual cues may be considered predictors of pain, leading to *contextual pain-related fear*. Meulders *et al.* [6] argued that translated to chronic pain conditions, cued pain-related fear (i.e. predictable pain), and chronic contextual pain-related fear (i.e. unpredictable pain), respectively, model regional pain syndromes (e.g. low back pain) and widespread pain such as fibromyalgia.

Cumulating experimental evidence supports the role of associative learning processes in the acquisition of both cued and contextual pain-related fear. Meulders *et al.* [7] developed a voluntary joystick movement paradigm to model pain-related fear in musculoskeletal pain disorders: in the predictable condition, neutral joystick movements (moving to left/right) served as CSs of which one direction predicted the painful electrocutaneous stimulus (pain-US); in the unpredictable condition, two other joystick movements (moving upward/downward) were never followed by pain. Results showed that the CS+ movement elicited elevated startle amplitudes, increased self-reported fear, and larger response latencies (as a proxy of behavioral avoidance) than the CS-movements. No such differences occurred for both unpredictable CSs, but fear accrued to the context (i.e. increased startle responses during the intertrial interval). Furthermore, we demonstrated similar conditioned responses to the mere intention to perform pain-associated movements, before actually performing these movements [8], suggesting that *imagining a painful movement* can activate the memory representation of the movement-pain association and in turn evoke conditioned fear [9]. Other evidence supports the idea that pain-related fear can be acquired without direct experience, for example through observation and instruction [10–12], by symbolically representing pain [13] or by virtue of derived relationships with pain such as conceptual equivalence between stimuli [14,15].

The generalization of pain-related fear

Learning which stimuli predict bodily threat is highly adaptive. Nevertheless, adaptive learners face the challenge of how to deal with variations in the appearances of signaling stimuli. Rapid adaptation to a dynamic environment requires balancing discrimination and generalization towards novel situations. *Stimulus generalization* [16] allows extrapolation of the predictive value of one stimulus to novel stimuli (generalization stimuli, GSs) based on (non-)perceptual similarities minimizing the necessity to learn everything anew. Unbridled fear generalization to safe situations may lead to sustained anxiety and excessive avoidance behavior initiating a pathway toward disability [17].

Using the voluntary joystick movement paradigm, we demonstrated that healthy participants were more afraid of and showed elevated startle responses to

movements that resembled the original CS+ compared to those resembling the CS– [6,18]; in the unpredictable condition, however, fear generalized in a non-differential way to all novel movements. In contrast with healthy controls, fibromyalgia patients showed nondifferential generalization in both conditions, suggesting that all GSs triggered strong conditioned responses, irrespective of the previous learning history [19,17]. Patients showed excessive fear-to-safe stimuli (GSs resembling the CS–) rather than to those resembling the threatening CS+, suggesting that *impaired safety learning* drives this ‘overgeneralization’ effect. In addition, after receiving disconfirming information during an extinction of generalization phase (i.e. GSs without US), healthy participants quickly update their beliefs leading to extinction of generalized fear whereas fibromyalgia patients continued to display elevated pain-related fear and expectancy to the unreinforced GSs [20].

Similar ‘overgeneralization’ effects exist in expectancy judgments of chronic unilateral hand pain patients in a contingency learning task [21]. Other evidence supporting the lack of selective learning in fibromyalgia patients stems from a clinical diary scenario study using a *blocking* design in which diary entries of a fictitious patient served as CSs and verbal outcomes ‘pain’/‘no pain’ as US [22]. First, one cue (Kim was stressed; A+) is paired with ‘pain’. Next, another cue (Kim slept badly; X), referred to as the *blocked stimulus*, presented in compound with the original pain-associated cue is also paired with ‘pain’ (AX+). During the crucial test of X alone, conditioned responding is typically weak or *blocked*, despite previous pairings with the US, because A suffices to predict the occurrence of the US. Compared to healthy controls, fibromyalgia patients did not show blocking to X.

Humans also can abstract conceptual details during a learning episode, which allows them to generalize fear to physically dissimilar stimuli that are semantically related, a process known as *category-based fear conditioning* [23]. Using an adapted version of the voluntary joystick movement paradigm, we paired exemplars from one action category with pain (e.g. ‘opening’ a small, rectangular, blue box with a left joystick movement) and observed fear generalization to conceptually similar exemplars (e.g. ‘opening’ a large, round, yellow box with a right joystick movement), but not to novel exemplars of another action category (e.g. ‘closing’ a small, round, green box with a left joystick movement) [24]. Bennett *et al.* [14] provided further support for non-perceptual generalization by showing that pain-related fear transferred from nonsense word CSs to joystick movements that were never paired with pain themselves but were trained to belong to the same stimulus equivalence category as the CSs.

Extinction and return of pain-related fear

When the CS is no longer followed by the US (or its mental representation), the CS loses its predictive value

and will no longer evoke conditioned responses. *Extinction* is a fragile and context-dependent process [25]. It is commonly accepted that extinction does not erase the original CS-US association, but that a new extinction memory is formed that inhibits the behavioral expression of the first-learned association. Intact acquisition memory is evidenced by the return of fear when the CS is encountered in a different context than the extinction context (i.e. renewal). Also unexpected encounters with the original pain-US (reinstatement) [26] such as pain flare-ups or other unanticipated non-pain stressors (cross-US reinstatement) [27] can restore pain-related fear. In addition, another challenge is the generalization of the extinction to novel but similar movements and other contexts. Research in back pain patients showed that exposure to physical movement did not generalize to novel dissimilar movements, suggesting that ‘under-generalization’ of extinction may contribute as well to the maintenance of chronic pain disability [28].

Exposure *in vivo* is the clinical analog of Pavlovian extinction and has been effectively used in patients with chronic musculoskeletal pain [29] such as low back pain [30,31] and CRPS [32]. In essence, this treatment entails confrontation with feared movement/activity, following an idiosyncratic fear hierarchy, until fear is reduced and daily functioning can be resumed. Notwithstanding the success of exposure to reduce pain-related fear and disability, relapse often occurs. The use of *safety-seeking behavior* during exposure (e.g. bending to pick something up with a straight back) has been theorized to impede fear extinction [33]. The idea is that when safety-seeking behavior is performed during extinction, the non-occurrence of the US can be attributed to this safety-seeking behavior hampering genuine extinction learning. However, there is a current debate as to whether allowing judicious safety behavior facilitates or jeopardizes treatment. According to a recent meta-analysis this question remains unsettled [34**] highlighting the need for systematic replications with sufficient statistical power and large sample sizes, using idiosyncratic safety-seeking behavior (not predetermined by the experimenter) and taking the motivational context into account. Current recommendations regarding the use of safety-seeking behavior in clinical practice should therefore be interpreted with caution.

We investigated whether performing an avoidance response would hamper extinction of fear of movement-related pain using a voluntary joystick movement paradigm [35]. During extinction, the safety group was instructed that they could avoid the US by pressing a ‘safety button’ on the joystick during the pain-associated movement, whereas a control group did not have this option. When the avoidance option was eliminated during the subsequent test, return of fear of movement-related pain was observed in the safety group, but not in the

control group, demonstrating *protection from extinction*. A follow-up study showed that the same behavior only reliably led to return of fear when the response was serving a pain-avoidance goal (avoiding the US), not when it was serving an achievement goal (attaining a reward) [36], underscoring the importance of the motivational context in understanding the role of safety-seeking behavior in exposure-based interventions.

The relationship between fear and avoidance in a motivational context

Avoidance is defined as overt behavior aimed to prevent the occurrence of a threatening event (e.g. pain), and the relief experienced when the anticipated catastrophe does not occur has been advanced as a reinforcer of this behavior [37]. Experimental research mainly focused on self-report and psychophysiological correlates of pain-related fear, leaving its behavioral component poorly understood. The lack of scientific attention for avoidance behavior itself is based on the tacit assumption that fear is the sole motivator of avoidance [38] implying that extinguishing fear (i.e. exposure to the feared stimulus without avoidance option) would suffice to abolish avoidance behavior, a procedure referred to as *response prevention with extinction* or *flooding* [39]. However, increasing evidence suggests that there is no one-to-one relationship between fear and avoidance, insinuating that more complex decision-making may be at play, involving weighing the value of pain avoidance against other life goals (e.g. going to work) [40]. Indeed pain-related fear typically occurs in a context of multiple, competing goals. Recent experimental research demonstrated that avoidance behavior, but not fear can be attenuated when another valued, competing goal (i.e. attaining a monetary reward) is introduced [41]. Pain-related fear induces discomfort but does not necessarily lead to disability; therefore, the role of behavioral avoidance is essential in explaining the pathway to disability, highlighting the need for more research on this topic.

Although avoidance behavior aims to reduce pain-related fear in the short run, it may paradoxically boost pain-related in the long run, possibly by increasing the *threat value of pain*. People can indeed use their own avoidance behavior to infer threat, that is ex-consequencia reasoning [42]: “*I’m avoiding, so it must be painful and/or I must be afraid*”. In a recent study [43] we provided support for the bidirectional relationship between fear and avoidance. Using a heat pain paradigm, participants in the avoidance group were led to believe that they could avoid full exposure to a painful heat stimulus by pressing the stop button, whereas control group participants believed they were always exposed to the full painful heat stimulus. In reality, the stimulation was identical in both groups. During the test without avoidance option, participants in the avoidance group reported higher levels of pain-related fear. Interestingly, receiving instructions that

avoidance was possible already increased fear, even before avoidance behavior was enacted.

Operant learning of pain-related avoidance behavior

Another mechanism that might contribute to chronic pain disability is *instrumental or operant conditioning* [44,2], that is, learning about one's own behavior and its consequences. Successful avoidance of feared movements (e.g. bending to pick something up with a straight back) can be negatively reinforced because the feared catastrophe (e.g. spine snapping) does not occur. Despite the importance of avoidance behavior, a lack of ecologically valid paradigms in humans hampers the progress in this field. Available paradigms have several limitations including: (1) *instructed avoidance* precluding the opportunity to study the acquisition of avoidance behavior itself; (2) using dichotomous responses (press/not press a button) that may lack sensitivity and accuracy; (3) *adaptive avoidance* is examined: the response effectively cancels the US; (4) using low-cost avoidance (button press with no response cost). Arguably, this does not reflect the clinical reality well because especially high-cost avoidance may contribute to disability in the long run. If there is no response cost to avoidance behavior, the underlying motivation remains elusive as well.

Striving for increased ecological validity, we developed a novel operant pain-related avoidance paradigm, in which we modeled a response cost of avoidance behavior using a trade-off between pain probability and effort. In a reaching task, participants moved a robotic arm to a target location choosing one of three possible trajectories. The shortest/easiest trajectory (T1) was always paired with pain. If they deviated from this trajectory, pain could be partly or totally prevented (T2 = 50% reinforcement; T3 = 0% reinforcement), but more effort was needed (T2 = moderate resistance and deviation; T3 = largest resistance and deviation). The yoked control group received the same reinforcement schedule irrespective of their own choice of trajectory. Avoidance behavior was operationalized as the deviation from the shortest/easiest trajectory and results confirmed larger deviations in the experimental group than in the yoked group [45^{••}]. Overcoming some limitations of existing methods, this paradigm may hold promise to advance our knowledge about how pain-related avoidance behavior is acquired. It also allows testing for generalization to assess when avoidance behavior becomes maladaptive or excessive.

Individual differences in fear learning: vulnerability and resilience

The role of vulnerability factors such as pain-related fear, catastrophizing, and avoidance have been well-documented in the development and maintenance of chronic pain disability. Far less attention has been dedicated to *resilience factors* such as optimism and positive affect and how these can modulate fear learning and pain-related outcomes [46,47[•],48].

Evidence suggests that positive affect may alter fear learning. First, positive affect may counteract excessive *fear generalization*. Using the voluntary joystick movement paradigm, we showed that experimentally increasing positive affect protected against generalization of fear-to-safe stimuli [49]. Interestingly, positive affect preserves adaptive fear generalization, but eliminates maladaptive 'overgeneralization'. Second, positive affect might enhance inhibitory learning [50], the key mechanism underlying exposure treatment. For example, we have shown that trait positive affect protects against deficient safety learning during extinction in individuals with high levels of subclinical anxiety [51]. Positive affect also has been shown to prevent return of fear after successful extinction [52,53]. Increasing positive affect in chronic pain patients thus may have beneficial effects through limiting excessive spreading of pain-related fear and avoidance as well as through the optimization of (inhibitory) learning processes during treatment [47[•]].

Conclusions

We reviewed recent developments in fear of movement-related pain and avoidance research, adopting a contemporary learning approach focusing on cognitive, emotional, and behavioral factors, and emphasizing the need for a motivational perspective to further our understanding of the development and persistence of chronic pain disability. Although fear generalization per se is adaptive, we have argued that *excessive generalization* of fear and avoidance may be a pathway to disability. Future research should focus on the bidirectional relationship between fear and avoidance, employ and develop novel ecologically valid methods to study avoidance behavior itself, and the relationship with individual vulnerability factors and resiliencies. Research efforts exploring possible avenues tackling generalization such as positive affect interventions, discrimination training, and neural stimulation techniques may strengthen our treatment repertoire for sufferers from chronic pain.

Conflict of interest statement

Nothing declared.

Acknowledgements

The author is grateful for the input of many colleagues over the years, and she particularly thanks Johan Vlaeyen for ongoing discussions on the topic. She also thanks Rini Masuy and Kristof Vandael for their help with the artwork.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Pavlov IP: *Conditioned Reflexes*. London: Oxford University Press; 1927.
2. Vlaeyen JW: **Learning to predict and control harmful events: chronic pain and conditioning**. *Pain* 2015, **156**(Suppl. 1):S86-S93.

3. Vlaeyen JW, Crombez G, Linton SJ: **The fear-avoidance model of pain.** *Pain* 2016, **157**:1588-1589.
This paper provides an update on the fear-avoidance model of pain, reviewing the acquisition, generalization, and extinction of pain-related fear and highlighting the role of the motivational context such as goal pursuit in chronic pain disability.
4. Vlaeyen JW, Linton SJ: **Fear-avoidance model of chronic musculoskeletal pain: 12 years on.** *Pain* 2012, **153**:1144-1147.
5. Johnson LM, Zautra AJ, Davis MC: **The role of illness uncertainty on coping with fibromyalgia symptoms.** *Health Psychol* 2006, **25**:696-703.
6. Meulders A, Vandebroek N, Vervliet B, Vlaeyen JW: **Generalization gradients in cued and contextual pain-related fear: an experimental study in healthy participants.** *Front Hum Neurosci* 2013, **7**:345.
7. Meulders A, Vansteenwegen D, Vlaeyen JW: **The acquisition of fear of movement-related pain and associative learning: a novel pain-relevant human fear conditioning paradigm.** *Pain* 2011, **152**:2460-2469.
8. Meulders A, Vlaeyen JWS: **Mere intention to perform painful movements elicits fear of movement-related pain: an experimental study on fear acquisition beyond actual movements.** *J Pain* 2013, **14**:412-423.
9. Meulders A, Harvie DS, Lorimer Moseley G, Vlaeyen JWS: **Generalization of pain-related fear using a left-right hand judgment conditioning task.** *Behav Ther* 2015, **46**:699-716.
10. Goubert L, Vlaeyen JW, Crombez G, Craig KD: **Learning about pain from others: an observational learning account.** *J Pain* 2011, **12**:167-174.
11. Helsen K, Vlaeyen JWS, Goubert L: **Indirect acquisition of pain-related fear: an experimental study of observational learning using coloured cold metal bars.** *PLoS One* 2015, **10**:e0117236.
12. Koban L, Jepma M, Geuter S, Wager TD: **What's in a word? How instructions, suggestions, and social information change pain and emotion.** *Neurosci Biobehav Rev* 2017, **81**:29-42.
13. Jepma M, Wager TD: **Conceptual conditioning: mechanisms mediating conditioning effects on pain.** *Psychol Sci* 2015, **26**:1728-1739.
14. Bennett MP, Meulders A, Baeyens F, Vlaeyen JW: **Words putting pain in motion: the generalization of pain-related fear within an artificial stimulus category.** *Front Psychol* 2015, **6**:520.
15. Meulders A, Bennett MP: **The concept of contexts in pain: generalization of contextual pain-related fear within a de novo category of unique contexts.** *J Pain* 2018, **19**:76-87.
16. Dymond S, Dunsmoor JE, Vervliet B, Roche B, Hermans D: **Fear generalization in humans: systematic review and implications for anxiety disorder research.** *Behav Ther* 2015, **46**:561-582.
17. Meulders A, Jans A, Vlaeyen JW: **Differences in pain-related fear acquisition and generalization: an experimental study comparing patients with fibromyalgia and healthy controls.** *Pain* 2015, **156**:108-122.
18. Meulders A, Vlaeyen JWS: **The acquisition and generalization of cued and contextual pain-related fear: an experimental study using a voluntary movement paradigm.** *Pain* 2013, **154**:272-282.
19. Harvie DS, Moseley GL, Hillier SL, Meulders A: **Classical conditioning differences associated with chronic pain: a systematic review.** *J Pain* 2017, **18**:889-898.
This paper reviews conditioning differences between healthy controls and chronic pain patients, suggesting altered differential and generalization learning in patients.
20. Meulders A, Meulders M, Stouten I, De Bie J, Vlaeyen JWS: **Extinction of fear generalization: a comparison between fibromyalgia patients and healthy control participants.** *J Pain* 2017, **18**:79-95.
21. Meulders A, Harvie DS, Bowering JK, Caragianis S, Vlaeyen JWS, Moseley GL: **Contingency learning deficits and generalization in chronic unilateral hand pain patients.** *J Pain* 2014, **15**:1046-1056.
22. Meulders A, Boddez Y, Blanco F, Van Den Houte M, Vlaeyen JWS: **Reduced selective learning in patients with fibromyalgia vs healthy controls.** *Pain* 2018, **159**:1268-1276.
This paper introduces a simple contingency learning task without experimental pain induction that is sensitive to pick up selective learning differences between chronic pain patients and healthy controls.
23. Dunsmoor JE, Murphy GL: **Categories, concepts, and conditioning: how humans generalize fear.** *Trends Cogn Sci* 2015, **19**:73-77.
24. Meulders A, Vandael K, Vlaeyen JWS: **Generalization of pain-related fear based on conceptual knowledge.** *Behav Ther* 2017, **48**:295-310.
25. Vervliet B, Craske M, Hermans D: **Fear extinction and relapse: state of the art.** *Ann Rev Clin Psychol* 2013, **9**:215-248.
26. den Hollander M, Meulders A, Jakobs M, Vlaeyen JW: **The effect of threat information on acquisition, extinction, and reinstatement of experimentally conditioned fear of movement-related pain.** *Pain Med* 2015, **16**:2302-2315.
27. Meulders A, Rousseau A, Vlaeyen J: **Motor intention as a trigger for fear of movement-related pain: an experimental cross-US reinstatement study.** *J Exp Psychopathol* 2015, **6**(3):206-228.
28. Goubert L, Francken G, Crombez G, Vansteenwegen D, Lysens R: **Exposure to physical movement in chronic back pain patients: no evidence for generalization across different movements.** *Behav Res Ther* 2002, **40**:415-429.
29. Vlaeyen JWS, Morley S, Linton SJ, Boersma K, De Jong JR: *Pain-Related Fear: Exposure-based Treatment for Chronic Pain.* Seattle: IASP Press; 2012.
30. Glombiewski JA, Holzapfel S, Riecke J, Vlaeyen JWS, de Jong J, Lemmer G, Rief W: **Exposure and CBT for chronic back pain: an RCT on differential efficacy and optimal length of treatment.** *J Consult Clin Psychol* 2018, **86**:533-545.
31. Goossens MEJB, de Kinderen RJA, Leeuw M, de Jong JR, Ruijgrok J, Evers SMAA, Vlaeyen JWS: **Is exposure in vivo cost-effective for chronic low back pain? A trial-based economic evaluation.** *BMC Health Serv Res* 2015, **15**:549.
32. den Hollander M, Goossens M, de Jong J, Ruijgrok J, Oosterhof J, Onghena P, Smeets R, Vlaeyen JW: **Expose or protect? A randomized controlled trial of exposure in vivo vs pain-contingent treatment as usual in patients with complex regional pain syndrome type 1.** *Pain* 2016, **157**:2318-2329.
33. Lovibond PF, Mitchell CJ, Minard E, Brady A, Menzies RG: **Safety behaviours preserve threat beliefs: protection from extinction of human fear conditioning by an avoidance response.** *Behav Res Ther* 2009, **47**:716-720.
34. Meulders A, Van Daele T, Volders S, Vlaeyen JW: **The use of safety-seeking behavior in exposure-based treatments for fear and anxiety: benefit or burden? A meta-analytic review.** *Clin Psychol Rev* 2016, **45**:144-156.
This paper reviews and provides a meta-analysis of the effect of safety-seeking behavior during exposure *in vivo*, concluding that the jury is still out, and thus current clinical recommendations are unwarranted.
35. Volders S, Meulders A, De Peuter S, Vervliet B, Vlaeyen JW: **Safety behavior can hamper the extinction of fear of movement-related pain: an experimental investigation in healthy participants.** *Behav Res Ther* 2012, **50**:735-746.
36. Volders S, Meulders A, De Peuter S, Vlaeyen JWS: **The reduction of fear of movement-related pain: does motivational context matter?** *Clin J Pain* 2015, **31**(11):933-945.
37. Volders S, Boddez Y, De Peuter S, Meulders A, Vlaeyen JWS: **Avoidance behavior in chronic pain research: a cold case revisited.** *Behav Res Ther* 2015, **64**:31-37.
38. Krypotos A-M, Effting M, Kindt M, Beckers T: **Avoidance learning: a review of theoretical models and recent developments.** *Front Behav Neurosci* 2015, **9**.
39. Mineka S: **The role of fear in theories of avoidance learning, flooding, and extinction.** *Psychol Bull* 1979, **86**:985.
40. Wiech K, Tracey I: **Pain, decisions, and actions: a motivational perspective.** *Front Neurosci* 2013, **7**:46.

41. Claes N, Karos K, Meulders A, Crombez G, Vlaeyen JW: **Competing goals attenuate avoidance behavior in the context of pain.** *J Pain* 2014, **15**:1120-1129.
42. Arntz A, Rauner M, van den Hout M: **"If I feel anxious, there must be danger": ex-consequentia reasoning in inferring danger in anxiety disorders.** *Behav Res Ther* 1995, **33**:917-925.
43. van Vliet CM, Meulders A, Vancleef LMG, Vlaeyen JWS: **The opportunity to avoid pain may paradoxically increase fear.** *J Pain* 2018, **19**(October (10)):1222-1230 <http://dx.doi.org/10.1016/j.jpain.2018.05.003>.
44. Skinner BF: *Science and Human Behavior*. New York: McMillan; 1953.
45. Meulders A, Franssen M, Fonteyne R, Vlaeyen JWS: **Acquisition and extinction of operant pain-related avoidance behavior using a 3 degrees-of-freedom robotic arm.** *Pain* 2016, **157**.
- This paper introduces a novel operant pain-related avoidance conditioning allowing to study not only the acquisition of avoidance, but also its extinction and generalization.
46. Finan PH, Garland EL: **The role of positive affect in pain and its treatment.** *Clin J Pain* 2015, **31**:177-187.
47. Hanssen MM, Peters ML, Boselie JJ, Meulders A: **Can positive affect attenuate (persistent) pain? State of the art and clinical implications.** *Curr Rheumatol Reports* 2017, **19**:80.

This paper reviews the potential of positive affect/optimism interventions in treatment of chronic pain, emphasizing the need to focus not only on resilience, but also on vulnerability.

48. Thong ISK, Tan G, Jensen MP: **The buffering role of positive affect on the association between pain intensity and pain related outcomes.** *Scand J Pain* 2017, **14**:91-97.
49. Geschwind N, Meulders M, Peters ML, Vlaeyen JW, Meulders A: **Can experimentally induced positive affect attenuate generalization of fear of movement-related pain?** *J Pain* 2015, **16**:258-269.
50. Zbozinek TD, Craske MG: **The role of positive affect in enhancing extinction learning and exposure therapy for anxiety disorders.** *J Exp Psychopathol* 2017, **8**:13-39.
51. Meulders A, Meulders M, Vlaeyen JWS: **Positive affect protects against deficient safety learning during extinction of fear of movement-related pain in healthy individuals scoring relatively high on trait anxiety.** *J Pain* 2014, **15**:632-644.
52. Zbozinek TD, Craske MG: **Positive affect predicts less reacquisition of fear: relevance for long-term outcomes of exposure therapy.** *Cogn Emotion* 2017, **31**:712-725.
53. Zbozinek TD, Holmes EA, Craske MG: **The effect of positive mood induction on reducing reinstatement fear: relevance for long term outcomes of exposure therapy.** *Behav Res Ther* 2015, **71**:65-75.