

Cardiorespiratory Fitness and Long-Term Mortality

Citation for published version (APA):

Stamatakis, E., Koster, A., & Mork, P. J. (2018). Cardiorespiratory Fitness and Long-Term Mortality: Targeting the Least Fit With Incidental Physical Activity. *Journal of the American College of Cardiology*, 72(9), 996-998. <https://doi.org/10.1016/j.jacc.2018.07.001>

Document status and date:

Published: 28/08/2018

DOI:

[10.1016/j.jacc.2018.07.001](https://doi.org/10.1016/j.jacc.2018.07.001)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

EDITORIAL COMMENT

Cardiorespiratory Fitness and Long-Term Mortality



Targeting the Least Fit With Incidental Physical Activity*

Emmanuel Stamatakis, PhD,^{a,b} Annemarie Koster, PhD,^c Paul Jarle Mork, PhD^d

Cardiorespiratory fitness (CRF), which is commonly expressed as relative maximal oxygen uptake in milliliters per minute per kilogram or metabolic equivalents (METs) (1 MET = approximately 3.5 ml/kg/min), is a vital physiological indicator that corresponds to the integrated ability of the cardiac, pulmonary, muscular, and cellular systems to transport oxygen to the mitochondria to perform physical activity (PA). CRF has a genetic component that is similar in magnitude to other major physiological risk factors for cardiovascular disease (CVD) (1), but it is also highly modifiable through PA. Paradoxically, CRF is perhaps the only major physiological risk factor that is not routinely assessed in clinical practice, and despite consistent evidence that shows that even algorithm-estimated CRF improves CVD risk prediction (2), it is not part of any major CVD risk calculation (1). The only known prescription for poor CRF is regular engagement in PA of a certain intensity.

SEE PAGE 987

In this issue of the *Journal*, Clausen et al. (3) examined the association between CRF and mortality in the Copenhagen Male Study (CMS) over an impressive follow-up period of 46 years. CMS

participants were employees from a range of diverse industries, including construction, railway, military, and health care (4). Among the 5,107 middle-aged (ages 40 to 59 years) men who completed the baseline examination in 1970 to 1971, 4,700 (92%) died during follow-up; of those deaths, 2,149 (45.7% of total events) were attributed to CVD mortality. Such an unusually high mortality rate is not surprising considering the long follow-up, the baseline age range of the cohort, and the life expectancy at age 50 years among men in Denmark in 1970 (25 years) (5).

CRF grouping was based on age-standardized z-scores distribution: below lower limit of normal (bottom 5% of z-distribution, mean CRF: 20.7 ± 2.0 ml/kg/min); low normal (between mean and bottom 5%, mean CRF: 28.3 ± 3.1 ml/kg/min); high normal (between mean and top 5%, mean CRF: 37.1 ± 4.1 ml/kg/min); and above the upper limit of normal (top 5% of z-distribution, mean CRF: 49.6 ± 4.8 ml/kg/min). By this classification, the categories of CRF were dependent on the distribution of CRF in this specific study sample rather than existing cut-offs. Compared with the bottom 5% of the CRF z-distribution, the multivariable-adjusted life expectancy gains were 2.1 and 2.9 years for the low and high normal ranges, respectively, and 4.9 years for the top 5% of the CRF distribution. However, when competing mortality risks were taken into account, there were no statistically significant gains specifically for CVD mortality.

Like most epidemiological evidence, this study was based on a single baseline examination of CRF. Reverse causation, which is defined as undiagnosed and/or prodromal and/or subclinical disease that both compromised the exposure (CRF) and predisposed subjects to an earlier death, is a serious issue in epidemiological studies and can lead to deceptive

*Editorials published in the *Journal of the American College of Cardiology* reflect the views of the authors and do not necessarily represent the views of *JACC* or the American College of Cardiology.

From the ^aCharles Perkins Centre, University of Sydney, Sydney, New South Wales, Australia; ^bSydney School of Public Health, Faculty of Medicine and Health, University of Sydney, Sydney, New South Wales, Australia; ^cDepartment of Social Medicine, CAPHRI Care and Public Health Research Institute, Maastricht University, Maastricht, the Netherlands; and the ^dDepartment of Public Health and Nursing, Norwegian University of Science and Technology (NTNU), Trondheim, Norway. The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

conclusions. The long follow-up and the large number of events permitted the investigators to examine possible reverse causation by repeating the all-cause mortality analyses without the 361 participants who died during the first 10 years of follow-up. However, life expectancy gains were only slightly attenuated (1.8, 2.6, and 4.3 years) compared with the analyses in the full sample described above. Although the reassurance offered by this additional analysis was a notable strength of the study, the long follow-up also added some uncertainty to the interpretation of the data, which merits discussion. Drawing potentially causal inferences between states and events that are far apart by nearly 5 decades is difficult, because of the likely plethora of changes that are bound to occur (e.g., changes in participant behavior, life circumstances, and also broader societal changes). One of the most dramatic ecological changes that occurred in Copenhagen during the study period was a major shift toward physically active transportation. Between 1970 and 2016, there was an approximately 2.5-fold surge in cycling for transportation among Copenhagen residents (6). By 2016, 41% of all commuting trips in Copenhagen were done by bicycle, and nearly 20% of all trips were done on foot (6). Considering that uncompromised physical capacity, including adequate CRF, enables PA participation, it is plausible that those participants with a normal or high CRF at baseline in 1970 to 1971 were also more likely to increase incidental PA as the active transportation city infrastructure improved over time. Such a likely reciprocal association between CRF and PA is important to acknowledge in interpreting the current results, and it is unfortunate that no repeated measures of CRF were included in the analysis. The reported life years gained might reflect increased PA that would not only lead to a slower age-related decline in CRF but could also act through a number of cardiometabolic and psychological health pathways. Under such a scenario, the results found by Clausen et al. (3) might over-estimate the independent associations of CRF with future mortality risk. It was also unclear what the role of occupational PA was because only leisure-time PA was considered a confounder, despite previous CMS data that showed interactions between physical demands at work and CRF in terms of all-cause mortality risk (4).

The preceding life expectancy estimates (3) showed that there was relatively little variation (2.1 to 2.9 years) in the middle 90% of the age-adjusted CRF distribution, which corresponded to a wide range of average CRF levels (e.g., approximately 23 to 47 ml/kg/min for men aged 50 years). Essentially, the

results of the current study imply that the substantial longevity gains might be realized simply by moving away from the least-fit end of the CRF distribution. This is in line with previous observations in a predominantly male U.S. sample in which more than half the reduction in all-cause mortality occurred between the low and moderate CRF groups (7). Also, the present data (3) corroborated the well-established substantial gains attained at the lower spectrum of PA, whereby the largest reduction in mortality risk was seen when the groups who did little and/or no PA were compared with those who did some but did not meet the PA guidelines (8).

How does all this translate into a public health and clinical practice message? It is worth noting that the intensity of PA has a strong impact on CRF improvements (1). Higher intensities lead to more rapid and larger CRF gains even when the volume of PA is modest (e.g., high-intensity interval training programs) (9). This does not necessarily mean that those with the lowest CRF should be prescribed high-intensity exercise. In general, when the baseline CRF is lower, less vigorous intensity is needed to produce beneficial improvements in CRF. For example, in adults with a CRF <10 METs (35 ml/kg/min), an intensity of approximately 50% heart rate or theoretical maximum CRF reserve is adequate (1). For most middle-aged people, such an intensity could be walking at a pace that makes the person slightly out of breath when sustained for more than a few minutes. Even slower than brisk walking pace has been associated with substantial all-cause and CVD mortality gains (10): compared with a habitually slow walking pace in upper middle and older age, a medium pace has been shown to be associated with 44% lower CVD mortality risk, with little further gains for a brisk and/or fast pace (10). The 2018 U.S. Physical Activity Guidelines Advisory report (11) acknowledged the importance of light intensity PA for health, but little is currently known about its CRF-promoting potential. Although longitudinal epidemiological and experimental evidence is lacking, it is encouraging that beneficial associations of objectively measured incidental light intensity activity and CRF were reported in a cross-sectional study (12).

In summary, the study by Clausen et al. (3) made excellent use of a historical data resource and contributed important information on the question of whether CRF is associated with long-term mortality risk. However, a few inherent data limitations precluded conclusive assertions as to what extent low CRF is an independent risk factor for long-term

mortality, or if it acts through other pathways involving PA and other known or unknown changes that occurred during the long follow-up period. Regardless of whether the identified associations were causal or predictive, the current study supported the use of CRF as a clinical vital sign (1). Promoting incidental PA such as active transportation in the least fit and least physically active segments of the middle-aged adult population is a safe investment

that will likely lead to improvements in CRF and will certainly save lives.

ADDRESS FOR CORRESPONDENCE: Dr. Emmanuel Stamatakis, Charles Perkins Centre, The University of Sydney, Sydney, New South Wales 2006, Australia. E-mail: emmanuel.stamatakis@sydney.edu.au. Twitter: [@M_Stamatakis](https://twitter.com/M_Stamatakis), [@Sydney_Uni](https://twitter.com/Sydney_Uni), [@MaastrichtU](https://twitter.com/MaastrichtU).

REFERENCES

- Ross R, Blair SN, Arena R, et al. Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American Heart Association. *Circulation* 2016;134:e653-99.
- Stamatakis E, Hamer M, O'Donovan G, Batty GD, Kivimaki M. A non-exercise testing method for estimating cardiorespiratory fitness: associations with all-cause and cardiovascular mortality in a pooled analysis of eight population-based cohorts. *Eur Heart J* 2013;34:750-8.
- Clausen JSR, Marott JL, Holtermann A, Gyntelberg F, Jensen MT. Midlife cardiorespiratory fitness and the long-term risk of mortality: 46 years of follow-up. *J Am Coll Cardiol* 2018;72:987-95.
- Holtermann A, Mortensen OS, Burr H, Sogaard K, Gyntelberg F, Suadicani P. Physical demands at work, physical fitness, and 30-year ischaemic heart disease and all-cause mortality in the Copenhagen Male Study. *Scand J Work Environ Health* 2010;36:357-65.
- Glei D, Mestlé F, Vallin J. Diverging trends in life expectancy at age 50: a look at causes of death. In: Crimmins EMPS, Cohen B, editors. *International differences in mortality at older ages: dimensions and sources*. Washington, DC: National Academies Press, 2010:17-48.
- City of Copenhagen: The Technical and Environmental Administration. *Copenhagen city of cyclists, facts and figures 2017*. Copenhagen, Denmark: City of Copenhagen, 2017.
- Blair SN, Kohl HW 3rd, Paffenbarger RS Jr., Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA* 1989;262:2395-401.
- O'Donovan G, Lee I, Hamer M, Stamatakis E. Association of "weekend warrior" and other leisure time physical activity patterns with risks for all-cause, cardiovascular disease, and cancer mortality. *JAMA Int Med* 2017;177:335-42.
- Batacan RB, Duncan MJ, Dalbo VJ, Tucker PS, Fenning AS. Effects of high-intensity interval training on cardiometabolic health: a systematic review and meta-analysis of intervention studies. *Br J Sports Med* 2017;51:494-503.
- Stamatakis E, Kelly P, Strain T, Murtagh EM, Ding D, Murphy MH. Self-rated walking pace and all-cause, cardiovascular disease and cancer mortality: individual participant pooled analysis of 50 225 walkers from 11 population British cohorts. *Br J Sports Med* 2018;52:761-8.
- 2018 Physical Activity Guidelines Advisory Committee. 2018 Physical Activity Guidelines Advisory Committee Scientific Report. Washington, DC: U.S. Department of Health and Human Services, 2018.
- Van Der Velde JHPM, Koster A, Van Der Berg JD, et al. Sedentary behavior, physical activity, and fitness—the Maastricht Study. *Med Sci Sports Exercise* 2017;49:1583-91.

KEY WORDS epidemiology, fitness, mortality, physical activity