

One-dimensional bargaining with unanimity rule

Citation for published version (APA):

Predtetchinski, A. (2007). *One-dimensional bargaining with unanimity rule*. METEOR, Maastricht University School of Business and Economics. METEOR Research Memorandum No. 011
<https://doi.org/10.26481/umamet.2007011>

Document status and date:

Published: 01/01/2007

DOI:

[10.26481/umamet.2007011](https://doi.org/10.26481/umamet.2007011)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Arkadi Predtetchinski

One-dimensional bargaining with unanimity rule

RM/07/011

JEL code: C78



Maastricht research school of **E**conomics
of **T**Echnology and **O**Rganizations

Universiteit Maastricht
Faculty of Economics and Business Administration
P.O. Box 616
NL - 6200 MD Maastricht

phone : ++31 43 388 3830
fax : ++31 43 388 4873

One-dimensional bargaining with unanimity rule

Arkadi Predtetchinski*

March 27, 2007

Abstract

The paper examines bargaining over a one-dimensional set of social states, with a unanimity acceptance rule. We consider a class of δ -equilibria, i.e. subgame perfect equilibria in stationary strategies that are free of coordination failures in the response stage. We show that along any sequence of δ -equilibria, as δ converges to one, the proposal of each player converges to the same limit. The limit, called the bargaining outcome, is uniquely determined by the set of players, the recognition probabilities, and the utility functions, and it is independent of the choice of the sequence. We characterize the bargaining outcome as a unique solution of a characteristic equation.

JEL classification code: C78

Keywords: Bargaining, subgame perfect equilibrium, unanimity rule.

*I am most thankful to Jean-Jacques Herings to whom this work is dedicated. I would also like thank all the participants of a special seminar in the Sobolev Institute of Mathematics (Novosibirsk, Russia).

1 Introduction

This paper studies bargaining in a group of players over a one-dimensional set of alternatives or social outcomes. The alternatives may represent the level of taxation, the amount of the public good, or a location of a facility.

Bargaining proceeds as follows. First, nature chooses a player to make a proposal. Recognition probabilities (i.e. the distribution on the moves of nature) are time-invariant. The chosen player then puts forward a proposal that specifies one alternative. The players react sequentially, in an exogenously given order. Each player can accept or reject a proposal. If the players unanimously agree to a proposal, it is implemented and the procedure ends. Otherwise, a new time period begins, and a new proposer is chosen. Each time a new period begins, a discount factor δ applies.

The central question of the paper is, of course, what alternative or alternatives will be eventually chosen. The paper gives a surprisingly sharp answer to the question as it identifies a specific alternative that is a unique outcome, in a sense made precise below, of the bargaining procedure.

It is clear that the model of bargaining with the unanimity rule as described above is not equally applicable to all instances where negotiations take place. However, there are important real-life examples where the unanimity is indeed required. Thus, peace or disarmament talks are likely to achieve a viable agreement if it is approved upon by all parties.

It is well-known that bargaining procedures such as the one above typically have infinitely many subgame perfect equilibria, including equilibria where delay occurs. For that reason we shall restrict attention to a class of subgame perfect equilibria involving only stationary strategies. Stationarity means that an equilibrium proposal of any player does not depend on the history of play and that the reaction of any player to a proposal only depends on the proposal itself.

When there are finitely many players, in a subgame perfect equilibrium involving only stationary strategies there is no delay. However, with infinitely many players, stationarity alone does not guarantee immediate acceptance of equilibrium proposals and even a perpetual disagreement can be supported as a subgame perfect equilibrium in stationary strategies. Despite the fact that players respond to a proposal sequentially, coordination failures can occur when a proposal is rejected by infinitely many players. We shall only be concerned with subgame perfect equilibria in stationary strategies that are robust to coordination failures in the response stage, in the sense that, whenever a proposal is rejected, for at least one player rejection remains a best response even if all other players, by mistake, accept. In any such equilibrium, no delay ever occurs.

Even in this class of equilibria, the individual acceptance sets are indeterminate. I shall further narrow down the class of strategies by looking at δ -equilibria that pin down a specific functional form for the individual acceptance sets. Restricting attention to δ -equilibria is without loss of generality, however: in terms of equilibrium proposals and equilibrium utilities, the class of δ -equilibria effectively represents all subgame perfect equilibria in stationary strategies that are robust to coordination failures in the response

stage.

The concept of δ -equilibrium as a solution concept for a social choice problem suffers from two obvious disadvantages. One disadvantage is that a given δ -equilibrium does not unambiguously pin down a specific alternative, because different players make different proposals in equilibrium. Furthermore, it depends on the discount factor δ . This motivates us to consider the asymptotic behavior of δ -equilibria as δ converges to one.

We prove that along any sequence of δ -equilibria, as the discount factor δ converges to one, the equilibrium proposals of all players converge to the same limit, and the social acceptance set collapses to a point. This point, called the bargaining outcome, is independent of the choice of the sequence and is uniquely determined by the set of players, recognition probabilities and the utility functions. The main result of the paper is a characterization of the bargaining outcome as the unique solution of a characteristic equation. The characterization is particularly simple in the special case where the player's utility of an alternative x is a linear function of the distance of that player's ideal point from x . In this case, the bargaining outcome is the unique zero of the function g that assigns to each alternative x the mass of players with the ideal points in the interval $(x, 1]$.

In the special case of the model with two players, the bargaining outcome as defined above coincides with the asymmetric Nash bargaining solution, with weights equal to the respective recognition probabilities. In this case the characteristic equation is a first-order condition for the maximization of the asymmetric Nash product.

This paper is closely related to the work Cardona and Ponsati [2]. Cardona and Ponsati [2] study bargaining over a one-dimensional set of social outcomes, with a deterministic recognition rule where a passing of a proposal requires an approval of at least q players. The authors prove stationary equilibrium to be asymptotically unique: for a given quota q , as the discount factor converges to one, the equilibrium proposals of all players approach the same limit, the limit being independent of the recognition sequence. While Cardona and Ponsati [2] prove asymptotic uniqueness of stationary equilibria, they do not provide an explicit computation of the limit. Our results are therefore complementary. We contribute to the insights in [2] by showing that the bargaining outcome can be characterized as a unique zero of a characteristic equation, but our results apply only to a game of bargaining with random recognition and the unanimity acceptance rule.

The setup of this paper is close that in Banks and Duggan [1] and Cho and Duggan [3]. Banks and Duggan [1] consider bargaining over a set of social outcomes that is an arbitrary compact convex subset of an Euclidean space. The bargaining protocol examined in Banks and Duggan [1] has time-invariant recognition probabilities and a general voting rule. A voting rule is represented by a family of decisive coalitions, and the approval of a proposal by any of these coalitions is sufficient for the passing of a proposal. Unanimity rule considered in this paper is a voting rule where the entire player set is the only decisive coalition. Banks and Duggan [1] prove existence of stationary equilibrium and examine the equilibrium set in the case of perfectly patient players (i.e. when the discount factor equals one). The setting in Cho and Duggan [3] is similar to that in Banks and Duggan [1], but the paper focuses on the case of a one-dimensional set of alternatives.

The bargaining protocol studied in this paper is the same as the bargaining protocol

in Banks and Duggan [1] and Cho and Duggan [3] with the unanimity acceptance rule. In particular, the class of δ -equilibria coincides with a class of stationary equilibria as defined in Cho and Duggan [3], under the unanimity rule.

This work builds on a contribution of Herings and Predtetchinski [4]. Herings and Predtetchinski [4] study a bargaining protocol where the identity of a proposer follows a Markov process, and player's utility of an alternative x is a linear function of the distance of that player's ideal point from x . The setup of this paper is much more general with respect to the utility functions: it is only assumed that each utility function is continuous, concave and has a single peak (ideal point). On the other hand, we do not treat a general Markov recognition rule, but restrict attention to the case of time-invariant recognition probabilities.

As we have already mentioned, when there are only two players, the bargaining outcome in a model of one-dimensional bargaining coincides with the asymmetric Nash bargaining solution. With more than two players, the bargaining outcome does not in general maximize the (asymmetric) Nash product. The model of one-dimensional bargaining therefore provides a complementary set of the results to those in Miyakawa [5], where the asymmetric Nash bargaining solution for an n -person bargaining problem is obtained as a limit of stationary equilibria in a game of bargaining with time-invariant recognition probabilities, as the probability of the breakdown of negotiations converges to zero.

The rest of the paper is as follows. The next section presents the most important ideas and definitions including the definition of stationary strategy, a definition of δ -equilibrium, and of bargaining outcome. Section 3 illustrates the key insights by means of an example. We consider a world where player's utility of an alternative x depends linearly on the distance between the player's location and x . For this special case we prove that the bargaining outcome is the unique zero of the function g that maps each alternative x into the mass of players in the interval $(0, x]$.

Section 4 proves existence of δ -equilibrium. Banks and Duggan [1] prove the existence of no-delay stationary equilibria, but they assume finitely many players. We extend the existence result to environments with an arbitrary player set, under appropriate continuity assumptions.

In Section 5 the set of 1-equilibria is analyzed. It is demonstrated that in each 1-equilibrium the social acceptance set is a singleton. We also show that, given a weakly Pareto-efficient alternative, there exists a 1-equilibrium where each player proposes x . Furthermore, it is proved that along any sequence of δ -equilibria as δ converges to one, the social acceptance set collapses to a point.

Section 6 introduces the characteristic function and the main result of the paper. The next section is devoted to the proof of the main result.

Section 8 proves that each δ -equilibrium is a subgame perfect equilibrium. If the number of players is finite, for a given stationary subgame perfect equilibrium there exists a δ -equilibrium with the same proposals and equilibrium utilities. More generally, if a subgame perfect equilibrium in stationary strategies is robust to coordination failures in the response stage, then there exists a δ -equilibrium having the same proposals and equilibrium utilities. In this sense, δ -equilibria effectively represent all subgame perfect equilibria in

stationary strategies that are robust to coordination failures in the response stage.

2 The components of the model

2.1 A description of the world and a game of bargaining

We study a world ω described by the following variables: X , N , \mathcal{A} , μ , and u_\bullet . The symbol X denotes the unit interval $[0, 1]$. This is a space of alternatives or social states the players must choose from. The set N is a set of players that can be finite or infinite. The symbol \mathcal{A} denotes a sigma-algebra of subsets of N , and μ is the probability measure. Thus the triple (N, \mathcal{A}, μ) is a probability space. The probability measure represents the distribution of types within the population. The symbol u_\bullet denotes a collection of utility functions, one for each player. The utility function of player $t \in N$ is $u_t : X \rightarrow [0, 1]$.

Without further mentioning we shall assume that the characteristics of the world satisfy the following assumption.

- (A1) For each $t \in N$ the utility function $u_t : X \rightarrow [0, 1]$ is concave, continuous, and it attains its unique maximum at point \bar{x}_t . The function $\bar{x}_\bullet : N \rightarrow X$ given by $t \mapsto \bar{x}_t$ is \mathcal{A} -measurable.

Given a discount factor $\delta \in [0, 1]$ we define a game of bargaining $\Gamma(\delta)$ as follows. The game starts in period zero. Each period τ begins with nature randomly choosing a player from the set N to make a proposal. A probability for a proposer to be a member of a set $S \in \mathcal{A}$ is $\mu(S)$. The chosen player proposes an alternative x from X . All players (including the proposer) respond. We assume that the players respond sequentially, according to a total order $>$ on the player set N . The order $>$ is fixed throughout the game. Each responder can either accept or reject the current proposal. If the responders unanimously agree to the proposal, the game terminates and the proposal is implemented. Otherwise, period $\tau + 1$ begins.

If alternative x is agreed upon in period τ , player i receives a payoff of $\delta^\tau u_i(x)$. The payoff of perpetual disagreement to any player is zero.

It is well known that games of bargaining with more than two players typically have infinitely many subgame perfect equilibria, including equilibria with delay. In order to make a prediction about an outcome of the game, one would often restrict attention to a smaller class of equilibria, typically a class of subgame perfect equilibria involving only stationary strategies. This approach is also adopted in our paper. The exact definition of a profile of stationary strategies is as follows.

Definition 1 A joint strategy σ is said to be *stationary* if there exist an \mathcal{A} -measurable function $x_\bullet : N \rightarrow X$ and a collection A_\bullet of subsets A_t of X , one for each $t \in N$, with $\cap A_t$ a Borel-measurable set such that (a) whenever player t has to make a proposal, player t proposes x_t , and (b) whenever player t has to respond to a proposal x , player t accepts if and only if $x \in A_t$. The set A_t is called an *individual acceptance set* of player t and $\cap A_t$ is called a *social acceptance set*.

2.2 A δ -equilibrium and a bargaining solution

We shall use the following solution concept for the game $\Gamma(\delta)$, called for brevity δ -equilibrium.

Definition 2 Let $x_\bullet : N \rightarrow X$ be an \mathcal{A} -measurable function and $y_\bullet : N \rightarrow [0, 1]$ be an arbitrary function. Let A_\bullet be a collection of subsets A_t of X for $t \in N$. Let A be a non-empty subset of X . The tuple $(x_\bullet, y_\bullet, A_\bullet, A)$ is said to be a δ -equilibrium of the world ω if the following conditions are satisfied:

$$\begin{aligned} x_t &= \arg \max_{x \in A} u_t(x) \text{ for each } t \in N, \\ y_t &= \int u_t(x_i) d\mu(i) \text{ for each } t \in N, \\ A_t &= \{x \in X \mid u_t(x) \geq \delta y_t\} \text{ for each } t \in N, \\ A &= \cap A_t. \end{aligned}$$

In a δ -equilibrium no delay ever occurs, as all equilibrium proposals are unanimously accepted. As we show in Section 8, if $(x_\bullet, y_\bullet, A_\bullet, A)$ is a δ -equilibrium, then the joint stationary strategy (x_\bullet, A_\bullet) is a subgame perfect equilibrium of the game $\Gamma(\delta)$. Conversely, if the player set N is finite and the joint stationary strategy (x_\bullet, A_\bullet) is a subgame perfect equilibrium, then the equilibrium proposal map x_\bullet is a part of some δ -equilibrium $(x_\bullet, \dot{y}_\bullet, \dot{A}_\bullet, \dot{A})$. In particular, each x_t is unanimously accepted. In general, if the joint stationary strategy (x_\bullet, A_\bullet) is a subgame perfect equilibrium and is free of coordination failures in the response stage of the game (in the sense made precise in Section 8), then the equilibrium proposal map x_\bullet is a part of some δ -equilibrium $(x_\bullet, \dot{y}_\bullet, \dot{A}_\bullet, \dot{A})$.

Since we assume utility functions to be concave, each individual acceptance set is an interval. We shall use the notation $[x_t^-, x_t^+]$ to denote the individual acceptance set A_t of player t . The social acceptance set is also a closed interval, denoted by $[x^-, x^+]$ or $[x_-, x_+]$. Furthermore,

$$x^- = \sup\{x_t^-\} \text{ and } x^+ = \inf\{x_t^+\},$$

where the supremum and the infimum are taken over all $t \in N$. The equilibrium proposal x_t of player t is a point of $[x^-, x^+]$ closest to \bar{x}_t , the ideal point of individual t . Thus

$$x_t = \begin{cases} x^- & \text{if } \bar{x}_t \leq x^- \\ \bar{x}_t & \text{if } x^- \leq \bar{x}_t \leq x^+ \\ x^+ & \text{if } x^+ \leq \bar{x}_t. \end{cases}$$

Notice that the function x_\bullet thus defined is \mathcal{A} -measurable.

Definition 3 For each natural n let $(x_\bullet^n, y_\bullet^n, A_\bullet^n, A^n)$ be a δ^n -equilibrium of the world ω and let x^n be a point in A^n . Suppose that the sequence δ^n converges to one and x^n converges to x . Then the alternative x is called a bargaining outcome of ω . A collection of bargaining outcomes is called a bargaining solution.

The main result of the paper states that the bargaining outcome is unique. In particular, along any sequence of δ -equilibria, as δ converges to one, the social acceptance set collapses to a point, and in the limit all players make the same proposal. The next section illustrates this important insight by an example.

2.3 General remarks

For a subset B of \mathbb{R} we write $\text{Int}B$ to denote the interior of B . In particular, $\text{Int}X = (0, 1)$.

Assumption (A1) implies that each function u_t is positive on $\text{Int}X$ and that it is not a constant on any non-degenerate interval $I \subset X$.

A concave function $f : X \rightarrow \mathbb{R}$ has left and right derivatives at each point x of $\text{Int}X$ denoted by $f'(x-0)$ and $f'(x+0)$. We shall often use the following fact: Given $x \in \text{Int}X$ and $\dot{x} \in X$, the inequality $f(\dot{x}) - f(x) \leq s(\dot{x} - x)$ holds for each $s \in [f'(x-0), f'(x+0)]$.

3 An example

This section builds on the results in Herings and Predtetchinski [4]. Consider a world λ where $N \subset [0, 1]$ is a finite or infinite set of players containing 0 and 1. We assume that \mathcal{A} is a sigma-algebra of Borel subsets of N . The utility function of player $t \in N$ is $u_t(x) = 1 - |x - t|$. Define a function $g : [0, 1] \rightarrow [0, 1]$ by letting $g(x) = \mu(\{i \in N | x < i\})$. Then $1 - g$ is the cumulative distribution function on X induced by the probability measure μ . The function g is non-decreasing and it is continuous if μ is non-atomic, i.e. if $\mu(\{t\}) = 0$ for each $t \in N$. We need the following definition.

Definition 4 Point $x \in X$ is a generalized fixed point of the function $f : X \rightarrow X$ if there are sequences x_-^n and x_+^n of points in X converging to x such that $\lim f(x_-^n) \leq x \leq \lim f(x_+^n)$.

It is clear that any fixed point x of f is also a generalized fixed point of f (take $x_-^n = x$ and $x_+^n = x$). Conversely, if f is continuous, then any generalized fixed point of f is also its fixed point. If f is a non-increasing function, then it has exactly one generalized fixed point.

Proposition 1

- (i) For each $\delta \in [0, 1)$ the world λ has a unique δ -equilibrium. The equilibrium proposals are given by Figure 1 below, where $[x^-, x^+]$ is a social acceptance set.
- (ii) Let $N = [0, 1]$ and suppose that $\mu(B) = \mu(1 - B)$ for each Borel-measurable set $B \subset N$, where $1 - B = \{1 - t | t \in B\}$. Then $x^- = \delta/2$ and $x^+ = 1 - \delta/2$, and the unique bargaining outcome of the world λ is $1/2$.
- (iii) The world λ has a unique bargaining outcome being the generalized fixed point of the function g .




Figure 1: Equilibrium proposal x_t in the example of Section 3.

Proof. Let $(x_\bullet, y_\bullet, A_\bullet, A)$ be a δ -equilibrium of λ and let $A_t = [x_t^-, x_t^+]$ and $A = [x^-, x^+]$. First we show that the triple (x_\bullet, x^-, x^+) satisfies the following equations:

$$x_t = \begin{cases} x^- & \text{if } t \leq x^- \\ t & \text{if } x^- \leq t \leq x^+ \\ x^+ & \text{if } x^+ \leq t, \end{cases} \quad (1)$$

$$x^- = \delta E(x_\bullet) \text{ and } x^+ = 1 - \delta + \delta E(x_\bullet). \quad (2)$$

The proposal x_t of player t is the point of $[x^-, x^+]$ which is closest to t , as is illustrated in Figure 1 above, whence Equation 1. To derive Equation 2, we compute:

$$y_t = 1 - \int |x_i - t| d\mu(i),$$

$$x_t^- = (t - [1 - \delta y_t]) \vee 0 \text{ and } x_t^+ = (t + [1 - \delta y_t]) \wedge 1.$$

Now one can see that $x_\bullet^- : N \rightarrow X$ and $x_\bullet^+ : N \rightarrow X$ are non-decreasing functions. It follows that $x^- = x_1^-$ and $x^+ = x_0^+$. Because $u_1(x) = x$, we have $y_1 = E(x_\bullet)$ and therefore $x_1^- = \delta E(x_\bullet)$. Because $u_0(x) = 1 - x$, we have $y_0 = 1 - E(x_\bullet)$ and therefore $x_0^+ = 1 - \delta + \delta E(x_\bullet)$. Equation 2 follows.

Existence of δ -equilibrium will be proven more generally in Section 4. We prove that for each $\delta \in [0, 1)$ the world λ has at most one δ -equilibrium by showing that the system 1-2 has at most one solution. Suppose the triples (x_\bullet, x^-, x^+) and $(\hat{x}_\bullet, \hat{x}^-, \hat{x}^+)$ both satisfy Equations 1-2. Let $|\hat{x}_\bullet - x_\bullet| = \sup |\hat{x}_t - x_t|$. Then $|\hat{x}^- - x^-| = |\hat{x}^+ - x^+| = \delta |E(\hat{x}_\bullet - x_\bullet)| \leq \delta |\hat{x}_\bullet - x_\bullet|$. On the other hand, as we show in the proof of Proposition 3, $|\hat{x}_\bullet - x_\bullet| \leq |x^- - \hat{x}^-| \vee |x^+ - \hat{x}^+|$. It follows that $(x_\bullet, x^-, x^+) = (\hat{x}_\bullet, \hat{x}^-, \hat{x}^+)$, as desired.

To prove claim (ii) let $x^- = \delta/2$ and $x^+ = 1 - \delta/2$ and let x_\bullet be as in Figure 1. We show that $E(x_\bullet) = 1/2$. It is then clear that the triple (x_\bullet, x^-, x^+) satisfies the system 1-2. Since the system admits but one solution, x_\bullet must be the equilibrium proposal map and $[x^-, x^+]$ the social acceptance set in a δ -equilibrium of λ .

Since $x^+ + x^- = 1$, we have $x_{1-t} + x_t = 1$ for all $t \in N$. Let $\{s_t^n\}$ be a sequence of simple functions converging pointwise to x_\bullet . Define a function \dot{s}_\bullet^n by the equation $\dot{s}_t^n = 1/2(1 + s_t^n - s_{1-t}^n)$ for all $t \in N$. The function \dot{s}_\bullet^n is a simple function and the sequence $\{\dot{s}_\bullet^n\}$ converges pointwise to x_\bullet . Thus $E(\dot{s}_\bullet^n)$ converges to $E(x_\bullet)$.

It remains to show that $E(\dot{s}_\bullet^n) = 1/2$. Suppose $s_\bullet^n = \sum a_k 1_{B_k}$, where B_k are Borel-measurable subsets of N . Then $\dot{s}_\bullet^n = 1/2(1 + \sum a_k 1_{B_k} - \sum a_k 1_{1-B_k})$. Therefore, $E(\dot{s}_\bullet^n) = 1/2(1 + \sum a_k \mu(B_k) - \sum a_k \mu(1 - B_k)) = 1/2$, because $\mu(B_k) = \mu(1 - B_k)$ for all k .

We prove claim (iii). As before, let $(x_\bullet, y_\bullet, A_\bullet, A)$ be a δ -equilibrium of λ with $A = [x^-, x^+]$. We estimate the expected equilibrium proposal from below and from above, as follows. Define maps $z_\bullet^- : N \rightarrow X$ and $z_\bullet^+ : N \rightarrow X$ by the following equations:

$$z_t^- = \begin{cases} x^- & \text{if } t \in [0, x^+] \\ x^+ & \text{otherwise} \end{cases} \quad \text{and} \quad z_t^+ = \begin{cases} x^- & \text{if } t \in [0, x^-] \\ x^+ & \text{otherwise.} \end{cases}$$

Both maps are measurable and $z_t^- \leq x_t \leq z_t^+$ for all $t \in N$. Therefore,

$$E(z_\bullet^-) \leq E(x_\bullet) \leq E(z_\bullet^+), \text{ where}$$

$$E(z_\bullet^-) = (1 - g(x^+))x^- + g(x^+)x^+ = (1 - \delta)g(x^+) + \delta E(x_\bullet),$$

$$E(z_\bullet^+) = (1 - g(x^-))x^- + g(x^-)x^+ = (1 - \delta)g(x^-) + \delta E(x_\bullet).$$

Rewriting the above system yields

$$g(x^+) \leq E(x_\bullet) \leq g(x^-).$$

Let $(x_\bullet^n, y_\bullet^n, A_\bullet^n, A^n)$ be a δ^n -equilibrium of λ with $A = [x_-^n, x_+^n]$. Suppose δ^n converges to one. Without loss of generality assume that both sequences x_-^n and x_+^n converge. Because $x_+^n - x_-^n = 1 - \delta^n$, the sequences x_-^n and x_+^n converge to the same limit, say a point x . The point x is a bargaining outcome of λ . As $x_-^n \leq x_t^n \leq x_+^n$, the equilibrium proposal x_t^n of each player t also converges to x , and so does the expected value $E(x_\bullet^n)$. We know that $g(x_+^n) \leq E(x_\bullet^n) \leq g(x_-^n)$ for each n . Taking the limit, we find that x is a generalized fixed point of g , as desired. \square

4 Existence of δ -equilibrium

The results of this section rely on the following additional assumptions.

(A2) The inequality $\inf\{u_t(\bar{x}_t)\} > 0$ holds.

(A3) The family of functions u_\bullet is jointly continuous: Given an $x \in X$ and an $\epsilon > 0$ there exists a $\varepsilon = \varepsilon(x, \epsilon)$ such that $|u_t(x) - u_t(\dot{x})| < \epsilon$ for all $t \in N$ whenever $|x - \dot{x}| < \varepsilon$.

If the player set N is finite, assumptions **(A2)** and **(A3)** are automatically satisfied. Indeed, assumption **(A1)** implies that $u_t(\bar{x}_t) > 0$ for all $t \in N$, for otherwise the function u_t would have been identically zero. Also, any finite family of continuous functions is jointly continuous.

In the general case, assumption **(A1)** implies that the family u_\bullet is jointly continuous at any point x in the interior of X , so that joint discontinuity can occur only at points 0 and 1. As an example of a family of utility functions that violates **(A3)** consider

$$u_t(x) = \begin{cases} x/t, & \text{if } x \leq t \\ (1-x)/(1-t), & \text{if } x \geq t, \end{cases}$$

where the player set is $N = (0,1)$. Then the family u_\bullet satisfies assumptions **(A1)** and **(A2)** but it is not uniformly continuous at either point of the boundary of X .

Consider also the following assumption.

(A4) The family of functions u_\bullet is jointly uniformly continuous: Given an $\epsilon > 0$ there exists a $\varepsilon = \varepsilon(\epsilon)$ such that $|u_t(x) - u_t(\dot{x})| < \epsilon$ for all $t \in N$ whenever $|x - \dot{x}| < \varepsilon$.

Assumption **(A4)** differs from **(A3)** in that ε depends only on ϵ but not on x , whereas in **(A3)** it can depend on both. In fact, the two conditions are equivalent. To see this, let F denote the set of all functions $N \rightarrow X$ endowed with a usual sup-norm: $|x_\bullet - \dot{x}_\bullet| = \sup |x_t - \dot{x}_t|$ for x_\bullet and \dot{x}_\bullet in F . Let u_\bullet denote a map $X \rightarrow F$ given by $x \mapsto u_\bullet(x)$. Then assumption **(A3)** can be equivalently stated as saying that the map u_\bullet is continuous. Assumption **(A4)**, on the other hand, is equivalent to a requirement that the map u_\bullet be uniformly continuous. As X is a compact space, any map from X to a metric space is uniformly continuous if and only if it is continuous. Thus assumption **(A3)** can be replaced with **(A4)** without loss of generality.

Theorem 1 *Let $\delta \in [0,1)$. Suppose ω satisfies the assumptions **(A1)**, **(A2)** and **(A3)**. Then ω has a δ -equilibrium.*

To prove Theorem 1 we characterize the δ -equilibrium as a fixed point of a continuous map.

Let F denote the set of all functions $N \rightarrow X$ endowed with a sup-norm. Let D be a collection of all closed intervals $[a,b]$ in X . We shall identify D with a subspace $\{(a,b) \in X \times X | a \leq b\}$ of $X \times X$. Let M denote the set of all \mathcal{A} -measurable functions $x_\bullet : N \rightarrow X$. Let Y be a set of maps $y_\bullet : N \rightarrow X$ such that there exists an $x \in X$ with $y_t \leq u_t(x)$ for all $t \in N$. The sets M and Y are considered subspaces of F . Let S be a set of all families of closed intervals A_\bullet in X such that $\cap A_t$ is non-empty. Alternatively, one can think of S as a set of pairs $(x_\bullet^-, x_\bullet^+)$ of maps $x_\bullet^- : N \rightarrow X$ and $x_\bullet^+ : N \rightarrow X$, such that $x_t^- \leq x \leq x_t^+$ for all $t \in N$ for some $x \in X$. Then S can be considered a subspace of $F \times F$.

Define f to be a composite map

$$D \xrightarrow{f_1} M \xrightarrow{f_2} Y \xrightarrow{f_3} S \xrightarrow{f_4} D, \text{ where}$$

$$f_1(x^-, x^+)_t = \begin{cases} x^- & \text{if } \bar{x}_t \leq x^- \\ \bar{x}_t & \text{if } x^- \leq \bar{x}_t \leq x^+ \text{ for each } (x^-, x^+) \in D, \\ x^+ & \text{if } x^+ \leq \bar{x}_t \end{cases}$$

$$f_2(x_\bullet)_t = \int_i u_t(x_i) d\mu(i) \text{ for each } x_\bullet \in M,$$

$$f_3(y_\bullet)_t = \{x \in X \mid \delta y_t \leq u_t(x)\} \text{ for each } y_\bullet \in Y,$$

$$f_4(A_\bullet) = \cap A_t \text{ for each } A_\bullet \in S.$$

The map f_4 can also be written as

$$f_4(x_\bullet^-, x_\bullet^+) = (\sup\{x_t^-\}, \inf\{x_t^+\}) \text{ for each } (x_\bullet^-, x_\bullet^+) \in S.$$

It is obvious that an interval $[x^-, x^+]$ is a social acceptance set in a δ -bargaining outcome if and only if it is a fixed point of the map f . The map f_1 maps each interval A to an equilibrium proposal map. The map f_2 transforms equilibrium proposals into a collection of expected utilities, whereas f_3 maps expected utilities to individual acceptance sets. Finally, f_4 maps individual acceptance sets to a social acceptance set.

It remains to show that the map f is continuous. First we establish the following auxiliary proposition. Define

$$m = \inf \left\{ |u'_t(x-0)| \mid \begin{array}{l} (t, x) \in N \times \text{Int}X \\ u_t(x) \leq \delta u_t(\bar{x}_t) \end{array} \right\}.$$

Proposition 2 *If $\delta \in [0, 1)$, then $m > 0$.*

Proof. Let $t \in N$ and $x \in \text{Int}X$ be such that $u_t(x) \leq \delta u_t(\bar{x}_t)$. Then $u_t(\bar{x}_t) - u_t(x) \leq u'_t(x-0)(\bar{x}_t - x)$. We have thus the following chain of inequalities:

$$|u'_t(x-0)| \geq \frac{u_t(\bar{x}_t) - u_t(x)}{|\bar{x}_t - x|} \geq \frac{(1 - \delta)u_t(\bar{x}_t)}{|\bar{x}_t - x|} \geq (1 - \delta)u_t(\bar{x}_t).$$

It follows that

$$m \geq (1 - \delta) \inf\{u_t(\bar{x}_t)\}$$

Assumption **(A2)** now implies that m is positive. \square

Proposition 3 *If $\delta \in [0, 1)$, then the map f is continuous.*

Proof. The map f_1 is continuous. Let $A = [x^-, x^+]$ and $\dot{A} = [\dot{x}^-, \dot{x}^+]$. Let $f_1(A) = x_\bullet$ and $f_1(\dot{A}) = \dot{x}_\bullet$. The continuity of f_1 follows from the inequality

$$|x_\bullet - \dot{x}_\bullet| \leq |x^- - \dot{x}^-| \vee |x^+ - \dot{x}^+|.$$

To prove the inequality, fix a t and let $\epsilon = |x^- - \dot{x}^-| \vee |x^+ - \dot{x}^+|$. First we show that there is a point in A at a distance of at most ϵ from \dot{x}_t . Similarly, there is a point in \dot{A} at a distance of at most ϵ from x_t . To see this, write x_t as a convex combination of x^- and x^+ and \dot{x}_t as a convex combination of \dot{x}^- and \dot{x}^+ : $x_t = ax^- + (1-a)x^+$ and $\dot{x}_t = a\dot{x}^- + (1-a)\dot{x}^+$, and define the points $\dot{z}_t \in \dot{A}$ and $z_t \in A$ by the equations $\dot{z}_t = a\dot{x}^- + (1-a)\dot{x}^+$ and $z_t = ax^- + (1-a)x^+$. Then

$$\begin{aligned} |\dot{z}_t - x_t| &\leq a|\dot{x}^- - x^-| + (1-a)|\dot{x}^+ - x^+| \leq \epsilon \text{ and} \\ |z_t - \dot{x}_t| &\leq a|\dot{x}^- - x^-| + (1-a)|\dot{x}^+ - x^+| \leq \epsilon. \end{aligned}$$

We also have the following obvious property: Let $x \in A$. If $x < x_t$, then $x_t \leq \bar{x}_t$ and if $x_t < x$, then $\bar{x}_t \leq x_t$. A similar property holds for \dot{A} .

Now we prove that $x_t \leq \dot{x}_t + \epsilon$. Suppose not. Then we have $z_t \leq \dot{x}_t + \epsilon < x_t$. Since z_t is a point of A , this means that $x_t \leq \bar{x}_t$. On the other hand, we also have $\dot{x}_t < x_t - \epsilon \leq \dot{z}_t$. Since \dot{z}_t is a point of \dot{A} , we have $\bar{x}_t \leq \dot{x}_t$. We arrive at a contradiction, because $\bar{x}_t \leq \dot{x}_t < x_t - \epsilon < x_t \leq \bar{x}_t$. The proof that $\dot{x}_t \leq x_t + \epsilon$ is similar.

The map f_2 is continuous. Let $f_2(x_\bullet) = y_\bullet$ and $f_2(\dot{x}_\bullet) = \dot{y}_\bullet$. Let ϵ and ε be as in assumption (A4). Suppose $|x_\bullet - \dot{x}_\bullet| < \varepsilon$. Then $|u_t(x_i) - u_t(\dot{x}_i)| < \epsilon$ for all t and i in N . Therefore,

$$|y_t - \dot{y}_t| \leq \int_i |u_t(x_i) - u_t(\dot{x}_i)| d\mu(i) \leq \epsilon.$$

It follows that $|y_\bullet - \dot{y}_\bullet| \leq \epsilon$.

The map f_3 is continuous. Let $f_3(y_\bullet) = (x_\bullet^-, x_\bullet^+)$ and $f_3(\dot{y}_\bullet) = (\dot{x}_\bullet^-, \dot{x}_\bullet^+)$. The continuity of f_3 is implied by the inequalities

$$|x_\bullet^- - \dot{x}_\bullet^-| \leq \frac{\delta|y_\bullet - \dot{y}_\bullet|}{m} \text{ and } |x_\bullet^+ - \dot{x}_\bullet^+| \leq \frac{\delta|y_\bullet - \dot{y}_\bullet|}{m}.$$

These inequalities are proven by showing that

$$|x_t^- - \dot{x}_t^-| \leq \frac{\delta|y_t - \dot{y}_t|}{m} \text{ and } |x_t^+ - \dot{x}_t^+| \leq \frac{\delta|y_t - \dot{y}_t|}{m}$$

for all $t \in N$. We prove the first set of inequalities. The proof of the second set of inequalities is analogous.

Recall that

$$x_t^- = \inf\{x \in X | \delta y_t \leq u_t(x)\} \text{ and } \dot{x}_t^- = \inf\{x \in X | \delta \dot{y}_t \leq u_t(x)\}.$$

Thus, $\delta y_t \leq u_t(x_t^-)$ and $\delta \dot{y}_t \leq u_t(\dot{x}_t^-)$. Without loss of generality, assume $\dot{x}_t^- < x_t^-$. In particular, $0 < x_t^-$, implying the equality $\delta y_t = u_t(x_t^-)$. Notice that both points \dot{x}_t^- and x_t^- lie in the interval $[0, \bar{x}_t)$, where u_t is an increasing function. Thus the inequality $u_t(\dot{x}_t^-) < u_t(x_t^-)$ holds. Also, the left derivative of u_t at x_t^- is positive. Thus, we have the inequalities

$$0 < u_t'(x_t^- - 0)(x_t^- - \dot{x}_t^-) \leq u_t(x_t^-) - u_t(\dot{x}_t^-) \leq \delta(y_t - \dot{y}_t).$$

They yield the inequalities

$$0 < x_t^- - \dot{x}_t^- \leq \frac{\delta(y_t - \dot{y}_t)}{u_t'(x_t^- - 0)}.$$

Since y_\bullet is an element of Y , we have the inequality $y_t \leq u_t(\bar{x}_t)$. Therefore, $u_t(x_t^-) = \delta y_t \leq \delta u_t(\bar{x}_t)$. It follows that $m \leq |u_t'(x_t^- - 0)|$. This gives the desired inequality

$$0 < x_t^- - \dot{x}_t^- \leq \frac{\delta(y_t - \dot{y}_t)}{m}.$$

The map f_4 is continuous. Let $f_4(x_\bullet^-, x_\bullet^+) = (x^-, x^+)$ and $f_4(\dot{x}_\bullet^-, \dot{x}_\bullet^+) = (\dot{x}^-, \dot{x}^+)$. We have an obvious inequality $|x_t^- - \dot{x}_t^-| \leq |x_\bullet^- - \dot{x}_\bullet^-|$. Then $x_t^- - |x_\bullet^- - \dot{x}_\bullet^-| \leq \dot{x}_t^- \leq \dot{x}^-$ for all t , hence $x^- - |x_\bullet^- - \dot{x}_\bullet^-| \leq \dot{x}^-$. Also, $\dot{x}_t^- - |x_\bullet^- - \dot{x}_\bullet^-| \leq x_t^- \leq x^-$ for all t , hence $\dot{x}^- - |x_\bullet^- - \dot{x}_\bullet^-| \leq x^-$. Thus, $|x^- - \dot{x}^-| \leq |x_\bullet^- - \dot{x}_\bullet^-|$. Similarly one derives an inequality $|x^+ - \dot{x}^+| \leq |x_\bullet^+ - \dot{x}_\bullet^+|$. Together they imply the continuity of f_4 . \square

5 The asymptotic behavior of δ -equilibria

This section is devoted to Theorems 2 and 3 below. The first of these describes the set of δ -equilibria when $\delta = 1$ and shows that in each 1-equilibrium the social acceptance set is a singleton. The second theorem shows that along any sequence of δ -equilibria as δ converges to one, the social acceptance set collapses to a point.

An alternative x is said to be weakly Pareto-efficient if there is no $\dot{x} \in X$ such that $u_t(\dot{x}) > u_t(x)$ for all $t \in N$. Each alternative in the open interval $(\inf\{\bar{x}_t\}, \sup\{\bar{x}_t\})$ is weakly Pareto-efficient. Furthermore, all alternatives in $\cup\{\bar{x}_t\}$ are obviously weakly Pareto-efficient. On the other hand, each weakly Pareto-efficient alternative lies in the closed interval $[\inf\{\bar{x}_t\}, \sup\{\bar{x}_t\}]$. If the player set is infinite it is easy to construct a family of the utility functions in such a way that the alternatives $\inf\{\bar{x}_t\}$ and $\sup\{\bar{x}_t\}$ are not weakly efficient.

Theorem 2 *In each 1-equilibrium the social acceptance set is a singleton. Conversely, given a weakly Pareto-efficient alternative x , there exists a 1-equilibrium with a social acceptance set consisting of point x alone.*

As in the preceding section, we shall identify an interval $[a, b] \subset X$ with a point $(a, b) \in X \times X$. Thus the collection of all closed intervals in X can be seen as a metric space.

Theorem 3 *For each natural number n let A^n be a social acceptance set in a δ^n -equilibrium. If the sequence A^n converges to an interval A and the sequence δ^n converges to 1, then A is a singleton.*

The second part of Theorem 2 is easy to prove. Thus let x be a weakly Pareto-efficient alternative. Define $(x_\bullet, y_\bullet, A_\bullet, A)$ by letting x_\bullet be identically equal to x , $y_t = u_t(x)$ and

$A_t = \{\dot{x} \in X \mid u_t(\dot{x}) \geq u_t(x)\}$ for all $t \in N$ and $A = \cap A_t$. We show that the set A consists of point x alone. It then follows that $(x_\bullet, y_\bullet, A_\bullet, A)$ is a 1–equilibrium.

It is clear that x is an element of A . Suppose A contains an open interval I . For each point \dot{x} in I it must hold that $u_t(\dot{x}) > u_t(x)$, for otherwise the function u_t would be constant on I , contradicting assumption **(A1)**. But then x is not a weakly Pareto–efficient alternative, a contradiction. It follows that A is a singleton.

To prove the rest of Theorem 2 and Theorem 3 it is convenient to consider an auxiliary concept of pseudo–equilibrium.

Definition 5 Let $x_\bullet : N \rightarrow X$ be an \mathcal{A} –measurable function and $y_\bullet : N \rightarrow [0, 1]$ be an arbitrary function. Let A_\bullet be a collection of subsets A_t of X for $t \in N$. Let A be a non–empty subset of X . The tuple $(x_\bullet, y_\bullet, A_\bullet, A)$ is said to be a *pseudo–equilibrium* if it satisfies the first two conditions of Definition 2 and

$$A_t = \{x \in X \mid u_t(x) \geq y_t\} \text{ for each } t \in N,$$

$$A \subset \cap A_t.$$

It is clear that a 1–equilibrium is also a pseudo–equilibrium. Thus Proposition 4 below implies the first part of Theorem 2. Together Propositions 4 and 5 imply Theorem 3.

Proposition 4 *If $(x_\bullet, y_\bullet, A_\bullet, A)$ is a pseudo–equilibrium, then A is a singleton.*

Proof. Let $(x_\bullet, y_\bullet, A_\bullet, A)$ be a pseudo–equilibrium. For each natural number n define the set $A_t^n = \{x \in X \mid u_t(x) \geq y_t + 1/n\}$. Observe that A_t^n is Borel–measurable being a closed subset of X . Consequently, the set $\{i \in N \mid x_i \in A_t^n\}$ is \mathcal{A} –measurable. Now,

$$\begin{aligned} y_t &= \int_{\{i \in N \mid x_i \in A_t^n\}} u_t(x_i) d\mu(i) + \int_{\{i \in N \mid x_i \in A_t^n\}^c} u_t(x_i) d\mu(i) \geq \\ &\geq (y_t + 1/n)\mu(\{i \in N \mid x_i \in A_t^n\}) + y_t\mu(\{i \in N \mid x_i \in A_t^n\}^c) = \\ &= y_t + (1/n)\mu(\{i \in N \mid x_i \in A_t^n\}). \end{aligned}$$

Thus we must have $\mu(\{i \in N \mid x_i \in A_t^n\}) = 0$. Define $B_t = \{x \in X \mid u_t(x) > y_t\}$. Clearly, $B_t = \cup_{n=1}^\infty A_t^n$. Therefore,

$$\mu(\{i \in N \mid x_i \in B_t\}) \leq \sum_{n=1}^\infty \mu(\{i \in N \mid x_i \in A_t^n\}) = 0.$$

We shall prove that there is a player $t \in N$ such that $u_t(x_t) = y_t$. If this is indeed the case, then the function u_t is constant on the set A , because $u_t(x_t)$ is an upper bound for u_t on the set A , while y_t is a lower bound. Because A is convex and u_t is strictly quasi–concave, it would then follow that A is singleton.

Suppose that $u_t(x_t) > y_t$ for all $t \in N$. Thus $x_t \in B_t$, so that the family of sets $\{B_t\}_{t \in N}$ is an open cover of the set $\cup_{t \in N} \{x_t\}$. Then there exists a countable subset $C \subseteq N$ such

that the subfamily $\{B_t\}_{t \in C}$ covers $\cup_{t \in N} \{x_t\}$. This is a consequence of the fact that X has a countable base; thus any open cover has a countable subcover.

It follows that

$$N = \bigcup_{t \in C} \{i \in N \mid x_i \in B_t\}.$$

Therefore,

$$\mu(N) \leq \sum_{t \in C} \mu(\{i \in N \mid x_i \in B_t\}) = 0,$$

which is impossible because $\mu(N) = 1$. The result follows. \square

Proposition 5 *Let $(x_\bullet^n, y_\bullet^n, A_\bullet^n, A^n)$ be a sequence of δ^n -equilibria. Suppose the sequence A^n converges to an interval A and the sequence δ^n converges to 1. Then there exist maps x_\bullet and y_\bullet and a collection of sets A_\bullet such that $(x_\bullet, y_\bullet, A_\bullet, A)$ is a pseudo-equilibrium.*

Proof. Define x_\bullet , y_\bullet and A_\bullet by the following equations

$$x_t = \arg \max_{x \in A} u_t(x) \text{ for each } t \in N,$$

$$y_t = \int u_t(x_i) d\mu(i) \text{ for each } t \in N,$$

$$A_t = \{x \in X \mid u_t(x) \geq y_t\} \text{ for each } t \in N.$$

Let the map f_1 be as in Section 4. Recall that f_1 carries an interval A into a map x_\bullet where each x_t is the point of A closest to \bar{x}_t . Thus $x_\bullet^n = f_1(A^n)$ and $x_\bullet = f_1(A)$. As we have seen in the proof of Proposition 3, the map f_1 is continuous with respect to the topology of uniform convergence. Therefore, the sequence x_\bullet^n converges uniformly to x_\bullet .

Since each function u_t is continuous, the sequence $u_t(x_i^n)$ converges to $u_t(x_i)$ for all $i \in N$. Because the integral is continuous with respect to a topology of pointwise convergence, y_t^n converges to y_t for all $t \in N$. (We do not claim that the sequence y_\bullet^n converges uniformly to y_\bullet . This is not true, unless we assume that the family u_\bullet is jointly continuous).

Let $A^n = [x_-^n, x_+^n]$ and $A = [x^-, x^+]$. We know that $A^n \subset \cap A_t^n$. In particular, $u_t(x_-^n) \geq \delta^n y_t^n$ and $u_t(x_+^n) \geq \delta^n y_t^n$ for all $t \in N$. Taking the limits, we obtain the inequalities $u_t(x^-) \geq y_t$ and $u_t(x^+) \geq y_t$. This implies that $A \subset \cap A_t$, as desired. \square

6 The characteristic function

For each $x \in \text{Int}X$ we define

$$\varphi^-(x) = \inf \left\{ \frac{u'_t(x-0)}{u_t(x)} \right\}, \quad \varphi^+(x) = \sup \left\{ \frac{u'_t(x+0)}{u_t(x)} \right\} \text{ and}$$

$$\xi(x) = \mu(\{i \in N \mid \bar{x}_i < x\})\varphi^-(x) + \mu(\{i \in N \mid x < \bar{x}_i\})\varphi^+(x).$$

Claim (a) of Proposition 9 implies that the numbers $\varphi^-(x)$, $\varphi^+(x)$ and $\xi(x)$ are finite. The function ξ is referred to as a *characteristic function*. It will be convenient to extend the characteristic function to X by letting $\xi(0) = +\infty$ and $\xi(1) = -\infty$.

There is nothing particularly important about the use of left derivatives in the function φ^- and right derivatives in φ^+ . As the reader can verify, all results remain true if one replaces a derivative $u'_t(x-0)$ or $u'_t(x+0)$ by an arbitrary element in the subgradient $[u'_t(x+0), u'_t(x-0)]$ of the function u_t .

Proposition 6 *The function φ^+ is positive and decreasing on the interval $(0, \sup\{\bar{x}_i\})$. The function φ^- is negative and decreasing on the interval $(\inf\{\bar{x}_i\}, 1)$. The characteristic function ξ is decreasing on X .*

Proof. Let $\ell_t = \ln u_t$. The function is well-defined on $(0, 1)$ because $u_t(x) > 0$ for all $x \in (0, 1)$. Then we can write

$$\varphi^-(x) = \inf\{\ell'_t(x-0)\} \text{ and } \varphi^+(x) = \sup\{\ell'_t(x+0)\}.$$

Since ℓ_t is a concave function, both its left and right derivatives are non-increasing. It follows that both φ^- and φ^+ are non-increasing functions.

We prove that φ^+ is positive on the interval $(0, \sup\{\bar{x}_i\})$. Given $0 < x < \sup\{\bar{x}_i\}$, there exists a player t such that $x < \bar{x}_t$. For this player t it holds that $u'_t(x+0) > 0$, therefore also $\ell'_t(x+0) > 0$. It follows that $\varphi^+(x) > 0$, as desired.

We prove that φ^+ is decreasing on $(0, \sup\{\bar{x}_i\})$. Suppose not. Then there exist $0 < a < b < \sup\{\bar{x}_i\}$ such that $\varphi^+(a) \leq \varphi^+(b)$. As we already know that the function φ^+ is non-increasing, it has to be a constant (say, equal to m) on the interval $[a, b]$. Let t^q be a sequence in N such that $\ell'_{t^q}(b+0) \rightarrow m$. Then for each $x \in [a, b]$ there are the inequalities

$$m \geq \ell'_{t^q}(a+0) \geq \ell'_{t^q}(x+0) \geq \ell'_{t^q}(b+0),$$

where the right-hand side converges to m as q goes to infinity. The first of these inequalities follows from the fact that m is the supremum of $\ell'_{t^q}(a+0)$, while the other two are true because the right derivative of ℓ_t is a non-increasing function. It follows that $\ell'_{t^q}(x+0) \rightarrow m$. Now we have the inequalities

$$\ell'_{t^q}(x+0)(x-a) \leq \ell_{t^q}(x) - \ell_{t^q}(a) \leq \ell'_{t^q}(a+0)(x-a),$$

where the left- and the right-hand sides converge to $m(x-a)$. Thus $\ell_{t^q}(x) \rightarrow \ell_{t^q}(a) + m(x-a)$. Since $x \in [a, b]$ is arbitrary, it means that the sequence $\{\ell_{t^q}\}$ of functions converges pointwise on $[a, b]$ to a linear function $c + m(\bullet - a)$. Therefore, the sequence $\{u_{t^q}\}$ of functions converges pointwise on $[a, b]$ to an exponential function $\exp(c) \exp(m(\bullet - a))$. But this is impossible, because each u_{t^q} is a concave function, and the set of concave functions on a compact interval is closed in the topology of pointwise convergence. This contradiction establishes that φ^+ is decreasing on the interval $(0, \sup\{\bar{x}_i\})$.

The proof that φ^- is negative and decreasing on $(\inf\{\bar{x}_i\}, 1)$ is similar.

Write $\mu(\bar{x}_i < x)$ to denote $\mu(\{i \in N \mid \bar{x}_i < x\})$ and write $\mu(x < \bar{x}_i)$ to denote $\mu(\{i \in N \mid x < \bar{x}_i\})$. Consider the sets

$$X_- = \{x \in \text{Int}X \mid \mu(\bar{x}_i < x) > 0\} \text{ and } X_+ = \{x \in \text{Int}X \mid \mu(x < \bar{x}_i) > 0\},$$

and the sets $X_-^c = \text{Int}X - X_-$ and $X_+^c = \text{Int}X - X_+$. It is clear that the following inclusions hold:

$$X_- \subset (\inf\{\bar{x}_i\}, 1) \text{ and } X_+ \subset (0, \sup\{\bar{x}_i\}).$$

Furthermore, the set $X_-^c \cap X_+^c$ does not contain a non-degenerate interval. For suppose that $[a, b] \subset X_-^c \cap X_+^c$ where $a < b$. Then $\mu(N) \leq \mu(\bar{x}_i < b) + \mu(a < \bar{x}_i) = 0$, a contradiction.

Consider the function $\xi^+(\bullet) = \mu(\bullet < \bar{x}_i)\varphi^+(\bullet)$. The function $\mu(\bullet < \bar{x}_i)$ is identically zero on X_+^c . On the set X_+ the function $\mu(\bullet < \bar{x}_i)$ is positive and non-increasing while the function φ^+ is positive and decreasing. It follows that the function ξ^+ is identically zero on X_+^c and is positive and decreasing on X_+ . Similarly, consider the function $\xi^-(\bullet) = \mu(\bar{x}_i < \bullet)\varphi^-(\bullet)$. The function $\mu(\bar{x}_i < \bullet)$ is identically zero on X_-^c . Furthermore, $\mu(\bar{x}_i < \bullet)$ is positive and non-decreasing on X_- , while φ^- is negative and decreasing on X_- . It follows that ξ^- is identically zero on X_-^c and is negative and decreasing on X_- .

We now show that ξ is a decreasing function on $\text{Int}X$. Suppose not. Then there exist $0 < a < b < 1$ such that $\xi(a) \geq \xi(b)$. Because ξ , ξ^- and ξ^+ are non-increasing functions, all three must be constant on the interval $[a, b]$. From the fact that ξ^- is constant on $[a, b]$ it follows that $[a, b] \subset X_-^c$, while from the fact that ξ^+ is constant it follows that $[a, b] \subset X_+^c$. This contradiction establishes that ξ is a decreasing function on $\text{Int}X$.

Finally, ξ is a decreasing function on X because $\xi(0) = +\infty$ and $\xi(1) = -\infty$. \square

Definition 6 *The point $x \in X$ is a generalized zero of the function $f : X \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ if there are sequences x_-^n and x_+^n of points in X converging to x such that $\lim f(x_-^n) \leq 0 \leq \lim f(x_+^n)$.*

It is clear that any point $x \in X$ such that $f(x) = 0$ is a generalized zero of the function f (take $x_-^n = x$ and $x_+^n = x$). Conversely, if f is a continuous function, then each generalized zero x of f satisfies $f(x) = 0$. If f is a decreasing function, it has at most one generalized zero. We are now in a position to state the main result of the paper.

Theorem 4 (Main result) *Each world ω has a unique bargaining outcome. The bargaining outcome of ω is the unique generalized zero of the characteristic function ξ .*

Before we proceed with the proof, we verify for the world λ of Section 3 that the conclusion of Proposition 1 agrees with the conclusion of Theorem 4. Thus let λ be the world as in Section 3. Let $g(x) = \mu(\{i \in N \mid x < i\})$. Recall that Proposition 1 claims the unique bargaining outcome of the world λ to be a generalized fixed point x_* of the function g .

Proposition 7 *The characteristic function of the world λ can be written as*

$$\xi(x) = \frac{g(x) - x\mu(N \setminus \{x\})}{x(1-x)}$$

in the interior of X . The generalized fixed point x_ of the function g is the generalized zero of the function ξ .*

Proof. We compute:

$$\frac{u'_t(x \pm 0)}{u_t(x)} = \begin{cases} -1/(1-x+t) & \text{if } t < x \\ \mp 1 & \text{if } t = x \\ 1/(1-t+x) & \text{if } x < t. \end{cases}$$

Notice that for a fixed $x \in \text{Int}X$ the functions $u'_t(x \pm 0)/u_t(x)$ are non-decreasing in t . Therefore for each $x \in \text{Int}X$ we have

$$\varphi^-(x) = -\frac{1}{1-x}, \quad \varphi^+(x) = \frac{1}{x},$$

$$\begin{aligned} \xi(x) &= -\frac{\mu(\{i \in N | i < x\})}{1-x} + \frac{\mu(\{i \in N | x < i\})}{x} = \\ &= \frac{\mu(\{i \in N | x < i\}) - x\mu(N \setminus \{x\})}{x(1-x)} = \\ &= \frac{g(x) - x\mu(N \setminus \{x\})}{x(1-x)}. \end{aligned}$$

We argue that $g(x') \leq x_* \leq g(x'')$ whenever x' and x'' are points of X such that $x'' < x_* < x'$. To see this, let the sequences x_-^n and x_+^n be as in Definition 4. Then $x'' < x_+^n$ and $x_-^n < x'$ for n large. Since g is a non-increasing function, we have $g(x') \leq g(x_-^n)$ and $g(x_+^n) \leq g(x'')$. Therefore, we also have $g(x') \leq \lim g(x_-^n) \leq x_* \leq \lim g(x_+^n) \leq g(x'')$.

First we consider the case where x_* lies on the boundary of X . Suppose $x_* = 0$. It follows from the previous paragraph that $g(x) \leq 0$ whenever $0 < x \leq 1$. Therefore $\xi(x) \leq 0$ for all $x \in \text{Int}X$. Since $\xi(0) = +\infty$, it follows that 0 is a generalized fixed point of the function ξ , as desired.

If $x_* = 1$, then $1 \leq g(x)$ whenever $0 \leq x < 1$. As $x\mu(N \setminus \{x\}) \leq 1$ it follows that $0 \leq \xi(x)$ for all $0 \leq x < 1$. Since $\xi(1) = -\infty$, the point 1 is a generalized fixed point of ξ .

Finally, consider the case where x_* is in the interior of X . Let $z_+^n = x - 1/n$ and $z_-^n = x + 1/n$. Notice that $g(z_-^n) \leq x_* \leq g(z_+^n)$ for all n because $z_+^n < x_* < z_-^n$. Therefore, $\lim g(z_-^n) \leq x_* \leq \lim g(z_+^n)$.

We argue that $\lim \mu(\{z_-^n\}) = 0$ and $\lim \mu(\{z_+^n\}) = 0$. Suppose the first equation is false. Replacing, if necessary, the sequence $\{z_-^n\}$ by a subsequence we have $\mu(\{z_-^n\}) > \epsilon$. However,

since no two points in the sequence $\{z_-^n\}$ are the same, $\mu(\cup\{z_-^n\}) = \sum \mu(\{z_-^n\}) \geq +\infty$, a contradiction. It follows that $\lim \mu(N \setminus \{z_-^n\}) = 1$ and $\lim \mu(N \setminus \{z_+^n\}) = 1$.

Now, we compute

$$\begin{aligned}\lim \xi(z_-^n) &= \frac{\lim g(z_-^n) - x_*}{x_*(1 - x_*)} \leq 0, \\ \lim \xi(z_+^n) &= \frac{\lim g(z_+^n) - x_*}{x_*(1 - x_*)} \geq 0,\end{aligned}$$

establishing that x_* is a generalized zero of ξ . \square

7 A proof of the main result

Proposition 8 *Let $f : X \rightarrow \mathbb{R}$ be a non-negative concave function. Suppose that f is positive on the interior of X .*

(a) *For each $x \in \text{Int}X$ the following inequalities hold:*

$$-\frac{1}{1-x} \leq \frac{f'(x+0)}{f(x)} \leq \frac{f'(x-0)}{f(x)} \leq \frac{1}{x}.$$

(b) *Given an $x \in X$ and a number $0 < \kappa < 1$ define the sets $I(x, \kappa)$ and $B(x, \kappa)$ by the following equations:*

$$\begin{aligned}I(x, \kappa) &= \left[\frac{x}{1 - \ln \kappa}, \frac{x - \ln \kappa}{1 - \ln \kappa} \right], \\ B(x, \kappa) &= \{\dot{x} \in X \mid f(\dot{x}) \geq \kappa f(x)\}.\end{aligned}$$

Then $I(x, \kappa) \subset B(x, \kappa)$.

Proof. Consider a chain of inequalities

$$-\frac{f(x)}{1-x} \leq \frac{f(1) - f(x)}{1-x} \leq f'(x+0) \leq f'(x-0) \leq \frac{f(x) - f(0)}{x} \leq \frac{f(x)}{x}.$$

Dividing by $f(x)$ yields the inequalities of claim (a). Consider a concave function $\ell = \ln f : \text{Int}X \rightarrow \mathbb{R}$. We have

$$\ell'(x+0) = \frac{f'(x+0)}{f(x)} \text{ and } \ell'(x-0) = \frac{f'(x-0)}{f(x)}.$$

Given a point $\bar{x} \in X$, define a function $h : \text{Int}X \rightarrow \mathbb{R}$ by the equation

$$h(x) = \begin{cases} \frac{x - \bar{x}}{x} & \text{if } x \leq \bar{x} \\ \frac{\bar{x} - x}{1 - x} & \text{if } \bar{x} \leq x. \end{cases}$$

The function h is non-positive and concave and it attains the value of zero at point \bar{x} . Then for each $x \in \text{Int}X$ we have the inequalities $h(x) \leq \ell'(x+0)(x - \bar{x}) \leq \ell(x) - \ell(\bar{x})$, where the first inequality can be established using claim (a).

Both intervals $I(\bar{x}, \kappa)$ and $B(\bar{x}, \kappa)$ contain the point \bar{x} , and the interval $I(\bar{x}, \kappa)$ is non-degenerate. To prove the inclusion $I(\bar{x}, \kappa) \subset B(\bar{x}, \kappa)$, it is sufficient to prove that either endpoint of $I(\bar{x}, \kappa)$ is contained in $B(\bar{x}, \kappa)$. Thus let \dot{x} be an endpoint of $I(\bar{x}, \kappa)$. If $\dot{x} = 0$, then $\bar{x} = 0$, and if $\dot{x} = 1$, then $\bar{x} = 1$. In either case the point $\dot{x} = \bar{x}$ is an element of $B(\bar{x}, \kappa)$. If \dot{x} is in the interior of X , it is straightforward to verify that $h(\dot{x}) = \ln \kappa$. Therefore $\ln \kappa \leq \ell(\dot{x}) - \ell(\bar{x})$, implying that $\kappa f(\bar{x}) \leq f(\dot{x})$, as desired. \square

Proposition 9

- (a) For each $x \in \text{Int}X$ the inequalities $-1/(1-x) \leq \varphi^-(x) \leq \varphi^+(x) \leq 1/x$ hold.
- (b) Suppose $\delta \in (0, 1)$ and let $(x_\bullet, y_\bullet, A_\bullet, A)$ be a δ -equilibrium. Then $I(E(x_\bullet), \delta) \subset A$. Consequently, A is a non-degenerate interval.

Proof. Claim (a) of Proposition 9 follows immediately from claim (a) of Proposition 8. Consider the sets

$$B_t = \{x \in X \mid u_t(x) \geq \delta u_t(E(x_\bullet))\}.$$

By claim (b) of Proposition 8, $I(E(x_\bullet), \delta) \subset B_t$ for each t . On the other hand $B_t \subset A_t$, because by Jensen inequality $u_t(E(x_\bullet)) \geq E(u_t(x_\bullet)) = y_t$. The result follows. \square

Proposition 10 Suppose $\delta \in (0, 1)$ and let $(x_\bullet, y_\bullet, A_\bullet, A)$ be a δ -equilibrium. Let $A_t = [x_t^-, x_t^+]$ and $A = [x^-, x^+]$. If $x^- \in \text{Int}X$, then

$$\sup_{x^- < \bar{x}_t} \{\delta y_t / u_t(x^-)\} = 1,$$

where the supremum is taken over all players $t \in N$ with $x^- < \bar{x}_t$. If $x^+ \in \text{Int}X$, then

$$\sup_{\bar{x}_t < x^+} \{\delta y_t / u_t(x^+)\} = 1,$$

where the supremum is taken over all players $t \in N$ with $\bar{x}_t < x^+$.

Proof. We only prove the first claim, the proof of the second claim being similar.

Let $r_t^- = \delta y_t / u_t(x^-)$. We know that $r_t^- \leq 1$ for all t . Take a player t such that x_t^- is in the interior of X . Then $u_t(x_t^-) = \delta y_t$ and we have the following chain of inequalities:

$$\begin{aligned}
0 \leq 1 - r_t^- &= 1 - \frac{u_t(x_t^-)}{u_t(x^-)} = \\
&= \frac{u_t(x^-) - u_t(x_t^-)}{u_t(x^-)} \leq \\
&\leq \frac{u_t(x^-) - u_t(x_t^-)}{u_t(x_t^-)} \leq \\
&\leq \frac{u_t'(x_t^- + 0)}{u_t(x_t^-)} (x^- - x_t^-) \leq \\
&\leq \frac{x^- - x_t^-}{x_t^-},
\end{aligned}$$

where the last inequality follows from claim (a) of Proposition 8.

First we show that $\sup\{r_t^-\} = 1$, where the supremum is taken over all players is N . Since $x^- = \sup\{x_t^-\}$ there is an increasing sequence $x_{t^n}^-$ converging to x^- . Of course, $x_{t^n}^-$ is in the interior of X for n large enough, so the inequalities of the previous paragraph apply. In particular,

$$0 \leq 1 - r_{t^n}^- \leq \frac{x^- - x_{t^n}^-}{x_{t^n}^-}.$$

Since the sequence on the right side converges to zero, the sequence $r_{t^n}^-$ converges to one. This proves that $\sup\{r_t^-\} = 1$.

Now, for any player t with $\bar{x}_t \leq x^-$ the function u_t is non-increasing on the interval $A = [x^-, x^+]$. In particular, $u_t(x) \leq u_t(x^-)$ for any $x \in A$. It follows that $y_t \leq u_t(x^-)$, since y_t is an integral of $u_t(x_i)$ where each x_i is a point of A . Thus $r_t^- \leq \delta$ for any t with $\bar{x}_t \leq x^-$. This proves that $\sup_{x^- < \bar{x}_t} \{r_t^-\} = 1$. \square

Proposition 11 *Suppose $\delta \in (0, 1)$. Let $(x_\bullet, y_\bullet, A_\bullet, A)$ be a δ -equilibrium, where $A = [x^-, x^+]$. If $x^+ \in \text{Int}X$, then the following inequalities hold:*

$$\begin{aligned}
\delta \mu(x^+ < \bar{x}_i) \varphi^+(x^+) &\leq \frac{1 - \delta}{x^+ - x^-} \\
\delta \mu(\bar{x}_i < x^+) \varphi^-(x^+) &\leq -\frac{1 - \delta}{x^+ - x^-}.
\end{aligned}$$

If $x^- \in \text{Int}X$, then, the following inequalities hold:

$$\begin{aligned}
-\frac{1 - \delta}{x^+ - x^-} &\leq \delta \mu(\bar{x}_i < x^-) \varphi^-(x^-) \\
\frac{1 - \delta}{x^+ - x^-} &\leq \delta \mu(x^- < \bar{x}_i) \varphi^+(x^-)
\end{aligned}$$

Proof. Suppose that $x^+ \in \text{Int}X$. We prove the first inequality of Proposition 11. Let $x^+ \leq \bar{x}_t$. First we show that

$$\mu(x^+ < \bar{x}_i)[u_t(x^+) - u_t(x^-)] \leq y_t - u_t(x^-).$$

To see this, for $i \in N$ define \hat{x}_i to be x^- if $\bar{x}_i \leq x^+$ and x^+ otherwise. Clearly, \hat{x}_i is a measurable function of i . Furthermore, $\hat{x}_i \leq x_i$ for all $i \in N$. As the function u_t is non-decreasing on the interval $[x^-, x^+]$, we have the inequality $u_t(\hat{x}_i) \leq u_t(x_i)$ for all $i \in N$. The integration gives $\mu(\bar{x}_i \leq x^+)u_t(x^-) + \mu(x^+ < \bar{x}_i)u_t(x^+) \leq y_t$. Subtracting $u_t(x^-)$ from both sides yields the desired inequality.

As before, let $x^+ \leq \bar{x}_t$. From the fact that $\delta y_t \leq u_t(x^-) \leq u_t(x^+)$ it follows that

$$\delta[y_t - u_t(x^-)] \leq (1 - \delta)u_t(x^+).$$

Finally, we use the inequality

$$u'_t(x^+ + 0) \leq \frac{u_t(x^+) - u_t(x^-)}{x^+ - x^-}.$$

Combining them all, we conclude that that for any player t with $x^+ \leq \bar{x}_t$

$$\begin{aligned} \delta \mu(x^+ < \bar{x}_i) \frac{u'_t(x^+ + 0)}{u_t(x^+)} &\leq \delta \mu(x^+ < \bar{x}_i) \frac{[u_t(x^+) - u_t(x^-)]}{u_t(x^+)[x^+ - x^-]} \leq \\ &\leq \delta \frac{y_t - u_t(x^-)}{u_t(x^+)[x^+ - x^-]} \leq \\ &\leq \frac{1 - \delta}{x^+ - x^-}. \end{aligned}$$

For a player t with $\bar{x}_t \leq x^+$ the expression on the extreme left-hand side of the last inequality is non-positive, because $u'_t(x^+ + 0) \leq 0$. This proves the first inequality of Proposition 11.

We now prove the second inequality of Proposition 11. Let $\bar{x}_t < x^+$. First we show that

$$\mu(\bar{x}_i < x^+)[u_t(x^+) - u_t(x_t)] \leq u_t(x^+) - y_t.$$

To see this, for $i \in N$ define \hat{x}_i to be x_t if $\bar{x}_i < x^+$ and x^+ otherwise. Since x_t is the maximum of u_t on $[0, 1]$, $u_t(x_i) \leq u_t(\hat{x}_i)$ for all $i \in N$. The integration yields $y_t \leq \mu(\bar{x}_i < x^+)u_t(x_t) + \mu(x^+ \leq \bar{x}_i)u_t(x^+)$. Subtracting $u_t(x^+)$ from both sides and multiplying by minus one give the desired inequality.

Now since $x^- \leq x_t < x^+$, we have

$$u'_t(x^+ - 0) \leq \frac{u_t(x^+) - u_t(x_t)}{x^+ - x_t} \leq \frac{u_t(x^+) - u_t(x_t)}{x^+ - x^-} \leq 0.$$

Thus for each t with $\bar{x}_t < x^+$ we have the inequalities

$$\begin{aligned} \delta\mu(\bar{x}_i < x^+) \frac{u'_t(x^+ - 0)}{u_t(x^+)} &\leq \delta\mu(\bar{x}_i < x^+) \frac{[u_t(x^+) - u_t(x_t)]}{u_t(x^+)[x^+ - x^-]} \leq \\ &\leq \delta \frac{u_t(x^+) - y_t}{u_t(x^+)[x^+ - x^-]} = \\ &= -\frac{r_t^+ - \delta}{x^+ - x^-}, \end{aligned}$$

where $r_t^+ = \delta y_t / u_t(x^+)$. Using Proposition 10, we obtain

$$\begin{aligned} \delta\mu(\bar{x}_i < x^+) \varphi^-(x^+) &\leq \delta\mu(\bar{x}_i < x^+) \inf_{\bar{x}_t < x^+} \left\{ \frac{u'_t(x^+ - 0)}{u_t(x^+)} \right\} \leq \\ &\leq \inf_{\bar{x}_t < x^+} \left\{ -\frac{r_t^+ - \delta}{x^+ - x^-} \right\} = \\ &= -\frac{1 - \delta}{x^+ - x^-}. \end{aligned}$$

Suppose that $x^- \in \text{Int}X$. We now prove the third inequality of Proposition 11. This proof is similar to the proof of the first inequality. Let $\bar{x}_t \leq x^-$. First we show that

$$u_t(x^+) - y_t \leq \mu(\bar{x}_i < x^-)[u_t(x^+) - u_t(x^-)].$$

To prove this, notice that the function u_t is non-increasing on the interval $[x^-, x^+]$. The expected utility of player t can therefore be estimated from below as $\mu(\bar{x}_i < x^-)u_t(x^-) + \mu(x^- \leq \bar{x}_i)u_t(x^+) \leq y_t$. Subtracting $u_t(x^+)$ from both sides and multiplying by minus one yields the desired inequality. Secondly, the fact that $\delta y_t \leq u_t(x^+) \leq u_t(x^-)$ implies the inequality

$$-(1 - \delta)u_t(x^-) \leq \delta[u_t(x^+) - y_t].$$

Thirdly, we have

$$\frac{u_t(x^+) - u_t(x^-)}{x^+ - x^-} \leq u'_t(x^- - 0).$$

Combining these three inequalities, we conclude that for each player t with $\bar{x}_t \leq x^-$

$$-\frac{1 - \delta}{x^+ - x^-} \leq \delta\mu(\bar{x}_i < x^-) \frac{u'_t(x^- - 0)}{u_t(x^-)}.$$

This inequality also holds if $x^- < \bar{x}_t$ since in this case $0 \leq u'_t(x^- - 0)$. This proves the third inequality of Proposition 11.

The proof of the fourth inequality is similar to the proof of the second one. Let $x^- < \bar{x}_t$. First, we establish

$$y_t - u_t(x^-) \leq \mu(x^- < \bar{x}_i)[u_t(x_t) - u_t(x^-)].$$

This is done by estimating the expected utility of player t from above, as follows: $y_t \leq \mu(\bar{x}_i \leq x^-)u_t(x^-) + \mu(x^- < \bar{x}_i)u_t(x_t)$, and then subtracting $u_t(x^-)$ from both sides. Since $x^- < x_t \leq x^+$, we have

$$0 \leq \frac{u_t(x_t) - u_t(x^-)}{x^+ - x^-} \leq \frac{u_t(x_t) - u_t(x^-)}{x_t - x^-} \leq u'_t(x^- + 0).$$

Thus for each player t with $x^- < \bar{x}_t$ we have the inequalities

$$\begin{aligned} \frac{r_t^- - \delta}{x^+ - x^-} &= \delta \frac{y_t - u_t(x^-)}{u_t(x^-)[x^+ - x^-]} \leq \\ &\leq \delta \mu(x^- < \bar{x}_i) \frac{[u_t(x_t) - u_t(x^-)]}{u_t(x^-)[x^+ - x^-]} \leq \\ &\leq \delta \mu(x^- < \bar{x}_i) \frac{u'_t(x^- + 0)}{u_t(x^-)}. \end{aligned}$$

Finally, using Proposition 10, we obtain

$$\begin{aligned} \frac{1 - \delta}{x^+ - x^-} &= \sup_{x^- < \bar{x}_t} \left\{ \frac{r_t^- - \delta}{x^+ - x^-} \right\} \leq \\ &\leq \delta \mu(x^- < \bar{x}_i) \sup_{x^- < \bar{x}_t} \left\{ \frac{u'_t(x^- + 0)}{u_t(x^-)} \right\} \\ &\leq \delta \mu(x^- < \bar{x}_i) \varphi^+(x^-). \end{aligned}$$

This completes the proof of Proposition 11. \square

Proposition 12 *Each bargaining outcome of the world ω is a generalized zero point of the characteristic function ξ .*

Proof. Let x be a bargaining outcome of ω . By Definition 3, there exist sequences A^n , δ^n , and x^n such that A^n is a social acceptance set in a δ^n -equilibrium of ω , x^n is a point in A^n , the sequence δ^n converges to 1 and x^n converges to x . Let $A^n = [x_-^n, x_+^n]$. Replacing, if necessary, sequences by subsequences, assume that both x_-^n and x_+^n converge. Then Theorem 3 implies that both sequences converge to x .

First suppose that $x \in \text{Int}X$. Then x_-^n and x_+^n both lie in $\text{Int}X$ for n large enough. The first two inequalities in Proposition 11 then imply that $\xi(x_+^n) \leq 0$ while the last two imply $0 \leq \xi(x_-^n)$. It follows that x is a generalized fixed point of the function ξ .

Now suppose $x = 0$. Then $x_+^n < 1$ for n large enough. Since A^n is a non-degenerate interval by Proposition 9, we have $0 < x_+^n$. Then again the first two inequalities in Proposition 11 imply that $\xi(x_+^n) \leq 0$. Since $\xi(0) = +\infty$, it follows that x is a generalized fixed point of ξ . The argument in the case $x = 1$ is similar. \square

8 The analysis of the game

We start our analysis by computing the payoffs induced by a joint stationary strategy $\sigma = (x_\bullet, A_\bullet)$. Let y_t denote the expected payoff to player t at the beginning of the game, let $A = \cap A_t$ be a social acceptance set and N_a be the set of players whose proposal is accepted under σ i.e. $N_a = \{t \in N | x_t \in A\}$. Let N_r denote the set of players whose proposal is rejected, i.e. the complement of N_a . Notice that A is measurable set as it is an intersection of measurable sets, and the set N_a is \mathcal{A} -measurable as it is a preimage of an \mathcal{A} -measurable set under an \mathcal{A} -measurable map. If nature chooses a proposer i from N_a , player t receives a payoff of $u_t(x_i)$, while if nature chooses a proposer from N_r , then player t 's payoff is δy_t . Thus y_t satisfies the following relation:

$$y_t = \int_{N_a} u_t(x_i) d\mu(i) + \delta \mu(N_r) y_t.$$

Solving the equation, we obtain the following expression for y_t :

$$y_t = \begin{cases} \frac{1}{1 - \delta \mu(N_r)} \int_{N_a} u_t(x_i) d\mu(i) & \text{if } \mu(N_a) > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Of course, we are only interested in admissible, that is joint strategies that result in a well-defined payoff to each player at any node of the game. While any joint stationary strategy induces a well-defined payoff at each node of the game, finding sufficient conditions for a general joint strategy to be admissible is a difficult problem on its own and it is certainly beyond the scope of this paper. In what follows, we simply assume that all joint strategies involved are admissible.

Theorem 5 *Let $(x_\bullet, y_\bullet, A_\bullet, A)$ be a δ -equilibrium. Then a joint stationary strategy $\sigma = (x_\bullet, A_\bullet)$ is a subgame perfect equilibrium of $\Gamma(\delta)$.*

The proof of Theorem 5 consists of two steps. The first step is to show that strategy σ is robust to all one-shot deviations. Recall that a strategy $\bar{\sigma}_t$ for player t is said to be a one-shot deviation from σ at node h if it coincides with the strategy σ_t on all nodes but h . The second step establishes the one-shot deviation property for the game $\Gamma(\delta)$. The property states that if there is a profitable deviation from a joint strategy σ , then there is a profitable one-shot deviation. The one-shot deviation property is well-known for finite games, i.e. for games with a finite game tree and finitely many players and less so for infinite games such as $\Gamma(\delta)$.

Proposition 13 *Let $(x_\bullet, y_\bullet, A_\bullet, A)$ be a δ -equilibrium. Let σ denote a joint stationary strategy (x_\bullet, A_\bullet) . Then no player has a one-shot profitable deviation from σ_t in any subgame of $\Gamma = \Gamma(\delta)$.*

Proof. Let $\bar{\sigma}_t$ be a one-shot deviation from σ_t in a subgame $\Gamma(h)$ of Γ starting at node h .

Suppose player t has to make a proposal at node h . Under strategy σ_t player t proposes alternative x_t , which is accepted, leading to a payoff of $u_t(x_t)$ for player t . Suppose under strategy $\bar{\sigma}_t$ player t makes a proposal x . If x is not an element of the social acceptance set $A = \cap A_i$, the proposal x will be rejected. As $\bar{\sigma}_t$ coincides with σ_t on all nodes following h , player t will receive a payoff of δy_t . By the definition of δ -equilibrium, $\delta y_t \leq u_t(x_t)$. If x is an element of A , then x is accepted and player t receives a payoff of $u_t(x)$. However, $u_t(x) \leq u_t(x_t)$, because by the definition of δ -equilibrium, x_t maximizes the function u_t on the set A .

Suppose player t has to react to a proposal x at node h . Suppose x is accepted by player t under strategy σ_t but rejected under strategy $\bar{\sigma}_t$. Then strategy σ_t leads to a payoff of either $u_t(x)$ or δy_t , depending on whether other players accept or reject x , while strategy $\bar{\sigma}_t$ leads to payoff of δy_t . Then $\bar{\sigma}_t$ is not a profitable deviation, because $\delta y_t \leq u_t(x)$, since $x \in A_t$. Conversely, suppose x is rejected by player t under strategy σ_t but is accepted under strategy $\bar{\sigma}_t$. Then strategy σ_t leads to a payoff of δy_t , while strategy $\bar{\sigma}_t$ leads to payoff of either $u_t(x)$ or δy_t , depending on whether x is accepted or rejected by other players. Again, $\bar{\sigma}_t$ cannot be a profitable deviation, since $u_t(x) \leq \delta y_t$, as x is not an element of A_t . \square

Proposition 14 *Let σ be a profile of strategies. If player t has a profitable deviation from σ , then player t has a profitable one-shot deviation from σ .*

Proof. Given a node h we let $\tau(h)$ denote the period node h belongs to. Suppose player t has a profitable deviation $\bar{\sigma}_t$ from σ in the subgame $\Gamma(\bar{h})$ of the game Γ which increases the subgame payoff by $\varepsilon > 0$. First we show that player t has a profitable deviation in the subgame $\Gamma(\bar{h})$ that coincides with σ_t on all nodes corresponding to the first r periods of the subgame $\Gamma(\bar{h})$.

Since the payoff player t can get in period τ is bounded by δ^τ , the subgame payoff for any strategy that agrees with $\bar{\sigma}_t$ on all nodes corresponding to the first τ periods of the subgame differs from the payoff on $\bar{\sigma}_t$ by at most δ^τ . Set $r = \ln(\varepsilon)/\ln(\delta)$ and define the strategy σ_t^r as follows:

$$\sigma_t^r(h) = \begin{cases} \bar{\sigma}_t(h) & \text{if } \tau(h) \leq r + \tau(\bar{h}) \\ \sigma_t(h) & \text{otherwise.} \end{cases}$$

By definition, σ_t^r coincides with $\bar{\sigma}_t$ on all nodes corresponding to the first r periods of the subgame $\Gamma(\bar{h})$. Therefore, the subgame payoff on σ_t^r differs from the payoff on $\bar{\sigma}_t$ by $\delta^r = \varepsilon$ at the most. It follows that σ_t^r is a profitable deviation from σ_t in the subgame $\Gamma(\bar{h})$.

Suppose there is a node h in the subgame $\Gamma(\bar{h})$ with $\tau(h) = r + \tau(\bar{h})$ such that player t has to act at h and strategy σ_t^r is a profitable deviation from σ in the subsubgame $\Gamma(h)$. Then σ_t^r is a profitable one-shot deviation in the subsubgame $\Gamma(h)$. This follows from the fact that each player acts at most once every period. Thus if node h' follows node h and player t has to act at h' , then $\tau(h') > r + \tau(\bar{h})$, implying that $\sigma_t^r(h') = \sigma_t(h')$. In this case the argument is complete.

Suppose there is no node h in the subgame $\Gamma(\bar{h})$ with $\tau(h) = r + \tau(\bar{h})$ such that player t has to act at h and strategy σ_t^r is a profitable deviation in the subsubgame $\Gamma(h)$. In this case we define a new strategy for t as follows:

$$\sigma_t^{r-1}(h) = \begin{cases} \sigma_t^r(h) & \text{if } \tau(h) \leq r + \tau(\bar{h}) - 1 \\ \sigma_t(h) & \text{otherwise.} \end{cases}$$

Then the payoff on σ_t^{r-1} at any node of the subgame $\Gamma(\bar{h})$ is at least as high as the payoff on σ_t^r . In particular, σ_t^{r-1} is a profitable deviation from σ in the subgame $\Gamma(\bar{h})$. Iterating this argument, we shall be able to find a one-shot profitable deviation for player t . \square

We now turn to the converse of Theorem 5. Before we state the result a few remarks are in order concerning players' equilibrium behavior in the response stage of the game.

It is clear that individual acceptance sets are not uniquely determined. Take a subgame perfect equilibrium $\sigma = (x_\bullet, A_\bullet)$. If alternative x is not accepted by player t ($x \notin A_t$), each player i preceding player t in the response stage is indifferent between accepting and rejecting x , since by the stationarity of σ the continuation payoff is the same whether x is rejected by player i or player by t . Starting from σ , one can obtain a new subgame perfect equilibrium by adding x to or subtracting x from the individual acceptance set A_i of player $i < t$. The profile of strategies thus obtained is a subgame perfect equilibrium and it is equivalent to σ in the sense that the proposals, the social acceptance set and equilibrium payoffs are the same.

On top of the indeterminateness of the individual acceptance sets, when there are infinitely many players, coordination failure might occur in the response stage. The coordination failure may be present in equilibrium despite the fact that the players react to a proposal sequentially.

As an example, consider a world with a set of players $N = [0, 1]$. Suppose that the order in which the players react during the response stage is the natural order on N , i.e. player 0 reacts first, player 1 reacts last. Let ι be a sequence in N converging to 1. Let C be a subset of players $\cup \iota_n$. Then any joint strategy $\sigma = (x_\bullet, A_\bullet)$ is a subgame perfect equilibrium, provided only that the sets A_t are empty for each $t \in C$ and $A_1 = X$. The condition that A_t is empty means that player t rejects all proposals, while $A_1 = X$ means that player 1 accepts all proposals. Of course, σ induces perpetual disagreement. Each player $t < 1$ in the response stage is indifferent between accepting and rejecting a proposal, as player t knows that even if he will accept, a proposal will be later rejected by one of the players in C .

It is clear that the situation described above is peculiar to the case where there are infinitely many players. If there were finitely many players, since the last player accepts all proposals, the player who is second-last to respond should accept all proposals with a positive payoff. In the above example there is no second-last player in the sense that each player $t < 1$ is succeeded by a member of C .

To rule out coordination problems as described above we shall require that if alternative x is rejected, then for at least one player t rejection must be a best response not only to

the current strategy of his opponents, but also to a modified profile of strategies where all players following t accept a proposal in question, but continue playing according to the original profile later on. The definition is as follows.

Definition 7 Let σ be a joint strategy and let h be a history of player ending with player k proposing alternative x . Let h_t denote the history following h after which player t has to react to the proposal. Let σ_t^a denote a strategy for player t that coincides with σ_t on all histories other than h_t and $\sigma_t^a(h_t) = a$.

A joint strategy σ is said to be *free of coordination failures in the response stage* if either x is accepted under σ or there is a player t with $\sigma_t(h_t) = r$ such that the strategy σ_t is a best response against a profile of strategies σ_{-t}^a of player t 's opponents in the subgame $\Gamma(\delta)(h_t)$.

A subgame perfect equilibrium σ is free of coordination failures in the response stage provided that any rejected alternative is rejected by at most finitely many players. In this case player t who is the last player in the order $<$ to reject x satisfies the requirement of the above definition. In particular, if the player set N is finite, then any subgame perfect equilibrium is free of coordination failures. On the other hand, a joint strategy considered above violates the condition in Definition 7, as each player t should accept any alternative x having positive utility, provided that the following players following t accept.

Theorem 6 Suppose $0 \leq \delta < 1$. Consider a joint strategy $\sigma = (x_\bullet, A_\bullet)$ with the expected payoff map y_\bullet and the social acceptance set $A = \cap A_t$. Suppose σ is a subgame perfect equilibrium and that it is free of coordination failures in the response stage. Then there exists a δ -equilibrium $(x_\bullet, y_\bullet, B_\bullet, B)$ such that $\text{Int}B \subset A \subset B$.

Proof. The proof uses Proposition 8 of Section 7.

Define the collections of sets B_\bullet , C_\bullet , and the sets B and C as follows:

$$B_t = \{x \in X | u_t(x) \geq \delta y_t\} \text{ and } B = \cap B_t,$$

$$C_t = \{x \in X | u_t(x) > \delta y_t\} \text{ and } C = \cap C_t.$$

All sets are convex, the set B is closed and $\text{Int}B \subset C$.

First we prove the inclusions $C \subset A \subset B$. To prove the first inclusion, suppose x is not an element of A . Let the strategies σ_\bullet^a be as in Definition 7 above. There exists a player t with $\sigma_t(h_t) = r$ such that the strategy σ_t is a best response against a joint strategy σ_{-t}^a of player t 's opponents at node h_t . Playing σ_t when other players use σ_{-t}^a results leads to a payoff of δy_t for player t . On the other hand, playing σ_t^a against σ_{-t}^a leads to an acceptance of x and a payoff of $u_t(x)$. Since the strategy σ_t should be at least as good as the strategy σ_t^a at node h_t , we conclude that $u_t(x) \leq \delta y_t$. This means that $x \notin C_t$ and therefore x is not an element of C .

To prove the second inclusion, let $x \in A$. Let σ_t^r denote a strategy for player t that coincides with σ_t on all histories other than h_t and $\sigma_t^r(h_t) = r$. When other players use

strategies σ_{-t} , the payoff on strategy σ_t^r at node h_t is δy_t , while the payoff on σ_t is $u_t(x)$. Since σ_t is at least as good as σ_t^r when other players use σ_{-t} , we conclude that $u_t(x) \geq \delta y_t$. Thus $x \in B_t$. Since t is arbitrary, $x \in B$.

We prove that the set A is non-empty. As before, let $N_a = \{t \in N | x_t \in A\}$ and let N_r be a complement of N_a . We show that $\mu(N_a) > 0$. If $\mu(N_a) = 0$, then y_\bullet is identically zero. In this case $\text{Int}X \subset C \subset A$. Thus each alternative in the interior of X is unanimously accepted. Take $t \in N_r$ and $x \in \text{Int}X$. Then the strategy $\sigma_t' = (x, A_t)$ improves the payoff of player t at each node where t has to make a proposal, contradicting the assumption that σ is a subgame perfect equilibrium.

We prove that the set B is not a singleton. Suppose B is a singleton consisting of point \bar{x} alone. Then A consists of \bar{x} alone as well and $x_t = \bar{x}$ for all $t \in N_a$. Then and $y_t = \lambda u_t(\bar{x})$, where $\lambda = \mu(N_a)/[1 - \delta\mu(N_r)]$. Let $\kappa = \delta\lambda$. Notice that $0 \leq \kappa < 1$. The set B can now be written as

$$B = \cap \{x \in X | u_t(x) \geq \kappa u_t(\bar{x})\}.$$

If $\kappa = 0$, then $B = X$, and if $0 < \kappa < 1$, then Proposition 8 applies to show that B contains a non-degenerate interval $I(\bar{x}, \kappa)$. In either case, B is a non-singleton, a contradiction.

Since B is a non-empty convex subset of X and it is not a singleton, it has a non-empty interior. Therefore, the set C is non-empty.

We now show that $N_a = N$. Suppose not. Take $t \in N_r$ and $x \in C$. Then strategy $\sigma_t' = (x, A_t)$ improves the payoff to player t at each node where t has to make a proposal, contradicting the subgame perfection hypothesis. Since $N_a = N$, the expected utility of player t can be written as

$$y_t = \int u_t(x_i) d\mu(i).$$

Finally, we have to show that the proposal x_t of player t maximizes the function u_t on B . Suppose not. Then there is an $x \in \text{Int}B$ such that $u_t(x) > u_t(x_t)$. As $\text{Int}B \subset C \subset A$, the point x is unanimously accepted. The strategy $\sigma_t' = (x, A_t)$ is then a profitable deviation from σ_t by player t at any node where t has to make a proposal, a contradiction. \square

References

- [1] JEFFREY S. BANKS AND JOHN DUGGAN: A Bargaining Model of Collective Choice. *American Political Science Review*, **94**: 73–88, 2000.
- [2] DANIEL CARDONA AND CLARA PONSATÍ: Bargaining One-Dimensional Social Choices. *Journal of Economic Theory*, doi: 10.1016/j.jet.2006.12.001, 2007.
- [3] SEOK-JU CHO AND JOHN DUGGAN: Uniqueness of Stationary Equilibria in a One-Dimensional Model of Bargaining. *Journal of Economic Theory*, **113**: 118–130, 2003.
- [4] P. JEAN-JACQUES HERINGS AND ARKADI PREDTETCHINSKI: One-Dimensional Bargaining With Markov Recognition Probabilities. Working paper.

- [5] TOSHIJI MIYAKAWA Non-cooperative Foundation of n -Person Asymmetric Nash Bargaining Solution. Osaka University of Economics, working paper No. 2006-2.