

Protein Supplementation after Exercise and before Sleep Does Not Further Augment Muscle Mass and Strength Gains during Resistance Exercise Training in Active Older Men

Citation for published version (APA):

Holwerda, A. M., Overkamp, M., Paulussen, K. J. M., Smeets, J. S. J., van Kranenburg, J., Backx, E. M. P., Gijsen, A. P., Goessens, J. P. B., Verdijk, L. B., & van Loon, L. J. C. (2018). Protein Supplementation after Exercise and before Sleep Does Not Further Augment Muscle Mass and Strength Gains during Resistance Exercise Training in Active Older Men. *Journal of Nutrition*, 148(11), 1723-1732. <https://doi.org/10.1093/jn/nxy169>

Document status and date:

Published: 01/11/2018

DOI:

[10.1093/jn/nxy169](https://doi.org/10.1093/jn/nxy169)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 25 Apr. 2024

Protein Supplementation after Exercise and before Sleep Does Not Further Augment Muscle Mass and Strength Gains during Resistance Exercise Training in Active Older Men

Andrew M Holwerda,^{1,2} Maarten Overkamp,¹ Kevin JM Paulussen,¹ Joey SJ Smeets,¹ Janneau van Kranenburg,¹ Evelien MP Backx,¹ Annemie P Gijsen,¹ Joy PB Goessens,¹ Lex B Verdijk,^{1,2} and Luc JC van Loon^{1,2}

¹NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Center+, Maastricht, Netherlands and ²Top Institute Food and Nutrition, Wageningen, Netherlands

Abstract

Background: The proposed benefits of protein supplementation on the skeletal muscle adaptive response to resistance exercise training in older adults remain unclear.

Objective: The present study assessed whether protein supplementation after exercise and before sleep augments muscle mass and strength gains during resistance exercise training in older individuals.

Methods: Forty-one older men [mean \pm SEM age: 70 \pm 1 y; body mass index (kg/m²): 25.3 \pm 0.4] completed 12 wk of whole-body resistance exercise training (3 sessions/wk) and were randomly assigned to ingest either protein (21 g protein, 3 g total leucine, 9 g carbohydrate, 3 g fat; n = 21) or an energy-matched placebo (0 g protein, 25 g carbohydrate, 6 g fat; n = 20) after exercise and each night before sleep. Maximal strength was assessed by 1-repetition-maximum (1RM) strength testing, and muscle hypertrophy was assessed at the whole-body (dual-energy X-ray absorptiometry), upper leg (computed tomography scan), and muscle fiber (biopsy) levels. Muscle protein synthesis rates were assessed during week 12 of training with the use of deuterated water (²H₂O) administration.

Results: Leg-extension 1RM increased in both groups (placebo: 88 \pm 3 to 104 \pm 4 kg; protein: 85 \pm 3 to 102 \pm 4 kg; P < 0.001), with no differences between groups. Quadriceps cross-sectional area (placebo: 67.8 \pm 1.7 to 73.5 \pm 2.0 cm²; protein: 68.4 \pm 1.4 to 72.3 \pm 1.4 cm²; P < 0.001) increased in both groups, with no differences between groups. Muscle fiber hypertrophy occurred in type II muscle fibers (placebo: 5486 \pm 418 to 6492 \pm 429 μ m²; protein: 5367 \pm 301 to 6259 \pm 391 μ m²; P < 0.001), with no differences between groups. Muscle protein synthesis rates were 1.62% \pm 0.06% and 1.57% \pm 0.05%/d in the placebo and protein groups, respectively, with no differences between groups.

Conclusion: Protein supplementation after exercise and before sleep does not further augment skeletal muscle mass or strength gains during resistance exercise training in active older men. This study was registered at the Netherlands Trial Registry (www.trialregister.nl) as NTR5082. *J Nutr* 2018;148:1723–1732.

Keywords: aging, exercise, dietary protein, muscle mass, muscle protein synthesis

Introduction

The age-related decline in skeletal muscle mass and strength, termed sarcopenia, is accompanied by impairments in functional capacity and an increased risk of developing chronic metabolic diseases (1, 2). Physical activity, and resistance exercise in particular, may be the only effective strategy to increase muscle mass and strength. Meta-analyses have shown that protein supplementation may further augment the skeletal muscle adaptive response to resistance exercise training (3, 4). However, although some studies have shown greater increases in muscle mass, strength, or both after protein

supplementation during resistance exercise training (5–8), most individual studies have been unable to show such benefits of protein supplementation (9–18). The apparent discrepancy between studies may be attributed to the differences in study design, including the population studied, the exercise training program, and characteristics of the protein supplementation regimen.

Over the past decade, the application of stable isotope tracer methodology has taught us that the postprandial stimulation of muscle protein synthesis is modulated by the type (19–24), amount (23, 25–28), and timing (29–31) of dietary protein




FIGURE 1 CONSORT flow diagram. CONSORT, Consolidated Standards of Reporting Trials.

ingestion. The ingestion of a rapidly digestible protein source (e.g., whey) has been shown to augment the muscle protein synthetic response when compared with more slowly digestible protein sources (20, 23, 24). Furthermore, coingestion of free leucine has been shown to further augment the muscle protein synthetic response to protein ingestion in older individuals (32–35). In addition to enhancing the muscle protein synthetic

response to ingestion of the main meals, our laboratory has shown efficacy for dietary protein ingestion before sleep to increase overnight muscle protein synthesis rates (36, 37) and support recovery from resistance exercise (38–40). We followed up on this work by showing that dietary protein ingestion before sleep augments muscle mass and strength gains during 12 wk of resistance exercise training in healthy, younger men (41). Therefore, we hypothesized that leucine-enriched whey protein supplementation, provided immediately after training and before sleep, would maximize the muscle protein synthetic response to exercise and, as such, further augment muscle mass and strength gains after more prolonged resistance exercise training in older individuals.

Methods

Subjects. A total of 44 healthy, normoglycemic older men volunteered to complete 12 wk of whole-body resistance exercise training (3 sessions/wk) and ingested either protein or an energy-matched placebo after exercise and each night before sleep (Figure 1). Subjects' characteristics are presented in Table 1. All of the subjects were

Supported by the Top Institute Food and Nutrition. Study products were provided by Nutricia Research, Nutricia Advanced Medical Nutrition, Utrecht, Netherlands.

Author disclosures: AMH, MO, KJMP, JSJS, JvK, EMPB, APG, and JPBG, no conflicts of interest. LBV received speaker's fees from Friesland Campina and Nutricia Research. LJCvL has received research grants, consulting fees, speaking honoraria, or a combination of these, from Friesland Campina, Nutricia Research, and PepsiCo. The project is funded by Top Institute Food and Nutrition, a public-private partnership on precompetitive research in food and nutrition. The researchers were responsible for the study design, data collection and analysis, decision to publish, and preparation of the manuscript. The industrial partners contributed to the project through regular discussion.

Address correspondence to LJCvL (e-mail: l.vanloon@maastrichtuniversity.nl).

Abbreviations used: CSA, cross-sectional area; IGF-I, insulin-like growth factor I; OGTT, oral-glucose-tolerance test; SHBG, sex hormone-binding globulin; SPPB, short physical performance battery; 1RM, 1-repetition maximum.

TABLE 1 Subjects' characteristics¹

	Placebo group (n = 20)	Protein-supplemented group (n = 21)	P
Age, y	71 ± 1	69 ± 1	0.10
Body mass, kg	78.7 ± 1.8	77.4 ± 1.9	0.60
Height, m	1.77 ± 0.02	1.74 ± 0.01	0.11
BMI, kg/m ²	25.1 ± 0.5	25.5 ± 0.6	0.56
SMI, kg/m ²	8.5 ± 0.2	8.6 ± 0.1	0.67
Leg volume, L	8.6 ± 0.2	8.3 ± 0.1	0.42
Fasting plasma glucose, mmol/L	5.7 ± 0.1	5.7 ± 0.1	0.89
Fasting plasma insulin, mU/L	7.7 ± 0.8	9.8 ± 1.3	0.21
HbA1c, %	5.6 ± 0.1	5.4 ± 0.1	0.13
HOMA-IR	2.0 ± 0.2	2.5 ± 0.4	0.27
Systolic blood pressure, mm Hg	142 ± 4	142 ± 3	0.97
Diastolic blood pressure, mm Hg	75 ± 2	80 ± 2	0.07
Resting heart rate, beats/min	64 ± 3	61 ± 1	0.32

¹Values are means ± SEMs. Data were analyzed by using a Student's unpaired *t* test. No significant differences were detected between treatment groups. HbA1c, glycated hemoglobin; SMI, skeletal muscle mass index; 1RM, 1-repetition maximum.

screened for medical issues and excluded if any gastrointestinal, cardiovascular, neurological, or renal diseases were present. Fasting and rested blood glucose and glycated hemoglobin were also assessed to screen for the presence of type 2 diabetes. Subjects were cleared to perform resistance exercise by a cardiologist who examined electrocardiograms measured at rest and during submaximal cycling (performed at 70% of age-predicted heart rate maximum). All of the subjects were living independently and had not been participating in any structured, progressive resistance exercise training program within the past 3 y. All of the subjects were informed of the nature and possible risks of the experimental procedures before their written informed consent was obtained. The study was approved by the Medical Ethical Committee of the Maastricht University Medical Center+, Netherlands (METC 15-3-003) and conformed to standards for the use of human subjects in research as outlined in the most recent version of the Helsinki Declaration. This study was registered at the Netherlands Trial Registry (www.trialregister.nl) as NTR5082.

Study design. After inclusion in the study, subjects were randomly allocated in a double-blinded fashion to either a protein or a placebo group. Before and after the exercise training intervention, 2 separate experimental test days were performed within the same week. On the first experimental test day, anthropometric measurements (height, body mass, and leg volume), DXA, and computed tomography scans were performed and a muscle biopsy sample was collected. The second experimental test day was performed 4 d later and included an oral-glucose-tolerance test (OGTT), short physical performance battery (SPPB) testing, and maximal strength assessments [1-repetition maximum (1RM) on leg press and leg extension]. During week 12 of the training, deuterated water (²H₂O) was provided to assess myofibrillar protein synthesis rates.

Diet and physical activity. All of the subjects were instructed to refrain from any exhaustive physical activity and to keep their diet as consistent as possible 72 h before the experimental test days. On all test days, the subjects arrived at the laboratory by car or public transportation in a fasting and rested state. Before the onset of training and during week 12 of training, subjects recorded 3-d dietary intake records to assess potential changes in daily food intake that might have occurred during the intervention period. Food intake records were evaluated by a dietitian and analyzed online with the use of a publicly available nutrition database (NEVO 2016).

Exercise intervention program. Supervised whole-body resistance exercise training was performed 3 times/wk (Monday, Wednesday, and Friday) during the 12-wk exercise training intervention. All sessions were performed between 0800 and 1100. Training consisted of a 5-min warm-up on a cycle ergometer, followed by a warm-up and 4 sets on both the leg press and leg extension machines. Upper body exercises were paired (chest press with lateral pulldown and shoulder press with horizontal row) and were performed in an alternating manner between training sessions, with 2 sets of each exercise performed. After training, subjects performed 5 min of cool-down on the cycle ergometer. During the first 4 wk of training, the workload was increased from 70% 1RM (8 repetitions in each set) to 80% 1RM (10 repetitions). Resting periods of 2 and 3 min were allowed between sets and exercises, respectively. The workload was adjusted under the guidance of the investigators, which was primarily based on the 1RM estimations (weeks 0, 4, and 8). In addition, the workload was increased when >10 repetitions could be performed. Compliance with the resistance exercise training program was 95% ± 1%.

Protein supplementation. Subjects were randomly assigned to ingest either 21 g leucine-enriched whey protein (3 g total leucine) or an energy-matched placebo after exercise and each night before sleep, including rest days. The nutritional composition of the test beverages is shown in **Table 2**. The intervention was performed in a double-blinded fashion whereby treatment randomization and beverage preparation and labeling were performed independent of the study team. Subjects received labeled packages containing dried contents of their randomly assigned beverages, which were dissolved in 250 mL water and ingested by the subjects. All of the beverages were flavored (vanilla or strawberry) to mask their contents. On training days, no food or drinks were allowed between breakfast and lunch other than the experimental beverages. The subjects were allowed to drink water before, during, and after each exercise session. Compliance to the supplementation was assessed by supplementation logs and was confirmed by counting returned supplement packages. Compliance for protein and placebo supplementation was 99.6% ± 0.1%.

TABLE 2 Nutritional composition of the test beverages ingested in a double-blind, randomized fashion after each resistance exercise training session and each night before sleep during 12 wk of resistance exercise training in active older men¹

Component	Protein ²	Placebo ³
Energy, kJ	628	628
Energy, kcal	150	150
Protein, g	20.7	—
EAA, g	10.6	—
Leucine, g	2.8	—
Carbohydrate, g	9.4	24.5
Fat, g	3	5.8
Fiber, g	1.3	—
Cholecalciferol, μg	20	—

¹Dried beverage contents were dissolved in 250 mL water. EAA, essential amino acid (Leu, Ile, Val, Phe, Met, His, Trp, Thr, and Lys).

²The protein drink also contained other micronutrients: sodium (150 mg), potassium (279 mg), chloride (70 mg), calcium (500 mg), phosphorus (250 mg), magnesium (37 mg), iron (2.4 mg), zinc (2.2 mg), copper (270 μg), manganese (0.5 mg), fluoride (0.15 mg), molybdenum (15 μg), selenium (15 μg), chromium (7.5 μg), iodine (20 μg), vitamin A (152 μg), vitamin E (7.5 mg), phyloquinone (12 μg), thiamin (0.23 mg), riboflavin (0.25 mg), nicotinic acid (8.8 mg), pantothenic acid (0.81 mg), pyridoxine (0.76 mg), folic acid (203 μg), cyanocobalamin (3 μg), biotin (6.1 μg), ascorbic acid (32 mg), provitamin A carotenoids (0.3 mg), and choline (56 mg).

³The placebo drink also contained other micronutrients: sodium (123 mg), potassium (154 mg), chloride (329 mg), calcium (0.71 mg), phosphorus (0.3 mg), and iodine (0.3 μg).

Body composition. Body composition was measured at the whole-body and regional level using DXA (Discovery A; Hologic). Anthropometric measurements were assessed using standardized procedures, body mass by a digital scale to within 100 g, and height by a stadiometer to within 0.5 cm. Anatomic cross-sectional area (CSA) of the quadriceps muscle was assessed by computed tomography scanning (Philips Brilliance 64; Philips Medical Systems) before and after the 12-wk exercise training intervention, as described previously (42).

Muscle biopsy sampling. Muscle biopsy samples were obtained from the middle region of the *vastus lateralis*, 15 cm above the patella and 4 cm below entry through the fascia, using the percutaneous needle biopsy technique (43). Muscle samples were prepared for analyses as previously described (41).

Immunohistochemistry. Muscle biopsy specimens were sliced into 5-mm-thick cryosections and stained for muscle fiber type, and type I and II muscle fiber CSA and composition were determined as described in detail previously (13, 18). No differences in fiber circularity were observed in response to training or between groups. The mean fiber numbers included in analyses were 206 ± 16 and 161 ± 11 from the muscle biopsy samples collected before and after the 12-wk training intervention, respectively.

Myofibrillar fractional synthesis rates. Deuterated water ($^2\text{H}_2\text{O}$) was provided throughout week 12 of exercise training to assess average daily myofibrillar protein synthesis rates as described previously (44). In short, subjects ingested 400 mL of 70% deuterated water spread evenly over a 12-h period on the Friday of week 11 of the training intervention. Subjects ingested 50 mL of 70% deuterated water in the morning of each of the 10 remaining days of the intervention period to maintain the body water deuterium enrichment. A muscle biopsy sample was collected in the morning before the Monday training of week 12. Myofibrillar proteins were isolated from this muscle biopsy sample and from the posttraining muscle biopsy sample collected the following Monday, and the increase in ^2H -alanine enrichment was assessed by using GC-pyrolysis-isotope-ratio mass spectrometer (44). Samples were measured in quadruplicate along with a series of known standards every

12 injections. Myofibrillar protein synthesis rates were calculated as previously described (44).

Muscle strength and physical performance. Maximal strength was assessed by 1RM strength tests on leg press and leg extension machines (Technogym) before and after the training intervention. During the subject intake visit, proper lifting technique was demonstrated and practiced and maximal strength was estimated by using the multiple repetitions testing procedure (45). On the second experimental test day, each subject's 1RM was determined as described previously (46). To ensure progressive loading during training, 1RM was estimated at the end of weeks 4 and 8. Posttraining maximal strength was assessed 7 d after the last training session. Physical performance was assessed by the SPPB, which consists of 3 components: balance, gait speed, and chair-rise ability (47).

Glucose tolerance and plasma hormone concentrations. A standard OGTT was performed before and after the training intervention, as described previously (18). Testosterone, free testosterone, growth hormone, insulin-like growth factor I (IGF-I), and sex hormone-binding globulin (SHBG) were assessed in serum samples collected in a resting and fasting state before and after the training intervention. Serum testosterone was measured by using an electrochemiluminescent immunoassay (Cobas 8000 instrument; Roche Diagnostics). Growth hormone and IGF-I were measured by using a chemiluminescent immunometric assay (IDS iSYS instrument; Immunodiagnostic Systems). SHBG was determined in serum by using a chemiluminescent immunometric assay (XPi instrument; Siemens Medical Solutions Diagnostics). Free testosterone was calculated according to Vermeulen et al. (48).

Statistical analysis. All of the data are expressed as means \pm SEMs. Baseline characteristics between groups were compared by using a Student's unpaired *t* test. A 2-factor repeated-measures ANOVA (time \times treatment) with time (pre-post) as a within-subjects factor and treatment group (placebo-protein) as a between-subjects factor was performed for the analysis of dietary intake, appendicular lean body mass, quadriceps CSA, muscle fiber CSA, muscle strength and performance measures, serum hormone concentrations, and the AUC of glucose and insulin

TABLE 3 Energy intake and macronutrient composition of the diet before and after 12 wk of resistance exercise training in active older men who did or did not receive protein supplementation¹

	Placebo (n = 20)		Protein (n = 21)		P		
	0 wk	12 wk	0 wk	12 wk	Treatment \times time interaction	Treatment effect	Time effect
Energy intake, MJ/d							
Diet	9.7 \pm 0.5	10.3 \pm 0.5	9.4 \pm 0.4	9.1 \pm 0.3	0.062	0.151	0.525
Total	9.7 \pm 0.5	11.2 \pm 0.5*	9.4 \pm 0.4	10.0 \pm 0.3*	0.062	0.151	<0.001
Protein intake, g/d							
Diet	93 \pm 4	94 \pm 4	87 \pm 3	81 \pm 3	0.153	0.052	0.299
Total	93 \pm 4	94 \pm 4	87 \pm 3	111 \pm 3*	<0.001	0.204	<0.001
Protein intake, g/kg body mass/d							
Diet	1.19 \pm 0.06	1.17 \pm 0.06	1.14 \pm 0.05	1.04 \pm 0.04	0.164	0.191	0.061
Total	1.19 \pm 0.06	1.17 \pm 0.06	1.14 \pm 0.05	1.43 \pm 0.04*	<0.001	0.138	<0.001
Protein intake, % of energy							
Diet	16 \pm 1	14 \pm 1	16 \pm 1	15 \pm 1	0.664	0.771	0.044
Total	16 \pm 1	14 \pm 1	16 \pm 1	19 \pm 1	<0.001	0.005	0.308
Total fat intake, g/d	89 \pm 6	98 \pm 7	86 \pm 4	80 \pm 5	0.045	0.168	0.701
Total fat intake, % of energy	34 \pm 1	32 \pm 1	34 \pm 1	30 \pm 1	0.336	0.331	0.005
Total carbohydrate intake, g/d	242 \pm 16	303 \pm 16*	235 \pm 13	253 \pm 11*	0.022	0.11	<0.001
Total carbohydrate intake, % of energy	42 \pm 1	45 \pm 2	42 \pm 1	43 \pm 1	0.137	0.531	0.093
Vitamin D intake, $\mu\text{g}/\text{d}$							
Diet	3.7 \pm 0.5	3.8 \pm 0.7	3.5 \pm 0.4	3.0 \pm 0.03	0.531	0.314	0.624
Total	3.7 \pm 0.5	3.8 \pm 0.7	3.5 \pm 0.4	31.6 \pm 0.3*	<0.001	<0.001	<0.001

¹Values are means \pm SEMs. "Diet" indicates only dietary intake data. "Total" indicates dietary intake plus protein or placebo supplementation. *Different from 0 wk, $P < 0.05$.

TABLE 4 Body composition before and after 12 wk of resistance exercise training in active older men who did or did not receive protein supplementation¹

	Placebo (n = 20)			Protein (n = 21)			P		
	0 wk	12 wk	Difference	0 wk	12 wk	Difference	Treatment × time interaction	Treatment effect	Time effect
Body mass, kg	78.7 ± 1.8	80.5 ± 1.8*	+1.8 ± 0.3	77.4 ± 1.9	78.1 ± 1.8*	+0.8 ± 0.2	0.007	0.463	<0.001
Height, m	1.77 ± 0.02	1.77 ± 0.02	0 ± 0	1.74 ± 0.01	1.74 ± 0.01	0 ± 0		0.11	
BMI, kg/m ²	25.1 ± 0.5	25.6 ± 0.5*	+0.6 ± 0.1	25.5 ± 0.6	25.8 ± 0.6*	+0.3 ± 0.1	0.008	0.695	<0.001
Lean body mass, kg	60.8 ± 1.3	62.4 ± 1.3	+1.6 ± 0.3	59.2 ± 1.1	60.2 ± 1.2	+1.1 ± 0.2	0.183	0.278	<0.001
Appendicular lean body mass, kg	26.7 ± 0.7	27.7 ± 0.7	+1.0 ± 0.2	26.0 ± 0.5	26.6 ± 0.5	+0.6 ± 0.1	0.126	0.303	<0.001
Leg lean mass, kg	13.4 ± 0.4	13.9 ± 0.4	+0.7 ± 0.1	13.0 ± 0.2	13.4 ± 0.2	+0.4 ± 0.1	0.146	0.316	<0.001
Fat mass, kg	16.7 ± 1.0	16.3 ± 0.9	-0.5 ± 0.2	16.8 ± 0.9	16.2 ± 0.8	-0.6 ± 0.2	0.690	0.999	<0.001
Body fat, %	20.7 ± 0.9	19.8 ± 0.9	-0.9 ± 0.3	21.1 ± 0.7	20.3 ± 0.7	-0.8 ± 0.2	0.880	0.679	<0.001
Bone mineral content, g	2925 ± 69	2918 ± 69	-8 ± 14	2727 ± 72	2698 ± 69	-30 ± 14	0.279	0.040	0.069
Bone mineral density, g/cm	1.235 ± 0.025	1.232 ± 0.026	-0.003 ± 0.007	1.179 ± 0.024	1.175 ± 0.022	-0.003 ± 0.006	0.921	0.100	0.527

¹Values are means ± SEMs. *Different from 0 wk, $P < 0.05$.

concentrations during the OGTT. In case of a significant interaction, Student's paired t tests were performed to determine time effects within groups. Myofibrillar protein synthesis rates were compared between groups by using a Student's unpaired t test. Significance was set at $P < 0.05$. All calculations were performed with the use of SPSS (version 21.0; IBM Corporation).

Results

Subjects. Subjects' characteristics are presented in Table 1. In total, 44 subjects were included in the experimental protocol. Three subjects were excluded from analyses; one was excluded due to insufficient exercise compliance (>6 missed sessions), one dropped out due to knee aggravation, and one dropped out due to an adverse event that occurred during a training session. Therefore, analysis was performed in 41 subjects, 20 in the placebo group and 21 in the protein group. Training volume load throughout the entire training intervention did not differ between the placebo and protein groups for any of the exercises performed. At baseline, no differences in age, body mass, height, BMI, skeletal muscle mass index, or leg volume were observed between the placebo and protein groups (Table 1). We observed a significant increase in body mass in response to 12 wk of resistance exercise training ($P < 0.001$), with a greater increase observed in the placebo group ($P = 0.007$). We observed a significant increase in BMI in response to 12 wk of resistance exercise training ($P < 0.001$), with a greater increase observed in the placebo group ($P = 0.008$).

Dietary intake. Dietary intake data are shown in Table 3. Analysis of the 3-d dietary intake records collected before and during week 12 of the experimental protocol showed no baseline differences in total daily energy intake between groups but an increase during the intervention ($P < 0.001$). Dietary macronutrient composition did not differ at baseline between groups and did not change during the experimental protocol (Table 3). In particular, daily protein intake from the diet averaged 1.2 ± 0.1 and 1.1 ± 0.1 g/kg body mass/d at baseline in the placebo and protein groups, respectively, and did not change during the training intervention. Supplementation caused daily protein intake relative to body mass to significantly increase in the protein group when compared with the placebo group (treatment × time interaction, $P < 0.001$). Supplementation also resulted in greater daily vitamin D intake in the protein group than in the placebo group (treatment × time interaction, $P < 0.001$).

Body composition. At baseline, no significant differences in body-composition measurements were observed between the placebo and protein groups (Table 4). Whole-body lean mass significantly increased in response to 12 wk of resistance exercise training in the placebo and the protein groups ($P < 0.001$), with no differences between groups. In accordance, appendicular lean mass and leg lean mass significantly increased in response to 12 wk of resistance exercise training in the placebo and protein groups ($P < 0.001$), with no differences between groups. Whole-body fat mass and body fat percentage declined in response to 12 wk of resistance exercise training in the placebo and protein groups ($P < 0.001$), with no differences between groups.

Skeletal muscle hypertrophy. At baseline, no significant differences in quadriceps muscle CSA were observed between the placebo and protein groups (Figure 2). Quadriceps muscle CSA increased in response to 12 wk of resistance exercise training (Figure 2; $P < 0.001$), with no differences between groups. Before training, no significant differences were observed




FIGURE 2 Quadriceps muscle CSA before and after 12 wk of resistance exercise training in active older men who did or did not receive protein supplementation. Values are means ± SEMs, $n = 20$ (placebo) or 21 (protein). Data were analyzed with the use of 2-factor ANOVA. *Different from values before exercise training, $P < 0.05$. CSA, cross-sectional area.




FIGURE 3 Type I (A) and type II (B) muscle fiber CSA before and after 12 wk of resistance exercise training in active older men who did or did not receive protein supplementation. Values are means \pm SEMs, $n = 20$ (placebo) or 20 (protein). Data were analyzed with the use of 2-factor ANOVA. *Different from values before exercise training, $P < 0.05$. CSA, cross-sectional area.

in type I and type II muscle fiber size between placebo and protein groups. Type I muscle fiber CSA (Figure 3A) did not change in response to 12 wk of resistance exercise training in both the placebo and protein groups. Type II muscle fiber CSA (Figure 3B) increased in response to 12 wk of resistance exercise training in both groups ($P < 0.001$), with no differences between groups.

Muscle fiber-type composition. At baseline, no significant differences in the percentage of type I and type II muscle fibers and percentage of CSA occupied by type I and II muscle fibers were observed between the placebo and protein groups (Table 5). The percentage of type I and type II muscle fibers and the percentage of CSA occupied by type I and II muscle fibers did not change in response to 12 wk of resistance exercise training in either group.

Myofibrillar protein synthesis rates. Myofibrillar protein synthesis rates during week 12 of the training protocol were assessed by administration of deuterated water and by measuring myofibrillar protein-bound ^2H -alanine enrichments in muscle biopsy samples collected on Monday of week 12 and the following Monday (posttraining muscle biopsy sample). Body water deuterium enrichment over the 7-d assessment period averaged $0.75\% \pm 0.01\%$ in both groups, with no differences between groups. The increase in myofibrillar protein-bound ^2H -alanine enrichments averaged 0.315 ± 0.015 and 0.308 ± 0.016 mole percent excess in the placebo group and protein group, respectively. Myofibrillar protein synthesis rates (Figure 4) were not different between groups.

TABLE 5 Muscle fiber-type composition before and after 12 wk of resistance exercise training in active older men who did or did not receive protein supplementation¹

	Placebo ($n = 20$)		Protein ($n = 20$)		<i>P</i>		
	0 wk	12 wk	0 wk	12 wk	Treatment \times time interaction	Treatment effect	Time effect
Type I fibers, %	50 \pm 4	48 \pm 3	55 \pm 4	53 \pm 3	0.737	0.252	0.432
Type II fibers, %	50 \pm 4	52 \pm 3	45 \pm 4	47 \pm 3	0.737	0.252	0.432
Type I CSA, %	52 \pm 4	49 \pm 4	58 \pm 4	52 \pm 3	0.662	0.326	0.161
Type II CSA, %	48 \pm 4	51 \pm 4	42 \pm 4	48 \pm 3	0.662	0.326	0.161

¹Values are means \pm SEMs. CSA, cross-sectional area.

Muscle strength and physical performance. At baseline, no significant differences in leg extension or leg press 1RM (Table 6) were observed between the placebo and protein groups. Leg press and leg extension 1RM significantly increased in response to 12 wk of resistance exercise training, with no differences detected between groups. Likewise, physical performance (SPPB) scores (Table 6; $P < 0.01$) and sit-to-stand time decreased in response to 12 wk of resistance exercise training (Table 5; $P < 0.001$), with no differences between groups.

Glucose tolerance. Glucose and insulin concentrations during the OGTT are shown in Figure 5A. Glucose tolerance improved in response to 12 wk of resistance exercise training, as shown by a significant reduction in plasma glucose AUC during the OGTT (Figure 5B; $P = 0.027$), with no differences for plasma insulin AUC and no differences observed between groups.

Plasma hormone concentrations. Serum testosterone, free testosterone, growth hormone, IGF-I, and SHBG concentrations are shown in Table 7. Plasma hormone concentrations did not differ between groups and did not change in response to 12 wk of resistance exercise training.

Discussion

The present study shows that 12 wk of resistance exercise training increased muscle mass and strength in active older men. Protein supplementation after each exercise session (3 times/wk) and every night before sleep did not further




FIGURE 4 Myofibrillar protein FSRs assessed throughout week 12 of the resistance exercise training intervention in active older men who did or did not receive protein supplementation. Values are means \pm SEMs, $n = 20$ (placebo) or 21 (protein). Data were analyzed with the use of Student's unpaired t test. FSR, fractional synthesis rate.

enhance the training-induced gains in skeletal muscle mass or strength in older men.

Twelve weeks of progressive whole-body resistance exercise training resulted in a combined average gain of 1.3 ± 0.2 kg lean body mass and a concomitant decrease in whole-body fat mass. These results are in line with previous studies that showed that 12 wk of resistance exercise training results in a gain of 1.0 kg lean body mass (4, 10, 13, 18). The increase in lean body mass primarily occurred in the lower extremities, with quadriceps CSA being increased by $7\% \pm 1\%$. This increase in muscle mass was accompanied by $15\% \pm 1\%$ and $21\% \pm 3\%$ increases in leg press and leg extension strength, respectively. As expected, our findings are in line with previous studies that showed strength increases of $\geq 25\%$ after a 12-wk resistance exercise training program (13, 18, 49, 50). The observed increase in muscle strength and shift in body composition also contributed to improvements in functional capacity (Table 5) and oral-glucose tolerance (Figure 5).

In the present study, we tested the hypothesis that supplementation with leucine-enriched whey protein ingested after each exercise session and every night before sleep would further augment gains in muscle mass and strength during resistance exercise training in active older men. Despite the significant gains in lean body mass observed over the course of the

training period, we observed no differences between the protein-supplemented and placebo groups (1.1 ± 0.2 and 1.6 ± 0.3 kg, respectively; $P = 0.18$). These findings correspond with more direct assessments of skeletal muscle hypertrophy, because we observed no differences between groups in quadriceps CSA (Figure 2; $P = 0.09$) or type I (Figure 3A; $P = 0.53$) and type II (Figure 3B; $P = 0.84$) muscle fiber CSA. We also observed no differences in leg press 1RM (Table 6; $P = 0.77$), leg extension 1RM (Table 6; $P = 0.80$), or sit-to-stand time (Table 6; $P = 0.74$) between groups.

In addition to our assessments of skeletal muscle mass and strength, we also provided subjects with deuterated water ($^2\text{H}_2\text{O}$) to assess the muscle protein synthetic response to protein supplementation throughout the last week of the training intervention. With the use of this approach, we and others have shown that resistance exercise stimulates a robust increase in muscle protein synthesis rates over days or weeks (44, 51, 52). In line with the present findings on muscle mass, we observed no differences in myofibrillar protein synthesis rates between the placebo and protein groups. Daily fractional muscle protein synthesis rates averaged $1.57\% \pm 0.05\%$ and $1.62\% \pm 0.06\%/d$ in the protein and placebo groups, respectively (Figure 4; $P = 0.54$). Our findings seem to be in contrast with observations of higher muscle protein synthesis rates during recovery from a single bout of exercise when whey protein (20, 23, 24), leucine (34, 35, 53, 54), or leucine-enriched protein (55) are ingested. The deuterated water approach applied in this study allowed the assessment of muscle protein synthesis rates during 7 d of the exercise training intervention. During this week, other factors such as regular food intake, habitual physical activity, stress, and sleep may have modulated muscle protein synthesis rates. The present findings indicate that protein supplementation does not further augment muscle protein accretion during the latter stages (week 12) of a 12-wk exercise training program in active older men.

Dietary protein intake during the intervention period was 1.2 ± 0.1 g/kg body mass/d in the placebo group and 1.4 ± 0.1 g \cdot kg body mass⁻¹ \cdot d⁻¹ in the protein-supplemented group. Instead of merely increasing dietary protein intake, the present study focused on timed protein supplementation, with supplements being provided immediately after exercise and before sleep. The ingestion of dietary protein in close temporal proximity to exercise has been shown to further improve postexercise net muscle protein balance (56–59). In addition, we have previously shown that the ingestion of 40 g casein before sleep increases overnight muscle protein synthesis rates (36, 37, 40) and could therefore serve to promote skeletal muscle mass gains when combined with resistance exercise training (38, 60).

TABLE 6 Muscle strength and physical performance before and after 12 wk of resistance exercise training in active older men who did or did not receive protein supplementation¹

	Placebo ($n = 19$)			Protein ($n = 21$)			<i>P</i>		
	0 wk	12 wk	Difference	0 wk	12 wk	Difference	Treatment \times time interaction	Treatment effect	Time effect
Leg press 1RM, kg	162 \pm 6	188 \pm 7	+25 \pm 3	157 \pm 5	180 \pm 6	+24 \pm 3	0.765	0.397	<0.001
Leg extension 1RM, kg	88 \pm 3	105 \pm 4	+16 \pm 3	85 \pm 3	102 \pm 4	+17 \pm 3	0.798	0.481	<0.001
Total 1RM, kg	250 \pm 7	292 \pm 9	+41 \pm 5	241 \pm 7	282 \pm 8	+41 \pm 5	0.979	0.359	<0.001
SPPB score, points	11.2 \pm 0.2	11.6 \pm 0.1	+0.4 \pm 0.2	11.3 \pm 0.1	11.5 \pm 0.2	+0.2 \pm 0.2	0.340	0.719	0.006
4-m Walking test, s	3.57 \pm 0.12	3.60 \pm 0.09	+0.03 \pm 0.11	3.70 \pm 0.08	3.72 \pm 0.11	+0.03 \pm 0.09	0.877	0.273	0.604
Sit-to-stand test time, s	12.03 \pm 0.31	11.20 \pm 0.32	-0.83 \pm 0.37	11.57 \pm 0.27	10.80 \pm 0.27	-0.77 \pm 0.25	0.739	0.233	<0.001

¹Values are means \pm SEMs. SPPB, short physical performance battery; 1RM, 1-repetition maximum.




FIGURE 5 Plasma glucose (A) and insulin (C) concentrations and AUCs (B, D) in response to an OGTT before and after 12 wk of resistance exercise training in active older men who did or did not receive protein supplementation. Values are means \pm SEMs, $n = 20$ (placebo) or 21 (protein). Data for AUCs were analyzed with the use of 2-factor ANOVA. *Different from values before exercise training, $P < 0.05$. OGTT, oral-glucose-tolerance test.

In support, we have previously shown that presleep protein supplementation (~ 30 g/d) augments gains in muscle mass and strength after 12 wk of resistance exercise training in healthy young men (41). In the present study, however, we failed to observe greater increases in skeletal muscle mass or strength after postexercise and presleep protein supplementation during 12 wk of resistance exercise training in active older men. The absence of any surplus benefits of protein supplementation on muscle mass and strength gains during prolonged resistance exercise training in older compared with younger adults may be explained by age-related factors, which include a lower absolute workload, lower habitual physical activity, and the prevalence of anabolic resistance. In support, exercise workload and absolute gains in muscle mass and strength tend to be greater in younger individuals than in older individuals (7, 8, 41). Consequently, protein supplementation appears to provide a greater benefit to the skeletal muscle adaptive response to resistance exercise training in younger individuals (3, 4, 6–8).

It has been suggested that the prevalence of anabolic resistance may be compensated for by ingesting even larger

amounts (i.e., 40 g) of protein than what was provided in the present study (20 g whey + 1 g leucine). However, previous work has shown that the ingestion of 20 g whey should be effective at increasing postexercise muscle protein synthesis rates in the healthy older population (26). Furthermore, numerous studies have shown leucine coingestion to further augment the muscle protein synthetic response to protein ingestion (32–35). Therefore, we can only speculate on whether supplementation with a larger protein dose could have further increased muscle mass and strength gains after resistance training in these healthy, active older men.

The present findings add to the growing number of studies from our group (13, 18) as well as others (10–12, 14–17) that show that protein supplementation does not further augment skeletal muscle mass or strength gains during resistance exercise training in active older men who habitually consume ample amounts of dietary protein (1.0–1.2 g/kg body mass/d). In contrast, protein supplementation has been reported to increase muscle mass and strength gains during resistance exercise training in prefrail older men and women who consume less than optimal amounts of dietary protein (5). Therefore, protein

TABLE 7 Fasting serum hormone concentrations before and after 12 wk of resistance exercise training in active older men who did or did not receive protein supplementation¹

	Placebo ($n = 20$)			Protein ($n = 21$)			P		
	0 wk	12 wk	Difference	0 wk	12 wk	Difference	Treatment \times time interaction	Treatment effect	Time effect
Testosterone, nmol/L	17.7 \pm 1.8	18.5 \pm 1.8	+0.8 \pm 1.0	16.0 \pm 1.2	16.8 \pm 1.4	+0.7 \pm 0.9	0.972	0.534	0.258
Free testosterone, pmol/L	247 \pm 18	251 \pm 15	+5 \pm 13	242 \pm 15	262 \pm 19	+17 \pm 14	0.509	0.867	0.258
Growth hormone, μ g/L	3.6 \pm 1.0	4.3 \pm 1.0	+0.7 \pm 1.0	4.3 \pm 1.3	2.6 \pm 0.8	-1.6 \pm 1.1	0.171	0.974	0.583
IGF-I, nmol/L	16.6 \pm 1.4	17.6 \pm 1.3	+0.9 \pm 0.6	18.2 \pm 0.8	17.9 \pm 0.9	-0.1 \pm 0.5	0.163	0.733	0.399
SHBG, nmol/L	50.3 \pm 3.9	51.9 \pm 3.9	+1.5 \pm 1.2	44.3 \pm 3.6	43.1 \pm 3.5	-1.2 \pm 1.0	0.062	0.284	0.629

¹Values are means \pm SEMs. IGF-I, insulin-like growth factor I; SHBG, sex hormone-binding globulin.

supplementation may be of greater benefit in more clinically compromised older populations when gains in muscle mass, strength, or both are restricted by habitual protein consumption (61). In conclusion, protein supplementation after exercise and before sleep does not augment skeletal muscle mass or strength gains after resistance exercise training in active older men who consume ample dietary protein.

Acknowledgments

We thank Jean Nyakayiru, Tobias Lampert, and Finn Frielink for assistance during the experimental test days. The authors' responsibilities were as follows—AMH, LBV, and LJCvL: designed the research; AMH, KJMP, MO, JSJS, and EMPB: conducted the research; AMH, JvK, APG, and JPBG: performed sample analyses; AMH: analyzed the data; AMH and LBV: performed the statistical analysis; AMH and LJCvL: wrote the manuscript and hold primary responsibility for the final content; and all authors: read and approved the final manuscript.

References

- Baumgartner RN, Waters DL, Gallagher D, Morley JE, Garry PJ. Predictors of skeletal muscle mass in elderly men and women. *Mech Ageing Dev* 1999;107:123–36.
- Mitchell WK, Williams J, Atherton P, Larvin M, Lund J, Narici M. Sarcopenia, dynapenia, and the impact of advancing age on human skeletal muscle size and strength; a quantitative review. *Front Physiol* 2012;3:260.
- Cermak NM, Res PT, de Groot LCPGM, Saris WHM, van Loon LJC. Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: a meta-analysis. *Am J Clin Nutr* 2012;96:1454–64.
- Morton RW, Murphy KT, McKellar SR, Schoenfeld BJ, Henselmans M, Helms E, Aragon AA, Devries MC, Banfield L, Krieger JW, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med* 2018;52(6):376–84.
- Tieland M, Dirks ML, van der Zwaluw N, Verdijk LB, van de Rest O, de Groot LCPGM, van Loon LJC. Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 2012;13:713–9.
- Walker TB, Smith J, Herrera M, Lebegue B, Pinchak A, Fischer J. The influence of 8 weeks of whey-protein and leucine supplementation on physical and cognitive performance. *Int J Sport Nutr Exerc Metab* 2010;20:409–17.
- Hartman JW, Tang JE, Wilkinson SB, Tarnopolsky MA, Lawrence RL, Fullerton AV, Phillips SM. Consumption of fat-free fluid milk after resistance exercise promotes greater lean mass accretion than does consumption of soy or carbohydrate in young, novice, male weightlifters. *Am J Clin Nutr* 2007;86:373–81.
- Kerksick CM, Rasmussen CJ, Lancaster SL, Magu B, Smith P, Melton C, Greenwood M, Almada AL, Earnest CP, Kreider RB. The effects of protein and amino acid supplementation on performance and training adaptations during ten weeks of resistance training. *J Strength Cond Res* 2006;20:643–53.
- Bemben MG, Witten MS, Carter JM, Eliot KA, Knehans AW, Bemben DA. The effects of supplementation with creatine and protein on muscle strength following a traditional resistance training program in middle-aged and older men. *J Nutr Health Aging* 2010;14:155–9.
- Campbell WW, Crim MC, Young VR, Joseph LJ, Evans WJ. Effects of resistance training and dietary protein intake on protein metabolism in older adults. *Am J Physiol* 1995;268:E1143–53.
- Holm L, Olesen JL, Matsumoto K, Doi T, Mizuno M, Alsted TJ, Mackey AL, Schwarz P, Kjaer M. Protein-containing nutrient supplementation following strength training enhances the effect on muscle mass, strength, and bone formation in postmenopausal women. *J Appl Physiol* 2008;105:274–81.
- Kukuljan S, Nowson CA, Sanders K, Daly RM. Effects of resistance exercise and fortified milk on skeletal muscle mass, muscle size, and functional performance in middle-aged and older men: an 18-mo randomized controlled trial. *J Appl Physiol* 2009;107:1864–73.
- Verdijk LB, Jonkers RA, Gleeson BG, Beelen M, Meijer K, Savelberg HH, Wodzig WK, Dendale P, van Loon LJ. Protein supplementation before and after exercise does not further augment skeletal muscle hypertrophy after resistance training in elderly men. *Am J Clin Nutr* 2009;89:608–16.
- Iglay HB, Thyfault JP, Apolzan JW, Campbell WW. Resistance training and dietary protein: effects on glucose tolerance and contents of skeletal muscle insulin signaling proteins in older persons. *Am J Clin Nutr* 2007;85:1005–13.
- Meredith CN, Frontera WR, O'Reilly KP, Evans WJ. Body composition in elderly men: effect of dietary modification during strength training. *J Am Geriatr Soc* 1992;40:155–62.
- Candow DG, Chilibeck PD, Facci M, Abeysekara S, Zello GA. Protein supplementation before and after resistance training in older men. *Eur J Appl Physiol* 2006;97:548–56.
- Godard MP, Williamson DL, Trappe SW. Oral amino-acid provision does not affect muscle strength or size gains in older men. *Med Sci Sports Exerc* 2002;34:1126–31.
- Leenders M, Verdijk LB, van der Hoeven L, van Kranenburg J, Nilwik R, Wodzig WKWH, Senden JMG, Keizer HA, van Loon LJC. Protein supplementation during resistance-type exercise training in the elderly. *Med Sci Sports Exerc* 2013;45:542–52.
- Koopman R, Crombach N, Gijsen AP, Walrand S, Fauquant J, Kies AK, Lemosquet S, Saris WHM, Boirie Y, van Loon LJC. Ingestion of a protein hydrolysate is accompanied by an accelerated in vivo digestion and absorption rate when compared with its intact protein. *Am J Clin Nutr* 2009;90:106–15.
- Pennings B, Boirie Y, Senden JMG, Gijsen AP, Kuipers H, van Loon LJC. Whey protein stimulates postprandial muscle protein accretion more effectively than do casein and casein hydrolysate in older men. *Am J Clin Nutr* 2011;93:997–1005.
- Paddon-Jones D, Sheffield-Moore M, Katsanos CS, Zhang X-J, Wolfe RR. Differential stimulation of muscle protein synthesis in elderly humans following isocaloric ingestion of amino acids or whey protein. *Exp Gerontol* 2006;41:215–9.
- Gorissen SH, Horstman AM, Franssen R, Crombag JJ, Langer H, Bierau J, Respondek F, van Loon LJ. Ingestion of wheat protein increases in vivo muscle protein synthesis rates in healthy older men in a randomized trial. *J Nutr* 2016;146:1651–9.
- Yang Y, Churchward-Venne TA, Burd NA, Breen L, Tarnopolsky MA, Phillips SM. Myofibrillar protein synthesis following ingestion of soy protein isolate at rest and after resistance exercise in elderly men. *Nutr Metab* 2012;9:57.
- Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, Phillips SM. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J Appl Physiol* 2009;107:987–92.
- Pennings B, Groen B, de Lange A, Gijsen AP, Zorenc AH, Senden JMG, van Loon LJC. Amino acid absorption and subsequent muscle protein accretion following graded intakes of whey protein in elderly men. *Am J Physiol Endocrinol Metab* 2012;302:E992–9.
- Yang Y, Breen L, Burd NA, Hector AJ, Churchward-Venne TA, Josse AR, Tarnopolsky MA, Phillips SM. Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br J Nutr* 2012;108:1780–8.
- Moore DR, Robinson MJ, Fry JL, Tang JE, Glover EI, Wilkinson SB, Prior T, Tarnopolsky MA, Phillips SM. Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr* 2009;89:161–8.
- Witard OC, Jackman SR, Breen L, Smith K, Selby A, Tipton KD. Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr* 2014;99:86–95.
- Mamerow MM, Mettler JA, English KL, Casperson SL, Arentson-Lantz E, Sheffield-Moore M, Layman DK, Paddon-Jones D. Dietary protein distribution positively influences 24-h muscle protein synthesis in healthy adults. *J Nutr* 2014;144:876–80.
- Areta JL, Burke LM, Ross ML, Camera DM, West DWD, Broad EM, Jeacocke NA, Moore DR, Stellingwerff T, Phillips SM, et al. Timing and distribution of protein ingestion during prolonged recovery

- from resistance exercise alters myofibrillar protein synthesis. *J Physiol* 2013;591:2319–31.
31. West DW, Burd NA, Coffey VG, Baker SK, Burke LM, Hawley JA, Moore DR, Stellingwerf T, Phillips SM. Rapid aminoacidemia enhances myofibrillar protein synthesis and anabolic intramuscular signaling responses after resistance exercise. *Am J Clin Nutr* 2011;94:795–803.
 32. Atherton PJ, Kumar V, Selby AL, Rankin D, Hildebrandt W, Phillips BE, Williams JP, Hiscock N, Smith K. Enriching a protein drink with leucine augments muscle protein synthesis after resistance exercise in young and older men. *Clin Nutr* 2017;36:888–95.
 33. Bukhari SSI, Phillips BE, Wilkinson DJ, Limb MC, Rankin D, Mitchell WK, Kobayashi H, Greenhaff PL, Smith K, Atherton PJ. Intake of low-dose leucine-rich essential amino acids stimulates muscle anabolism equivalently to bolus whey protein in older women at rest and after exercise. *Am J Physiol Endocrinol Metab* 2015;308:E1056–65.
 34. Katsanos CS, Kobayashi H, Sheffield-Moore M, Aarsland A, Wolfe RR. A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly. *Am J Physiol Endocrinol Metab* 2006;291:E381–7.
 35. Wall BT, Hamer HM, de Lange A, Kiskini A, Groen BBL, Senden JMG, Gijsen AP, Verdijk LB, van Loon LJC. Leucine co-ingestion improves post-prandial muscle protein accretion in elderly men. *Clin Nutr* 2013;32:412–9.
 36. Groen BBL, Res PT, Pennings B, Hertle E, Senden JMG, Saris WHM, van Loon LJC. Intra-gastric protein administration stimulates overnight muscle protein synthesis in elderly men. *Am J Physiol Endocrinol Metab* 2012;302:E52–60.
 37. Kouw IW, Holwerda AM, Trommelen J, Kramer IF, Bastiaanse J, Halson SL, Wodzig WK, Verdijk LB, van Loon LJ. Protein ingestion before sleep increases overnight muscle protein synthesis rates in healthy older men: a randomized controlled trial. *J Nutr* 2017;147:2252–61.
 38. Holwerda AM, Kouw IW, Trommelen J, Halson SL, Wodzig WK, Verdijk LB, van Loon LJ. Physical activity performed in the evening increases the overnight muscle protein synthetic response to presleep protein ingestion in older men. *J Nutr* 2016;146:1307–14.
 39. Trommelen J, Kouw IWK, Holwerda AM, Snijders T, Halson SL, Rollo I, Verdijk LB, van Loon LJC. Pre-sleep dietary protein-derived amino acids are incorporated in myofibrillar protein during post-exercise overnight recovery. *Am J Physiol Endocrinol Metab* 2018;314(5):E457–67.
 40. Res PT, Groen B, Pennings B, Beelen M, Wallis GA, Gijsen AP, Senden JMG, van Loon LJC. Protein ingestion before sleep improves postexercise overnight recovery. *Med Sci Sports Exerc* 2012;44:1560–9.
 41. Snijders T, Res PT, Smeets JSJ, van Vliet S, van Kranenburg J, Maase K, Kies AK, Verdijk LB, van Loon LJC. Protein ingestion before sleep increases muscle mass and strength gains during prolonged resistance-type exercise training in healthy young men. *J Nutr* 2015;145:1178–84.
 42. Backx EMP, Hangelbroek R, Snijders T, Verschuijden M-L, Verdijk LB, Groot LCPGM, Loon LJC. Creatine loading does not preserve muscle mass or strength during leg immobilization in healthy, young males: a randomized controlled trial. *Sports Med* 2017;47:1661–71.
 43. Bergström J. Percutaneous needle biopsy of skeletal muscle in physiological and clinical research. *Scand J Clin Lab Invest* 1975;35:609–16.
 44. Holwerda AM, Paulussen KJM, Overkamp M, Smeets JSJ, Gijsen AP, Goossens JPB, Verdijk LB, van Loon LJC. Daily resistance-type exercise stimulates overall muscle protein synthesis rates in vivo in young males. *J Appl Physiol* 2018;124(1):66–75.
 45. Mayhew JL, Prinster JL, Ware JS, Zimmer DL, Arabas JR, Bembem MG. Muscular endurance repetitions to predict bench press strength in men of different training levels. *J Sports Med Phys Fitness* 1995;35:108–13.
 46. Verdijk LB, Koopman R, Schaart G, Meijer K, Savelberg HHCM, van Loon LJC. Satellite cell content is specifically reduced in type II skeletal muscle fibers in the elderly. *Am J Physiol Endocrinol Metab* 2006;292:E151–7.
 47. Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, Scherr PA, Wallace RB. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol* 1994;49:M85–94.
 48. Vermeulen A, Verdonck L, Kaufman JM. A critical evaluation of simple methods for the estimation of free testosterone in serum. *J Clin Endocrinol Metab* 1999;84:3666–72.
 49. Kosek DJ. Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *J Appl Physiol* 2006;101:531–44.
 50. Martel GF, Roth SM, Ivey FM, Lemmer JT, Tracy BL, Hurlbut DE, Metter EJ, Hurley BF, Rogers MA. Age and sex affect human muscle fibre adaptations to heavy-resistance strength training. *Exp Physiol* 2006;91:457–64.
 51. Wilkinson DJ, Franchi MV, Brook MS, Narici MV, Williams JP, Mitchell WK, Szwedczyk NJ, Greenhaff PL, Atherton PJ, Smith K. A validation of the application of D2O stable isotope tracer techniques for monitoring day-to-day changes in muscle protein subfraction synthesis in humans. *Am J Physiol Endocrinol Metab* 2014;306:E571–9.
 52. Robinson MM, Turner SM, Hellerstein MK, Hamilton KL, Miller BF. Long-term synthesis rates of skeletal muscle DNA and protein are higher during aerobic training in older humans than in sedentary young subjects but are not altered by protein supplementation. *FASEB J* 2011;25:3240–9.
 53. Dickinson JM, Gundermann DM, Walker DK, Reidy PT, Borack MS, Drummond MJ, Arora M, Volpi E, Rasmussen BB. Leucine-enriched amino acid ingestion after resistance exercise prolongs myofibrillar protein synthesis and amino acid transporter expression in older men. *J Nutr* 2014;144:1694–702.
 54. Murphy CH, Saddler NI, Devries MC, McGlory C, Baker SK, Phillips SM. Leucine supplementation enhances integrative myofibrillar protein synthesis in free-living older men consuming lower- and higher-protein diets: a parallel-group crossover study. *Am J Clin Nutr* 2016;104:1594–606.
 55. Kramer IF, Verdijk LB, Hamer HM, Verlaan S, Luiking YC, Kouw IWK, Senden JM, van Kranenburg J, Gijsen AP, Bierau J, et al. Both basal and post-prandial muscle protein synthesis rates, following the ingestion of a leucine-enriched whey protein supplement, are not impaired in sarcopenic older males. *Clinical Nutrition* 2017;36:1440–9.
 56. Tipton KD, Rasmussen BB, Miller SL, Wolf SE, Owens-Stovall SK, Petrini BE, Wolfe RR. Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrinol Metab* 2001;281:E197–206.
 57. Beelen M, Zorenc A, Pennings B, Senden JM, Kuipers H, van Loon LJC. Impact of protein co-ingestion on muscle protein synthesis during continuous endurance type exercise. *Am J Physiol Endocrinol Metab* 2011;300:E945–54.
 58. Tipton KD, Elliott TA, Cree MG, Aarsland AA, Sanford AP, Wolfe RR. Stimulation of net muscle protein synthesis by whey protein ingestion before and after exercise. *Am J Physiol Endocrinol Metab* 2006;292:E71–6.
 59. Rasmussen BB, Tipton KD, Miller SL, Wolf SE, Wolfe RR. An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J Appl Physiol* 2000;88:386–92.
 60. Trommelen J, Holwerda AM, Kouw IWK, Langer H, Halson SL, Rollo I, Verdijk LB, van Loon LJC. Resistance exercise augments postprandial overnight muscle protein synthesis rates. *Med Sci Sports Exerc* 2016;48:2517–25.
 61. Bauer JM, Verlaan S, Bautmans I, Brandt K, Donini LM, Maggio M, McMurdo MET, Mets T, Seal C, Wijers SL, et al. Effects of a vitamin D and leucine-enriched whey protein nutritional supplement on measures of sarcopenia in older adults: the PROVIDE study: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 2015;16:740–7.