

Beta2-adrenergic receptor polymorphisms and salbutamol-stimulated energy expenditure

Citation for published version (APA):

Oomen, J. M., van Rossum, C. T., Hoebee, B., Saris, W. H., & van Baak, M. A. (2005). Beta2-adrenergic receptor polymorphisms and salbutamol-stimulated energy expenditure. *Journal of Clinical Endocrinology & Metabolism*, 90(4), 2301-2307. <https://doi.org/10.1210/jc.2004-1356>

Document status and date:

Published: 01/01/2005

DOI:

[10.1210/jc.2004-1356](https://doi.org/10.1210/jc.2004-1356)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

β_2 -Adrenergic Receptor Polymorphisms and Salbutamol-Stimulated Energy Expenditure

J. M. Oomen, C. T. M. van Rossum, B. Hoebee, W. H. M. Saris, and M. A. van Baak

Department of Human Biology/NUTRIM (J.M.O., W.H.M.S., M.A.v.B.), Maastricht University, 6200 MD Maastricht, The Netherlands; and Department of Chronic Diseases Epidemiology (C.T.M.v.R.) and Laboratory of Health Effects Research (B.H.), National Institute of Public Health and the Environment, 3720 BA Bilthoven, The Netherlands

The β -adrenergic system is involved in the control of energy metabolism and expenditure. The β_2 -adrenergic receptor (β_2 -AR) gene shows polymorphisms that have been associated with obesity in several studies. *In vitro* and *in vivo* studies suggest differences in β_2 -AR-mediated function between these polymorphisms. The aim of this study was to investigate the influence of genetic variation in codon 16 of the β_2 -AR gene on energy metabolism in humans.

Thirty-four subjects were recruited [Gly16Gly (n = 13), Gly16Arg (n = 16), or Arg16Arg (n = 5)]. The β_2 -AR was stimulated with two doses of salbutamol (50 and 100 ng/kg fat-free mass per minute) after blockade of the β_1 -adrenergic receptors with atenolol. Energy expenditure and plasma substrate and hormone concentrations were measured.

The increase in energy expenditure (Δ EE) was significantly

different among groups in which the Arg16Arg group showed the lowest increase ($P < 0.05$ vs. Gly carriers). In a multiple regression model, variations in the increase in nonesterified fatty acid concentration during salbutamol infusion (partial $r = 0.51$) and the polymorphism contributed significantly to the variation in Δ EE. Thirty-five percent of the variation in Δ EE was explained by these two factors.

We conclude that subjects with the Arg16Arg polymorphism of the β_2 -AR gene have a reduced thermogenic response to β_2 -adrenergic stimulation. Although this relatively small study needs confirmation, the findings support a role for this polymorphism in the development and maintenance of overweight and obesity. (*J Clin Endocrinol Metab* 90: 2301–2307, 2005)

IT IS WELL DOCUMENTED that both environmental and genetic factors are involved in the onset and progression of obesity in humans. Severe obesity appears to have a particularly strong genetic component and is polygenic in nature (1). An increasing number of polymorphisms is assumed to be associated with obesity and has been described in literature (1). Despite intense effort, the pathways underlying these associations with obesity remain elusive. This is largely due to the complexities circumventing the process of developing obesity, which include age of onset; polygenic inheritance; genetic heterogeneity; incomplete penetrance; unknown mode of action of disease alleles; effect of ethnicity, age, gender, and the interaction with environmental factors, such as diet, physical activity, or smoking status.

Of all contributing factors, the components of the sympathetic nervous system are of interest because this system, in particular its β -adrenergic component, is involved in the control of energy metabolism and expenditure (2, 3).

One of the receptor subtypes of the β -adrenergic system is the β_2 -adrenergic receptor (β_2 -AR). It is coupled to a stimulatory G protein that promotes cAMP production, activating protein kinase A, which mediates a variety of responses, depending on the cell type. For instance, the lipolytic and gly-

cogenolytic effects of catecholamines are mediated through members of the β_2 -AR family (2, 4). One study found that the β_2 -mediated thermogenic and lipolytic responses are blunted in obese, compared with lean (5). In addition, several studies (6–13) have shown an association between β_2 -AR polymorphisms and weight gain, obesity, or obesity-related phenotypes. Nine polymorphisms have been identified in the coding region of the human β_2 -AR, three of which are nonsynonymous (14). The two most common variants are located at codon 16 (Arg16Gly) and codon 27 (Gln27Glu) and have been studied most in relation to obesity. Both polymorphisms result in variations at the amino-terminal site of the receptor.

In vitro studies have shown differences in agonist-promoted down-regulation (15) but no differences in agonist binding or agonist-stimulated adenylyl cyclase activities among these polymorphisms of the β_2 -AR gene (16). Functional consequences of these polymorphisms with respect to adipocyte lipolysis have been reported (17, 18), which may play a role in obesity. So far, the role of the β_2 -AR polymorphisms in human energy expenditure (EE) has not been studied.

We investigated whether the polymorphism at codon 16 of the β_2 -AR gene is of functional importance in human energy metabolism by direct stimulation of this receptor with a β_2 -AR agonist in an experimental setting.

Subjects and Methods

Subjects

Thirty-four healthy, normal-weight to moderately obese volunteers [body mass index (BMI) range 19.1–37.1 kg/m²] participated in this

First Published Online February 1, 2005

Abbreviations: Δ , Change; β_2 -AR, β_2 -adrenergic receptor; BMI, body mass index; BP, blood pressure; EE, energy expenditure; FFM, fat-free mass; HOMA, homeostasis model assessment; NEFA, nonesterified fatty acid; RER, respiratory exchange ratio.

JCEM is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.

study (32 men and two women, aged 45 ± 0.8 yr, BMI 27.8 ± 0.8 kg/m²). In part (n = 14) they were recruited from an existing cohort that has been described previously (13), in which the polymorphisms in the β_2 -AR were known; in part they were newly recruited by advertisement (n = 20). The subjects did not use medication. Physical characteristics of the subjects are summarized in Table 1.

Experimental design

One week before the experimental day, subjects came to the laboratory after an overnight fast, and body composition was determined by hydrostatic weighing with simultaneous lung volume measurements (Volugraph 2000, Mijndhardt, Bunnik, The Netherlands). Body composition was calculated according to the equation of Siri (19).

Subjects were asked to refrain from strenuous exercise during 24 h before the experiment. After an overnight fast, subjects arrived at the laboratory by bus or car to minimize activity before the measurements. On arrival, a cannula was inserted into a forearm vein of each arm. One cannula was used for the infusion of drugs and the other cannula for the sampling of blood. Next, indirect calorimetry measurements were started with a ventilated hood system with the subject in recumbent position and continued for the remainder of the experiment.

The study consisted of three study periods of 45 min. Before the first period, subjects received a priming dose of 50 μ g/kg fat-free mass (FFM) atenolol (β_1 -adrenergic receptor antagonist) (Tenormin, Zeneca, Ridderkerk, The Netherlands) in 5 min, after which a continuous infusion of 1.2 μ g/kg FFM per minute atenolol was started for the remainder of the experiment. After 45 min atenolol infusion, subjects additionally received consecutive infusions of 50 and 100 ng/kg FFM per minute salbutamol (Ventolin, GlaxoWellcome, Zeist, The Netherlands). Each infusion period lasted 45 min. After 30 and 45 min of each study period, a blood sample was taken.

Measurements

Whole-body EE and respiratory exchange ratio (RER) were measured by indirect calorimetry, using an open-circuit ventilated hood system. In the study a homemade system, which is based on the analysis system for respiration chambers, which has been described previously (20), was used. The volume of air drawn through the hood was measured by a dry-gas meter (Schlumberger, Dordrecht, The Netherlands), and the composition of the in- and outflowing air was analyzed by a paramagnetic O₂ analyzer (Servomex, Crowborough, UK) and an infrared CO₂ analyzer (Hartmann and Braun, Frankfurt, Germany). The airflow rate and the O₂ and CO₂ concentrations of the in- and outflowing air were used to compute O₂ consumption and CO₂ production online through an automatic acquisition system connected to a personal computer. EE was calculated according to the formula proposed by Weir (21). EE and RER values were averaged over the last 15 min of each period. The reproducibility of the measurement was within 5% as described before (22).

TABLE 1. Subject characteristics of groups with different polymorphisms at codon 16 of the β_2 -adrenergic receptor gene (mean \pm SEM)

	Gly16Gly (n = 13)	Gly16Arg (n = 16)	Arg16Arg (n = 5)
Gender (M/F)	12/1	16/0	4/1
Weight (kg)	83.6 \pm 4.3	87.0 \pm 3.5	86.5 \pm 12.9
Height (m)	1.73 \pm 0.03	1.78 \pm 0.02	1.74 \pm 0.05
Age (yr)	46.9 \pm 1.4	45.5 \pm 1.3	41.5 \pm 1.1
Fat percentage (%)	28.0 \pm 2.6	27.4 \pm 1.8	35.0 \pm 1.1
Fat mass (kg)	23.9 \pm 2.7	24.6 \pm 2.4	30.7 \pm 5.1
FFM (kg)	59.6 \pm 3.0	62.4 \pm 1.7	55.9 \pm 7.8
BMI (kg/m ²)	27.8 \pm 1.2	27.7 \pm 1.3	28.1 \pm 3.0
HOMA index	2.25 \pm 0.52	2.15 \pm 0.33	4.90 \pm 1.84 ^{a,b}

^a Significantly different between Arg16Arg and Gly16Arg ($P < 0.05$).

^b Tendency for difference between Arg16Arg and Gly16Gly ($P = 0.06$).

Total carbohydrate and fat oxidation were calculated using stoichiometric equations (23):

$$\text{Total fat oxidation (grams per minute)} = 1.695 \text{ VO}_2 \text{ (liters per minute)} - 1.701 \text{ VCO}_2 \text{ (liters per minute)}.$$

$$\text{Total carbohydrate oxidation (grams per minute)} = 4.585 \text{ VCO}_2 \text{ (liters per minute)} - 3.226 \text{ VO}_2 \text{ (liters per minute)}.$$

Substrate oxidation was expressed as percentage of the contribution to total EE (1 g fat = 37.8 kJ and 1 g carbohydrate = 16.9 kJ).

Heart rate was monitored by conventional electrocardiography. Blood pressure was measured three times by an automated blood pressure device (OMRON 705CP, Hamburg, Germany) during the last 5 min of each study period. The means of these three measurements were used for further analysis.

Analytical methods

Genomic DNA of the subjects was extracted from leukocytes by digestion with proteinase K followed by phenol/chloroform extraction. Determination of the polymorphism was performed using a PCR-restriction fragment length polymorphism analysis as described before (18).

Blood samples for the determination of nonesterified fatty acids (NEFAs), glycerol, glucose, and insulin were preserved with sodium-EDTA and those for norepinephrine and epinephrine with heparin plus glutathione (1.5% wt/vol). Blood samples were immediately centrifuged for 10 min at $800 \times g$ at 4 C. Plasma was rapidly frozen in liquid nitrogen and stored at -70 C until further analysis. Plasma NEFA concentration was measured with the NEFA C kit (99475409, Wako, Neuss, Germany), plasma glycerol concentration was measured with a glycerol kit (148270, Boehringer, Mannheim, Germany), and plasma glucose concentration was measured with a glucose kit (UniKit III, 07367204, Roche, Basel, Switzerland), all on a Cobas Fara centrifugal analyzer (Roche Diagnostics, Basel, Switzerland). Plasma insulin level was determined with a double-antibody RIA (Insulin RIA 100, Pharmacia, Uppsala, Sweden). The homeostasis model assessment (HOMA) index was calculated according to Matthews *et al.* (24) using baseline plasma glucose and baseline plasma insulin levels. Plasma norepinephrine and epinephrine levels were determined by HPLC with electrochemical detection according to the method of Alberts *et al.* (25) using a ClinPrep kit (Recipe, Munich, Germany).

Data analysis

All data are presented as mean \pm SEM. Data for EE were adjusted for FFM for group comparison (26). The change (Δ) in EE induced by the highest dose of salbutamol (100 ng/kg FFM per minute) was calculated by subtracting baseline EE from the EE at 100 ng/kg FFM per minute salbutamol. Changes in the other variables were calculated the same way.

One-way ANOVA with repeated measurements was used to analyze the effects of salbutamol administration in the total group of subjects and the three polymorphism groups. Differences among the codon 16 polymorphism groups at baseline and in response to salbutamol administration were analyzed by one-way ANOVA. *Post hoc* pairwise comparisons were made using Bonferroni correction.

Simple regression analysis was performed with Δ EE as the dependent variable and different parameters as independent variables (see Table 3). Multiple regression analysis was conducted to estimate the independent contributions of the variables associated with Δ EE. All variables that were correlated with Δ EE in the simple regression analysis with a $P < 0.20$ were included in the analysis. In addition, the polymorphism groups were entered as dummy variables into the multiple regression model. $P < 0.05$ was considered to be statistically significant.

Results

Responses to salbutamol infusion in the whole group

Table 2 shows changes in all parameters studied at baseline and during administration of increasing doses of salbutamol for the whole group of subjects. EE, heart rate, plasma norepinephrine, and NEFA and glycerol concentrations

TABLE 2. Metabolic parameters at different doses of salbutamol for the whole group (n = 34)

Salbutamol (ng/kg FFM · min)	0	50	100	P value (ANOVA)
Energy expenditure (kJ/min)	4.8 ± 0.1	5.1 ± 0.1	5.3 ± 0.1	0.037
RER	0.83 ± 0.01	0.85 ± 0.01	0.82 ± 0.01	0.183
Fat oxidation (% of total energy expenditure)	53.5 ± 2.2	48.5 ± 2.8	55.6 ± 3.2	0.179
Carbohydrate oxidation (% of total energy expenditure)	47.9 ± 2.8	53.6 ± 3.2	45.7 ± 3.5	0.181
Heart rate (beats/min)	55.9 ± 1.2	62.8 ± 1.2	71.2 ± 1.4	<0.001
Diastolic blood pressure (mm Hg)	83.9 ± 1.8	81.7 ± 1.7	78.2 ± 1.7	0.075
Systolic blood pressure (mm Hg)	118.1 ± 2.0	118.9 ± 1.9	121.1 ± 1.9	0.532
[Norepinephrine] (ng/liter)	456.0 ± 20.6	513.3 ± 33.2	579 ± 32.7	0.014
[Epinephrine] (ng/liter)	33.9 ± 3.5	24.1 ± 2.5	21.8 ± 2.1	0.006
[NEFA] (μ mol/liter)	366.6 ± 21.1	580.2 ± 30.8	724.9 ± 42.1	<0.001
[Glycerol] (μ mol/liter)	58.5 ± 3.7	84.3 ± 5.2	106.8 ± 7.2	<0.001
[Glucose] (mmol/liter)	5.40 ± 0.12	5.36 ± 0.10	5.49 ± 0.10	0.656
[Lactate] (mmol/liter)	0.97 ± 0.06	1.00 ± 0.05	1.15 ± 0.05	0.055
[Insulin] (mU/liter)	10.1 ± 1.3	13.5 ± 1.8	15.2 ± 1.6	0.075

Data represent mean \pm SEM.

showed statistically significant increases (at least $P < 0.05$). Plasma epinephrine concentrations were significantly lowered during salbutamol infusion ($P < 0.01$). No statistically significant changes in RER, blood pressure (BP), plasma glucose, lactate, and insulin were found.

Baseline values and responses to salbutamol infusion in the β_2 -AR codon 16 polymorphism groups

Baseline EE adjusted for FFM was similar in all polymorphism groups (4.89 ± 0.14 , 4.89 ± 0.06 , and 4.92 ± 0.23 kJ/min for Gly16Gly, Gly16Arg, and Arg16Arg, respectively, NS). There was a significant difference in the increase in EE (Δ EE) among groups (ANOVA, $P < 0.05$) (Fig. 1). *Post hoc* analysis showed that the response was significantly different between the Arg16Arg and the Gly16Arg groups ($P < 0.05$) and between Arg16Arg and Gly carriers ($P < 0.05$).

At baseline RER did not differ among groups, nor did Δ RER ($P = 0.41$) (Fig. 2). Fat oxidation at baseline was not significantly different among groups (54.4 ± 3.4 , 55.7 ± 3.2 , and $44.6 \pm 6.3\%$ of total EE for Gly16Gly, Gly16Arg, and Arg16Arg, respectively, $P = 0.25$) nor was carbohydrate oxidation (47.0 ± 3.9 , 45.6 ± 3.9 , and $57.9 \pm 7.0\%$ of total EE for

Gly16Gly, Gly16Arg, and Arg16Arg, respectively, $P = 0.25$). There was no difference in change of fat and carbohydrate oxidation (as percentage of total EE) in response to salbutamol among groups (ANOVA, both $P = 0.43$).

Plasma concentrations of NEFAs, glycerol, glucose, insulin, epinephrine, and norepinephrine did not differ significantly among groups at baseline nor during salbutamol infusion. The HOMA index, however, was significantly different among groups (ANOVA, $P < 0.05$), in which the Arg16Arg group showed the highest HOMA index.

During salbutamol infusion, plasma NEFA levels (Fig. 3) and norepinephrine levels (Fig. 4) increased significantly in all polymorphism groups ($P < 0.05$), with no significant difference among groups. Plasma concentrations of glycerol, glucose, insulin, and epinephrine changed significantly in the Gly16Gly and Gly16Arg groups ($P < 0.01$) but not in the Arg16Arg group; however, the responses to salbutamol infusion did not differ among groups. Plasma lactate levels were significantly higher at baseline and during salbutamol in the Arg16Arg group, compared with the Gly carriers (ANOVA, $P < 0.05$), but the lactate response to salbutamol infusion did not differ among groups (Fig. 3).




FIG. 1. EE adjusted for FFM for the codon 16 polymorphisms in the β_2 -AR at baseline and during infusion of salbutamol. *, $P < 0.05$ vs. baseline; #, $P < 0.05$ Arg16Arg vs. Gly carriers.




FIG. 2. RER for the codon 16 polymorphisms in the β_2 -AR at baseline and during infusion of salbutamol. #, $P < 0.05$ Arg16Arg vs. Gly carriers.

Baseline values for heart rate and systolic and diastolic BP were not significantly different among polymorphism groups. Systolic BP increased in the Gly16Gly and Arg16Arg groups ($P < 0.05$) but not in the Gly16Arg group ($P = 0.54$). Diastolic BP significantly decreased in the Gly16Gly and Gly16Arg groups ($P < 0.05$), but the decrease did not reach statistical significance in the Arg16Arg group ($P = 0.08$). However, the changes in heart rate and systolic and diastolic BP during salbutamol infusion did not differ significantly among groups (data not shown).

Factors contributing to the EE response to salbutamol infusion

Simple regression analysis with ΔEE as the dependent variable showed statistically significant correlations with basal NEFA and insulin concentration, $\Delta NEFA$, Δ glycerol, HOMA index, and ΔRER (Table 3). Multiple stepwise regression analysis with ΔEE as the dependent variable and basal plasma NEFA and insulin concentration, $\Delta NEFA$, HOMA index, ΔRER , age, and codon 16 polymorphism as independent variables was performed subsequently. Because glycerol and NEFAs are both parameters of lipolysis and are highly correlated ($R^2 = 0.838$), only basal NEFAs and $\Delta NEFAs$ and not basal glycerol and Δ glycerol were used in the multiple regression analysis. The analysis indicated statistically significant associations of $\Delta NEFAs$ and the codon 16 polymorphism with ΔEE . No significant association with the other variables was found. The partial correlation coefficient for $\Delta NEFAs$ was 0.51. The polymorphism added another 10% to the explained variance in ΔEE . The whole model therefore explained 35% of the variation in ΔEE (adjusted $R^2 = 0.354$, $P < 0.001$). Introducing basal and Δ glycerol instead of basal and $\Delta NEFAs$ into the model gave similar results, $\Delta NEFAs$ being replaced by Δ glycerol.

Discussion

The aim of the present study was to examine the influence of polymorphisms in codon 16 of the β_2 -AR gene on EE and

substrate oxidation during β_2 -AR stimulation by infusing the β_2 -AR agonist salbutamol. During salbutamol infusion subjects with the Arg16Arg variant of the β_2 -AR gene showed a blunted increase in EE, and their plasma lactate levels were higher. Fat oxidation as percentage of total EE and plasma NEFA and glycerol levels did not differ among groups, nor was there a difference in response to salbutamol. Multiple regression analysis showed that the polymorphism in codon 16 of the β_2 -AR gene and plasma NEFA or glycerol change but not RER change, percent body fat, BMI, age, and HOMA index were significantly associated with the change in EE.

The combined infusion of salbutamol and atenolol selectively stimulates the β_2 -AR, as shown previously (27). This selective stimulation resulted in the expected increases in EE, heart rate, plasma glycerol, NEFAs, lactate, insulin, and norepinephrine concentrations and in reductions in plasma epinephrine levels and diastolic BP (27, 28). There was no clear change of RER during salbutamol infusion in this study. However, changes in RER have been inconsistent in previous studies (27, 28).

The β_2 -AR gene contains the genetic code for the receptor protein. Several variants in this gene have been described in literature (14). Three of the single-nucleotide polymorphisms, at nucleotides 46, 79, and 491, lead to amino acid substitution at codons 16, 27, and 164 in the coding region of the gene and might therefore have consequences for receptor function (29). When expressed in cells, the receptors with polymorphisms at codons 16 and 27 have been associated with differences in cellular β_2 -AR trafficking (30). The more rare polymorphism at codon 164 is associated with depressed functional coupling to Gs and reduced β_2 -adrenergic receptor sequestration (31). Large *et al.* (18) showed that isolated abdominal sc fat cells from women homozygous for the Arg16 polymorphism of the β_2 -AR had a 5-fold lower sensitivity for β_2 -AR agonist-induced lipolysis than fat cells from women heterozygous or homozygous for Gly16, independent of percent body fat of the women. More recently it was shown that fat cells from subjects with different ho-




FIG. 3. Plasma NEFA concentration (micromoles per liter), plasma glycerol concentration (micromoles per liter), plasma glucose concentration (millimoles per liter), and plasma lactate concentration (millimoles per liter) for the codon 16 polymorphisms in the β_2 -AR at baseline and during infusion of salbutamol. *, $P < 0.05$ vs. baseline; #, $P < 0.05$ Arg16Arg vs. Gly carriers.

mozygous haplotypes of the β_2 -AR gene differed about 250-fold in sensitivity to terbutaline-induced lipolysis (17). The least sensitive homozygous haplotype group in this study contained the Arg variant at codon 16 (17).

This study extends these findings and demonstrates a reduced β_2 -AR agonist-induced thermogenic response in individuals homozygous for the Arg16 polymorphism of the β_2 -AR gene. On the basis of changes in plasma NEFA and glycerol concentrations, we could not confirm the reduced lipolytic response to β_2 -AR stimulation in the Arg16Arg group, which has been reported by Large *et al.* (18) based on *in vitro* lipolysis. This might be related to the limitations of plasma NEFA concentration as a measure of lipolysis because not only lipolysis but also reesterification and NEFA oxidation affect plasma NEFA concentrations. The higher RER and plasma lactate concentrations at baseline as well as during salbutamol infusion in the Arg16Arg group suggest that their energy production relies more on carbohydrate oxidation, which would be compatible with a reduced lipolysis.

This study shows that the codon16 polymorphism has a significant effect on the salbutamol-induced thermogenic response, independent of the change in NEFAs. The explana-

tion for this effect is not directly apparent from our study. It could be that the β_2 -AR-stimulated muscle and liver glycogen breakdown is also affected by the polymorphism, thus resulting in differences in carbohydrate substrate availability and oxidation (4).

It is not surprising that the study revealed a correlation between the change in NEFAs and the thermogenic response because we and others have previously shown an increase in EE during the infusion of a triglyceride emulsion, which elevates plasma NEFA levels (28, 32–34), although this is not found in all studies (35, 36), which may be related to differences in patient populations and study design. In addition, part of the increase in EE induced by dobutamine, a β_1 -adrenergic agonist, can be blocked by inhibiting lipolysis with nicotinic acid (37).

Previous studies have suggested that the β_2 -AR is involved in the reduced β -AR-mediated increase in lipolysis (5, 38, 39), lipid oxidation, and thermogenesis (5) in obese compared with lean individuals. This study suggests that variations in the lipolytic response to β_2 -adrenergic stimulation are associated with variations in the thermogenic response. BMI and fat percentage were negatively correlated with Δ NEFA in our study ($P < 0.01$ and $P = 0.06$, respectively),




FIG. 4. Plasma norepinephrine (nanograms per liter) and epinephrine (nanograms per liter) concentrations for the codon 16 polymorphisms in the β_2 -AR at baseline and during infusion of salbutamol. *, $P < 0.05$ vs. baseline; †, $P < 0.05$ vs. baseline for all three groups.

which supports these earlier findings. Whether the association between these variations in β_2 -AR-mediated Δ NEFA and the variation in body fat are completely independent of the codon 16 polymorphism remains to be determined, but as discussed above, no association between the polymorphism and the response of plasma NEFAs or glycerol was found in this study. Other factors that might have influenced the lipolytic response are age, insulin sensitivity, α_2 -adrenergic receptor sensitivity and other polymorphisms in the β_2 -AR gene or other genes associated with lipolysis. Aging is known to reduce β -adrenergic sensitivity (40). The Arg16Arg group tended to be slightly, although not significantly ($P = 0.11$), younger than the other two groups. This difference in age would only tend to reduce the difference in responsiveness between the polymorphism groups. It is therefore unlikely that the differences in thermogenic sensitivity among groups are due to age differences. HOMA index as a measure for insulin sensitivity was different among polymorphism groups (Table 1) and correlated with Δ EE but did not significantly contribute to the multiple regression model. As to be expected, the HOMA index was also associated with fat mass.

Subjects carrying the Arg16Arg polymorphism were all

TABLE 3. Pearson correlations with Δ EE for the whole group ($n = 34$)

Variables	Correlation	P value
Weight	-0.1	0.572
Fat percentage	-0.122	0.491
Fat mass	-0.117	0.512
FFM	-0.05	0.777
BMI	-0.181	0.305
Basal [NEFA]	0.397	0.022
Basal [glycerol]	0.302	0.093
Basal [insulin]	-0.408	0.025
Δ RER	-0.392	0.022
Δ [NEFA]	0.508	0.003
Δ [glycerol]	0.435	0.013
Δ [insulin]	0.224	0.234
HOMA index	-0.425	0.019
Age	0.241	0.170

carriers of the Gln27Gln polymorphism of the β_2 -AR gene. Of the Gly16Gly carriers, one subject had the Gln27Gln polymorphism and seven had the Gln27Glu polymorphism. Subjects with the Arg16Gly polymorphism were either Gln27Gln ($n = 7$) or Gln27Glu ($n = 9$). Limiting the analysis to the Gln27 homozygotes ($n = 13$) revealed the same difference in thermogenic response between Gly16 carriers ($n = 8$) and Arg16Arg carriers ($n = 5$) as in the total group. It is therefore unlikely that the polymorphism at codon 27 was responsible for the differences in thermogenic response between the Arg16Arg subjects and the Gly16 carriers.

We conclude that subjects with the Arg16Arg polymorphism of the β_2 -AR have a reduced thermogenic response to β_2 -adrenergic stimulation. Although this relatively small study needs confirmation, the findings support a role for this polymorphism in the development and maintenance of overweight and obesity, as suggested by association studies (12, 17, 18, 41).

Acknowledgments

Received July 13, 2004. Accepted January 20, 2005.

Address all correspondence and requests for reprints to: J. M. Oomen, Department of Human Biology, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands. E-mail: j.oomen@hb.unimaas.nl. This work was supported by Grant NWO 98-10-001, The Netherlands.

References

- Chagnon YC, Rankinen T, Snyder EE, Weisnagel SJ, Perusse L, Bouchard C 2003 The human obesity gene map: the 2002 update. *Obes Res* 11:313–367
- van Baak MA 2001 The peripheral sympathetic nervous system in human obesity. *Obes Rev* 2:3–14
- Ukkola O, Bouchard C 2004 Role of candidate genes in the responses to long-term overfeeding: review of findings. *Obes Rev* 5:3–12
- Hoeks J, van Baak MA, Hesselink MK, Hul GB, Vidal H, Saris WH, Schrauwen P 2003 Effect of β_1 - and β_2 -adrenergic stimulation on energy expenditure, substrate oxidation, and UCP3 expression in humans. *Am J Physiol Endocrinol Metab* 285:E775–E782
- Schiffelers SL, Saris WH, Boomsma F, van Baak MA 2001 β_1 - and β_2 -Adrenoceptor-mediated thermogenesis and lipid utilization in obese and lean men. *J Clin Endocrinol Metab* 86:2191–2199
- Hellstrom L, Large V, Reynisdottir S, Wahrenberg H, Arner P 1999 The different effects of a Gln27Glu β_2 -adrenoceptor gene polymorphism on obesity in males and in females. *J Intern Med* 245:253–259

7. Ukkola O, Tremblay A, Bouchard C 2001 β -2 adrenergic receptor variants are associated with subcutaneous fat accumulation in response to long-term over-feeding. *Int J Obes Relat Metab Disord* 25:1604–1608
8. Meirhaeghe A, Helbecque N, Cottel D, Amouyel P 1999 β_2 -Adrenoceptor gene polymorphism, body weight, and physical activity. *Lancet* 353:896
9. Echwald SM, Sorensen TI, Tybjaerg-Hansen A, Andersen T, Pedersen O 1998 Gln27Glu variant of the human β_2 -adrenoceptor gene is not associated with early-onset obesity in Danish men. *Diabetes* 47:1657–1658
10. Kortner B, Wolf A, Wendt D, Beisiegel U, Evans D 1999 Lack of association between a human β_2 adrenoceptor gene polymorphism (Gln27Glu) and morbid obesity. *Int J Obes Relat Metab Disord* 23:1099–1100
11. Gonzalez Sanchez JL, Proenza AM, Martinez Larrad MT, Ramis JM, Fernandez Perez C, Palou A, Serrano Rios M 2003 The glutamine 27 glutamic acid polymorphism of the β_2 -adrenoceptor gene is associated with abdominal obesity and greater risk of impaired glucose tolerance in men but not in women: a population-based study in Spain. *Clin Endocrinol (Oxf)* 59:476–481
12. Garenc C, Perusse L, Chagnon YC, Rankinen T, Gagnon J, Borecki IB, Leon AS, Skinner JS, Wilmore JH, Rao DC, Bouchard C 2003 Effects of β_2 -adrenoceptor gene variants on adiposity: the HERITAGE Family Study. *Obes Res* 11:612–618
13. van Rossum CT, Hoebee B, Seidell JC, Bouchard C, van Baak MA, de Groot CP, Chagnon M, de Graaf C, Saris WH 2002 Genetic factors as predictors of weight gain in young adult Dutch men and women. *Int J Obes Relat Metab Disord* 26:517–528
14. Reihnsaus E, Innis M, MacIntyre N, Liggett SB 1993 Mutations in the gene encoding for the β_2 -adrenoceptor in normal and asthmatic subjects. *Am J Respir Cell Mol Biol* 8:334–339
15. Green SA, Turki J, Innis M, Liggett SB 1994 Amino-terminal polymorphisms of the human β_2 -adrenoceptor impart distinct agonist-promoted regulatory properties. *Biochemistry* 33:9414–9419
16. Small KM, McGraw DW, Liggett SB 2003 Pharmacology and physiology of human adrenoceptor polymorphisms. *Annu Rev Pharmacol Toxicol* 43:381–411
17. Eriksson P, Dahlman I, Ryden M, Hoffstedt J, Arner P 2004 Relationship between β_2 adrenoceptor gene haplotypes and adipocyte lipolysis in women. *Int J Obes Relat Metab Disord* 28:185–190
18. Large V, Hellstrom L, Reynisdottir S, Lonnqvist F, Eriksson P, Lannfelt L, Arner P 1997 Human β_2 adrenoceptor gene polymorphisms are highly frequent in obesity and associated with altered adipocyte β_2 adrenoceptor function. *J Clin Invest* 100:3005–3013
19. Siri WE 1956 The gross composition of the body. *Adv Biol Med Physiol* 4:239–280
20. Schoffelen PF, Westerterp KR, Saris WH, Ten Hoor F 1997 A dual-respiration chamber system with automated calibration. *J Appl Physiol* 83:2064–2072
21. Weir JB 1949 New methods for calculating metabolic rate with special reference to protein metabolism. *J Physiol* 109:1–9
22. Adriaens MP, Schoffelen PF, Westerterp KR 2003 Intra-individual variation of basal metabolic rate and the influence of daily habitual physical activity before testing. *Br J Nutr* 90:419–423
23. Peronnet F, Massicotte D 1991 Table of nonprotein respiratory quotient: an update. *Can J Sport Sci* 16:23–29
24. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC 1985 Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412–419
25. Alberts G, Boomsma F, Man in't Veld AJ, Schalekamp MA 1992 Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorometric detection. *J Chromatogr* 583:236–240
26. Ravussin E, Bogardus C 1989 Relationship of genetics, age, and physical fitness to daily energy expenditure and fuel utilization. *Am J Clin Nutr* 49:968–975
27. Schiffellers SL, van Harmelen VJ, de Grauw HA, Saris WH, van Baak MA 1999 Dobutamine as selective $\beta(1)$ -adrenoceptor agonist in *in vivo* studies on human thermogenesis and lipid utilization. *J Appl Physiol* 87:977–981
28. Schiffellers SL, Saris WH, van Baak MA 2001 The effect of an increased free fatty acid concentration on thermogenesis and substrate oxidation in obese and lean men. *Int J Obes Relat Metab Disord* 25:33–38
29. Liggett SB 2000 Pharmacogenetics of β_1 - and β_2 -adrenoceptor polymorphisms. *Pharmacology* 61:167–173
30. Green SA, Turki J, Bejarano P, Hall IP, Liggett SB 1995 Influence of β_2 -adrenoceptor genotypes on signal transduction in human airway smooth muscle cells. *Am J Respir Cell Mol Biol* 13:25–33
31. Green SA, Cole G, Jacinto M, Innis M, Liggett SB 1993 A polymorphism of the human β_2 -adrenoceptor within the fourth transmembrane domain alters ligand binding and functional properties of the receptor. *J Biol Chem* 268:23116–23121
32. Sobotka L, Zadak Z, Bures J, Pidrman V 1991 Influence of rapid amino acid and lipid emulsion administration on gas exchange and resting energy expenditure. *Nutrition* 7:200–203;discussion 203–204
33. Jung RT, Shetty PS, James WP 1980 Heparin, free fatty acids and an increased metabolic demand for oxygen. *Postgrad Med J* 56:330–332
34. Thiebaud D, Acheson K, Schutz Y, Felber JP, Golay A, Defronzo RA, Jequier E 1983 Stimulation of thermogenesis in men after combined glucose-long-chain triglyceride infusion. *Am J Clin Nutr* 37:603–611
35. Mascioli EA, Randall S, Porter KA, Kater G, Lopes S, Babayan VK, Blackburn GL, Bistrian BR 1991 Thermogenesis from intravenous medium-chain triglycerides. *J Parenter Enteral Nutr* 15:27–31
36. Kjekshus JK, Ellekjaer E, Rinde P 1980 The effect of free fatty acids on oxygen consumption in man: the free fatty acid hypothesis. *Scand J Clin Lab Invest* 40:63–70
37. Schiffellers SL, Brouwer EM, Saris WH, van Baak MA 1998 Inhibition of lipolysis reduces β_1 -adrenoceptor-mediated thermogenesis in man. *Metabolism* 47:1462–1467
38. Lonnqvist F, Wahrenberg H, Hellstrom L, Reynisdottir S, Arner P 1992 Lipolytic catecholamine resistance due to decreased β_2 -adrenoceptor expression in fat cells. *J Clin Invest* 90:2175–2186
39. Reynisdottir S, Wahrenberg H, Carlstrom K, Rossner S, Arner P 1994 Catecholamine resistance in fat cells of women with upper-body obesity due to decreased expression of β_2 -adrenoceptors. *Diabetologia* 37:428–435
40. Kerckhoffs DA, Blaak EE, Van Baak MA, Saris WH 1998 Effect of aging on β -adrenergically mediated thermogenesis in men. *Am J Physiol* 274:E1075–E1079
41. Meirhaeghe A, Helbecque N, Cottel D, Amouyel P 2000 Impact of polymorphisms of the human β_2 -adrenoceptor gene on obesity in a French population. *Int J Obes Relat Metab Disord* 24:382–387

JCEM is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.